





CLASS _____

BOOK _____



H. J. Putnam.







B 19

HISTORICAL
OF
PENNSYLVANIA
The
Vol. II


Digitized by the Internet Archive
in 2010 with funding from
Lyrasis Members and Sloan Foundation

FOR
THE
INDEX

HISTORY
of
QUADRUPEDS,

The Third Edition.

Vol. II.



L O N D O N :

Printed for B.&J. WHITE , Fleet Street .

M D C C X C I I I .

* *

QL

703

P4

1793

v.2

VIRGINIA

AUG 13 1982

STATE LIBRARY

HISTORY

OF

QUADRUPEDS.

Six cutting teeth, and two canine, in each jaw.

Five toes before; five behind.

In walking rests on the hind feet, as far as the heel.

XX. BEAR.

Urfus. Plinii lib. viii. c. 36.

Aext. Oppian Cyneg. iii. 139.

Urfus. Gejner quad. 941. Agricola, An.

Subter. 486. Raii syn. quad. 171.

Niedzwiedz. Rzeczynski Polon. 225.

*Bär. Klein quad. 82. Schwencfeld The-
riotroph. 131. Ridinger Wild. Thiere.*

31. Art. Zool. i. No. 20.

*Urfus niger, cauda concolore. Brisson
quad. 187.*

Urfus, cauda abrupta. Lin. syst. 69.

Biorn. Faun. suec. No. 19.

L'Ours. De Buffon, viii. 248. tab. xxxi.

*xxxii. Schreber, cxxxix. cxi. Lev.
Mus.*

208. BROWN.

B. with a long head: small eyes: short ears, rounded at the top: strong, thick, and clumsy limbs: very short tail: large feet: body covered with very long and shaggy hair, various in its color: the largest of a rusty brown: some from the confines of *Russia*, black, mixed with white hairs, called by the *Germans*, *silver bear*; and some (but rarely) are found in *Tartary* of a pure white.

VOL. II.

B

Inhabits

PLACE AND
MANNERS.

Inhabits the north parts of *Europe* and *Asia*, the *Alps* of *Switzerland*, and *Dauphiné*; *Arabia**, *Japan*†, and *Ceylon*‡; and the northern parts of *North America*; and extends to the *Andes* of *Peru*: Doctor *Shaw* informs us, it is also found in *Barbary*. They must have been very plentiful, for *Pliny* says that *Demetrius Aenobarbus* produced at one of the shews a hundred *Numidian* bears, and as many *Ethiopian* hunters§. The brown bears are sometimes carnivorous, and will destroy cattle, and eat carrion; but their general food is roots, fruits, and vegetables: will rob the fields of pease; and when they are ripe, pluck great quantities up; beat the pease out of the husks on some hard place, eat them, and carry off the straw: they will also, during winter, break into the farmer's yard, and make great havock among his flock of oats: are particularly fond of honey.

They live on berries, fruits, and pulse, of all kinds; and feed much on the black mulberry: are remarkably fond of potatoes, which they very readily dig up with their great paws: make great havock in the fields of *maiz*; and are great lovers of milk and honey. They feed much on herrings, which they catch in the season when those fish come in shoals up the creeks; which gives their flesh a disagreeable taste; and the same effect is observed when they eat the bitter berries of the *Tupelo*.

Bears strike with their fore foot like a cat; seldom or never use their mouths in fighting; but seizing the assailant with their paws, and pressing him against their breast, almost instantly squeeze him to death.

The females, after conception, retire into the most secret

* *Forsskal*, iv. † *Kämpfer*, *Hist. Japan*, i. 126. ‡ *Knox*, *Hist. Ceylon*. 20.

§ *Lib. viii. c. 36.*

places; least, when they bring forth, the males should devour the young: it is affirmed for fact, that out of the several hundred bears that are killed in *America*, during winter, (which is their breeding season) that scarcely a female is found among * them; so impenetrable is their retreat during their pregnancy: they bring two, rarely three, young at a time: the cubs are deformed, but not a shapeless mass, to be licked into shape, as the ancients pretended†. The cubs even of the brown bears are of a jetty blackness, and often have round their necks a circle of white. The flesh of a bear in autumn, when they are most excessively fat, by feeding on acorns, and other mast, is most delicate food; and that of the cubs still finer; but the paws of the old bears are reckoned the most exquisite morsel: the fat white, and very sweet: the oil excellent for strains, and old pains.

The latter end of autumn, after they have fattened themselves to the greatest degree, the bears withdraw to their dens, where they continue for a great number of days in total inactivity, and abstinence from food, having no other nourishment than what they get by sucking their feet, where the fat lodges in great abundance. In *Lapland* they pass the long night in dens lined warmly with a vast bed of moss, in which they roll themselves, secure from the cold of the severe season‡. Their retreats are either in cliffs of rocks; in the deepest recesses of the thickest woods; or in the hollows of ancient trees, which they ascend and descend with surprizing agility: as they lay in no winter provisions, they

* Out of 500 bears that were killed in one winter, in two counties of *Virginia*, only two females were found, and those not pregnant. *Lawson*, 117.

† *Hi sunt candida informisque caro, paulo majoribus major, sine oculis, sine pilo; ungues tantum prominent: hanc lambendo parlatim figurant.* *Plinii lib. viii. c. 36.*

‡ *Fl. Lap. 313.* The moss is a variety of the *Polytrichum Commune*.

are in a certain space of time forced from their retreats by hunger, and come out extremely lean: multitudes are killed annually in *America*, for the sake of their flesh, or skins, which last makes a considerable article of commerce.

209. BLACK.

Arch. Zool. 2d. Ed. ii. N° 19.


B. with a long pointed nose, and narrow forehead: the cheeks, and throat of a yellowish brown color: hair over the whole body and limbs of a glossy black, smoother and shorter than that of the *European* kind.

They are usually smaller than those of the old world: yet Mr. *Bartram* gives an instance of an old he-bear killed in *Florida*, which was seven feet long, and, as he guessed, weighed four hundred pounds.

These animals are found in all parts of *North America*, from *Hudson's Bay* to the southern extremity; but in *Louisiana* and the southern parts they appear only in the winter, migrating from the north in search of food. They spread across the northern part of the *American* continent to the *Asiatic* isles. They are found in the *Kurilski* islands, which intervene between *Kamtshatka* and *Japan*, *Jeso*, *Masima*, which lies north of *Japan*, and probably *Japan* itself; for *Kämpfer* says, that a few small bears are found in the northern provinces.

It is very certain that this species of bear feeds on vegetables. *Du Rots*, who is a faithful as well as intelligent writer, relates, that in one severe winter, when these animals were forced in multitudes from the woods, where there was abundance of animal food, they rejected that, notwithstanding they were ready to perish with hunger, and migrating into the lower *Louisiana*, would often break





Magell Sculp

Polar Bear— . S. 210.

into the courts of houses. They never touched the butchers meat which lay in their way, but fed voraciously on the corn or roots they met with.

White bear. *Martin's Spitzberg.* 100. *App.* xxvi. *Arch. Zool.* i. No. 18.
Egede Greenl. 59 *Ellis voy.* 41. *Grantz* *Ursus albus. Martensii. Klein quad.* 82.
Greenl. i. 73. *Barentz voy.* 18. 45. *La* *L'Ours blanc. Brisson quad.* 188. *De Buf-*
Houtan voy. i. 235. *Catfish Carolina,* *fon,* xv. 128. *Schreber,* cxli. *Lev. Mus.*

210. POLAR.

B. with long head and neck: short round ears: end of the nose black: vast teeth; hair long, soft, white, tinged in some parts with yellow: limbs of great size and strength: grows to a vast size: the skins of some are thirteen feet long.

This animal is confined to the coldest part of the globe: it has been found as far as navigators have penetrated northwards, above *lat.* 80. The frigid climates only seem adapted to its nature. It is unknown, except on the shores of *Hudson's Bay, Greenland,* and *Spitzbergen.* The north of *Norway,* and the country of *Mesen,* in the north of *Russia,* are destitute of them: but they are met with again in great abundance in *Nova Zembla,* and from the river *Ob,* along the *Siberian* coast, to the mouths of the *Tenesi,* and *Lena,* but are never seen far inland, unless they lose their way in mists; none are found in *Kamtshatka,* or its islands.

PLACE.

They have been seen as far south as *Newfoundland;* but they are not natives of that country, being only brought there accidentally on the islands of ice.

During summer the white bears are either resident on islands of ice, or passing from one to another: they swim admirably, and can

MANNERS.

can continue that exercise* six or seven leagues; and dive with great agility. They bring two young at a time: the affection between the parents and them is so strong, that they would die rather than desert one another. Their winter retreats are under the snow†, in which they form deep dens, supported by pillars of the same, or else under some great eminence beneath the fixed ice of the frozen sea.

They feed on fish, seals, and the carcases of whales; and on human bodies, which they will greedily disinter: they seem very fond of human blood; and are so fearless as to attack companies of armed men, and even to board small vessels. When on land, they live on birds, and their eggs; and, allured by the scent of the seals flesh, often break into and plunder the houses of the *Greenlanders*: their greatest enemy in the brute creation is the *Morse*‡, with whom they have terrible conflicts, but are generally worsted; the vast teeth of the former giving it a superiority.

The flesh is white, and said to taste like mutton: the fat is melted for train oil, and that of the feet used in medicine; but the liver is very unwholesome, as three of *Barentz's* sailors experienced, who fell dangerously ill on eating some of it boiled.

One of this species was brought over to *England*, a few years ago: it was very furious, almost always in motion, roared loud, and seemed very uneasy, except when cooled by having pail-fulls of water poured on it.

Callixenus Rhodius§, in his description of the pompous pro-

* *La Hontan*, i. † *Egede*, 60. ‡ *Egede*, *Greenl.* 60. 83.

§ As quoted by *Athenæus*, lib. v. p. 201.

cession of *Ptolemaus Philadelphus* at *Alexandria*, speaks of one great white Bear, ἄρκτος λευκή μεγάλη μία, among other wild beasts that graced the shew: notwithstanding the local situation of this species at present, it is possible that *Ptolemy* might procure one; whether men could penetrate, in those early times, as far as the present residence of these *Arctic* animals, I will not venture to affirm, nor to deny; but since my friend, the Hon. *Daines Barrington* *, has clearly proved the intense cold that in former ages raged in countries now more than temperate, it is most probable that in those times they were stocked with animals natural to a rigorous climæ; which, since the alteration, have necessarily become extinct in those parts: the *Polar* bear might have been one; but that it was the species meant by *Callixenus* is clear to me, by the epithet μεγάλη, or great, which is very applicable to it; for the white *Tartarian* land bear (which *Ptolemy* might very easily procure) differs not in size from the black or brown kind, but the bulk of the other is quite characteristic.

Land bears, sometimes spotted with white; at other times wholly white; are sometimes observed on the parts of *Russia* bordering on *Siberia*, in a wandering state, supposed to have strayed out of the lofty snowy mountains, which divide the two countries †.

PIED
LAND BEARS.

* *Phil. Trans.* vol. lviii. p. 58.

† *Doctor Pallas.*

211. WOLVERENE.

Quickhatch *Catesby Carolina, App. xxx.*
Carcajou, or Quickhatch. *Dobbs Hud-*
son's B. y. 40.

Quickhatch, or Wolverine. *Ellis Hud-*
son's Trav. 42. *Clerk's v.* ii. 3.
Edw. 103.

Ursus luscus. U. cauda elongata, cor-
pore ferrugineo, rostro fusco, fronte

plagaue laterali corporis. *Lin. syst.*
71. *Art. Zool.* 1. N^o 21.

Ursus. *Freti Hudsonis.* U. castanei co-
loris, cauda unicolore, rostro pedi-
busque fuscis. *Briffon quad.* 188.
Schreber, cxliv.

Le Glouton. *De Buffon, Supplem.* iii.
244. *LEV. Mus.*

B with a black sharp-pointed visage: short rounded ears, al-
most hid in the hair: hairs on the head, back, and belly,
reddish, with black tips, so that those parts appear, on first
sight, quite black: sides of a yellowish brown, which passes in
form of a band quite over the hind part of the back, above the
tail: on the throat a white spot: on the breast a white mark, in
form of a crescent: legs very strong, thick and short, of a deep
black: five toes on each foot*, not deeply divided: on the fore
foot of that I examined were some white spots: the bottom of
the feet covered very thickly with hair: rests, like the bear, on its
foot, as far as the first joint of the leg; and walks with its back
greatly arched: claws strong and sharp, white at their ends: tail
cloathed with long coarse hairs; those at the base reddish, at the
end black: some of the hairs are six inches long: length from nose

* Mr. *Edwards* observed only four toes on the fore feet of the animal he de-
scribes. My description is taken from an entire skin, in very fine preservation,
communicated to me by the late Mr. *Astton Blackburne*, of *Orford, Lancashire*,
who, with indefatigable industry and great judgment, enriched the cabinets of
his friends with the rarest natural productions of that continent: as this work
has profited so greatly by that gentleman's labors, it would be ungrateful to
omit my acknowledgements.



Wolverine — . 8. 21.



to tail twenty-eight inches: length of the trunk of the tail seven inches, but the hairs reach six beyond its end: the tail in Mr. *Edwards's* figure not quite accurate: it is corrected in that which is borrowed from his admirable work. The whole body is covered with very long and thick hair, which varies in color, according to the season.

Inhabits *Hudson's-Bay*, and *Canada*, as far as the straits of *Michilimakinac*.

A most voracious animal: slow of foot, so is obliged to take its prey by surprize: in *America* is called the *Beaver-Eater*, for it watches those animals as they come out of their houses, and sometimes breaks into their habitations, and devours them.

In a wild state is vastly fierce; a terror to both wolf and bear, which will not prey on it when they find it dead*, perhaps on account of its being so very fœtid, smelling like a pole-cat: makes a strong resistance when attacked; will tear the flock from the gun, and pull the traps it is caught in to pieces: burrows†, and has its den under ground. Mr *Graham*, long resident in *Hudson's Bay*, has assured me, that it will lurk on a tree, and drop on the deer which pass beneath, and fasten on them till the animals are quite exhausted.

Charlevoix, in *Hist. Nouv. France*, v. 189, gives the name of this animal (*Carcajou*) to our 189th species, the *Puma*, or Brown Panther of *N. America*.

* *Clerk California*, ii. 3.

† *La Hontan's voy.* i. 62.

In conformity to the opinion of that respectable naturalist Doctor *Pallas*, I unite the *Wookwerene* and *Glutton*. I do not alter my description of the latter; but add both that and the synonyms; submitting to future times the propriety or impropriety of uniting these animals: there being distinctions that even now leave me very undetermined.

212. GLUTTON. Gulo. Olaii Magni gent. Septentr. 138. Hyæna. Brisson quad. 169. Ysbrandts Ides
Gulo, vieltrafs. Gesner quad. 554. Klein Trav. Harris's Coll. ii. 923.
quad. 83. tab. v. Mustela gulo M. pedibus fissis, corpore
Rosomak. Rzaczynski Polon. 218. Bell's rufo-tufco, medio dorfi nigro. Lin. syst.
Travels, i. 235. 67. Zimmerman. 311.
Muller's Rufs Samlung. iii. 549, 550. Jarf, Filfrefs. Faun. suec. No. 14.
Ritchkoff Topogr. Orenb. i. 295. Jærvæn. Gunner's Aft. Nidros. iii. 143.
Jerf, Fieldfrofs. Strom Sammor. 152. Pon- tab. iii.
top. Norway, ii. 22. Scheffer's Lapland, Le Glutton. De Buffon, xiii. 278.
134.

B. with a round head: thick blunt nose: short ears, rounded, except at the tip: limbs large: back strait; marked the whole length with a tawny line: tail short and very full of hair: the hair in all other parts black, finely damasked or watered like a silk, and very glossy; but sometimes varies into a browner color. *Klein* attributes to it five toes on each foot: that which Mr. *Zimmerman* describes, had but four, very thickly covered with hair.

SIZE. The length of one which was brought from *Siberia*, and kept alive at *Dresden*, was a yard and eight inches: the height from the top of the head was nineteen inches. Mr. *Zimmerman* de-

scribes another, rather lesser than the former, which was shot near *Helmstedt*, in *Wolfenbuttle*. Its length was three feet three: its height before fifteen inches; behind, sixteen: the tail six inches.

Inhabits *Lapland*, the northern and eastern parts of *Siberia*, and *Kamtchatka*. Those of *Kamtchatka* differ, and vary to white and yellowish, and their skins are esteemed by the natives before the black: they say, that the heavenly beings wear no other garments. The women wear the paws of the white sort in their hair: and esteem a skin as the most valuable present which their husbands or lovers can make.

PLACE.

They are excessively voracious; that which was confined at *Dresden* would eat thirteen pounds of flesh in a day, and not be satisfied. The report of their filling themselves so full, as to be obliged to go between two trees to force out part of the food, seems to be fabulous.

Like the Lynx, it lurks on the boughs of trees, and will fall on any animal which passes by, fasten on, and destroy it. Its game is chiefly deer; and about the *Lena*, horses. Is capable of being made tame.

It differs from the bear by its lean habit; by not lying inactive in winter; and by its living entirely on animal food. It is also more bold, voracious, and cunning.

The *Russians* call it *Rosomak*; the *Kamtchatkans*, *Timmi*; and the *Koratfski*, *Haepi*. An animal, called by the *Greenlanders*, *Amanki*, is said to be found in their country, which is supposed to be the Glutton; but as *Greenland* is destitute of wood, I suppose their *Amanki*, or *Amarok*, to be a fabulous animal*.

* See *Crantz Hist. Greenland.*

213. RACCOON.

Raccoon. *Lawson Carolina*, 121. *Catef-*
by Carolina, App. xxix.

Mapach, seu animal cuncta prætentante
manibus. *Hernandez, Nov. Hisp.* 1.
Nieremberg, 175.

Vulpi affinis Americana. *Raii syn. quad.*
179. *Sloane Jamaica*, ii. 329.

Coati. *Worm. Mus.* 319.

Coati. *Urfus cauda annulata* varie-

grata. *Briffon quad.* 189.

Urfus Lotor. U. cauda annulata, fascia
per oculos transversali nigra. Lin.
ssyl. 70. *Arch. Zool.* i. N^o 22.

Le Raton. *De Buffon*, viii. 337. *tab.*
xliii. *Schreber*, cxliiii.

Raccoon. *Kalm's Travels. Forster's Tr.*
i. 96. 208. *tab.* 11. *Lev. Mus.*

B. with a sharp-pointed black nose: upper jaw the longer: ears short, and rounded: eyes surrounded with two broad patches of black: from the forehead to the nose a dusky line: face, cheeks, and chin, white: upper part of the body covered with hair, ash-colored at the root, whitish in the middle, and tipped with black: tail very bushy, annulated with black: toes black, and quite divided. Sometimes this animal varies: I have seen one entirely of cream color*.

PLACE.

Inhabits the warm and temperate parts of *America*: found also in the mountains of *Jamaica*, and in the isles of *Maria*, between the S. point of *California* and *Cape Corientes*, in the S. Sea†: an animal easily made tame; very good-natured and sportive, but as unlucky as a monkey; almost always in motion; very inquisitive, examining every thing with its paws; makes use of them as hands: sits up to eat: is extremely fond of sweet things, and strong liquors, and will get excessively drunk: has all the cunning of a fox: very destructive to poultry; but will eat all sorts

* *Lev. Mus.*† *Dampier's voy.* i. 276.

of fruits, green corn, &c.: at low water feeds much on oysters; will watch their opening, and with its paw snatch out the fish; sometimes is caught in the shell, and kept there till drowned by the coming in of the tide: fond also of crabs: climbs very nimbly up trees: hunted for its skin; the fur next to that of the beaver, being excellent for making hats.

Wha Tapoua Row. *White's Bot. Bay*, 278.

214. NEW
HOLLAND.

B. of the same external form as the *American Raccoon* except the ears, which are pointed: six cutting teeth in the upper jaw; two ? in the lower: back of a dark grey; growing lighter on the sides: belly of a fine brown: tail as long as the body, covered with long hair; the lower part near the end is naked, and has a prehensile quality like some species of monkeys, or the common *Opossum*.

Inhabits *New Holland*.

PLACE.

XXI. BADGER.

Six cutting teeth, two canine, in each jaw.

Five toes before, five behind: very long strait claws on the fore feet.

A tranverse orifice between the tail and the anus.

215. COMMON.

Meles. *Plinii lib. viii. c. 38. Gesner quad. 327.*

Meles, five Taxus. *Raii syn. quad. 185.*

Meles, Taxus, Tassus, Blerellus; Jazwicz, Borsuk. *Rzacinski Polon. 233.*

Coati cauda brevi, Coati griseus, Taxus, Meles, Tax. *Klein quad. 73.*

Dachis. *Kramer Austr. 313.*

Meles pilis ex fordide albo et nigro variegatis vestita, capite tæniis alternatim albis et nigris variegato. *Briffon*

quad. 183.

Le Blaireau, ou Taifon. *De Buffon, viii. 104. tab. vii.*

Urfus meles U. cauda concolore, corpore supra cinereo, subtus nigro, fascia longitudinali per oculos auresque nigra. *Lin. syst. 70.*

Meles unguibus anticis longissimis. *Graf-fuin. Faun. suec. No. 20. Br. Zool. i. 64. Br. Zool. illustr. tab. lii. Schreber, cxlii. Lev. Mus.*

B. with small eyes: short rounded ears: short thick neck: with nose, chin, lower sides of the cheeks, and middle of the forehead, white: ears and eyes inclosed in a pyramidal bed of black: hairs on the body long and rude; their bottoms a yellowish white, middle black, ends ash-colored: throat, breast, belly, and legs black: tail covered with long hairs, colored like those on the body: legs very short and thick: claws on the fore feet very long: a foetid white matter exudes from the orifice beneath the tail: animal of a very clumsy make.

SIZE.

The length is commonly two feet six inches from the nose to the origin of the tail; of the tail six inches: the weight from fifteen to thirty four pounds. The last is rare; but I met with, in the winter of 1779, a male of that weight.

Inhabits

Inhabits most parts of *Europe*, as far north as *Norway**, and *Russia*; and the *step* or desert beyond *Orenburgh*, in the *Russian Asiatic* dominions; in *Great Tartary*, and in *Siberia* about the river *Tom*, and even about the *Lena*, but none in the north; inhabits also *China*, and is often found in the butchers shops in *Pekin*, the *Chinese* being fond of them†. A scarce animal in most countries: seldom appears in the day; confines itself much to its hole: is indolent and sleepy: generally very fat: feeds by night; eats roots, fruits, grubs, insects, and frogs: not carnivorous: its flesh makes good bacon: runs slowly; when overtaken comes to bay, and defends itself vigorously: its bite hard and dangerous: burrows under ground, makes several apartments, but forms only one entrance from the surface: hunted during night, for the skin, which serves for pistol furniture; the hair, for making brushes to soften the shades in painting. The division of this species into two, *viz.* the swine and the dog badger, unnecessary, there being only one.

Art. Zool. i. No. 23.

β. AMERICAN.

B. with a white line from the tip of the nose, passing between the ears, to the beginning of the back, bounded on each side, as far as the hind part of the head, with black, then by a white one, and immediately between that and the ears is another of black: hair long: back colored like that of the common badger: sides yellowish: belly cinereous: thighs dusky: tail covered with long dirty yellow hairs, tipped with white; the end dusky.

* *Pontop. hist. Norway*, ii. 28.

† *Bell's travels*, ii. 83.

The

The legs were wanting in the skin I took my description from. M. de Buffon's description, taken from a stuffed animal* brought from *Terra di Librador*, will supply that defect: he says there were only four toes on the fore feet; but he suspects (as I imagine was the case) that the fifth might have been rubbed off in stuffing.

Described from a skin from *Hudson's-Bay*, found in a furrier's shop in *London*: it was less than that of the *European* badger: the furrier said, he never met with one before from that country. *Kalm*† says, he saw the *European* badger in the province of *Pennsylvania*, where it is called the *Ground Hog*‡; and this proves to be no other, varying very little from it.

216. INDIAN.

B. with a small head, and pointed nose: scarcely any external ears; only a small prominent rim round the orifice, which was oval: color of the nose and face, a little beyond the eyes, black: crown, upper part of the neck, the back, and upper part of the tail, white, inclining to grey: legs, thighs, breast, belly, sides, and under part of the tail black.

Five toes on each foot; the inner small: claws very long and strait.

SIZE:

Length from nose to tail about two feet: tail four inches: hair short and smooth.

* He calls it *Le Carcejou*. Suppl. iii. 242. tab. xlix.

† *Kalm's travels, Forster's transl.* i. 189.

‡ M. *Briffon* describes a white Badger, with a yellowish white belly, and also much inferior in size to that of *Europe*, which M. *Reaumur* received from *New York*. Vide *Eriffon quad.* 185.

B A D G E R.

17

PLACE.

Inhabits *India*: feeds on flesh: is playful, lively, and good-natured: sleeps rolled up, with its head between its hind legs; sleeps little in the day: refused all commerce with the *English* badger which was turned to it, and lived some time in the same place: climbs very readily over a division in its cage.

XXII. OPOSSUM.


Two canine teeth in each jaw.
 Cutting teeth unequal in number in each jaw*.
 Five toes on each foot: hind feet formed like a hand, with a distinct thumb.
 Tail very long, slender, and usually naked.

217. VIRGINIAN. Tlaquatzin. *Hernandez Mex.* 330. *Nieremberg*, p. 136. and fig. 136. 135. *Catesby Carolina, App.* xxix. *Roche-
fort Atlas*, i. 283.
 Tajibi. *Margrave Brasil*, 222. *Taii syn.* Fara, ou Ravall? *Gumilla, Orenoque*, iii.
quad. 182. 185. 238. *Arch. Zool.* i. No. 24.
 Semi-vulpa. *Cesner quad.* 870. *Ken. An.* Le Manicou. *Feville obs. Peru*, iii. 206.
 90. Wood-rat. *Du Ratz Louisiana*, ii. 65.
 Opossum. *Pb. Tr. alridg* ii. 884. *tab.* Didelphis marsupialis. D. mammis octo
 xiii; iii. 593; and v. 169. 177. *Larv-* intra abdomen? *Liv syl.* 71. *Amæn.*
for Carolina, 120. *Beverley's Virginia,* *Acad.*? i. 561. *LEV. Mus.*

O with long sharp-pointed nose: large, round, naked, and very thin ears, black, edged with pure white: small, black, lively eyes: long stiff hairs each side the nose, and behind the eyes: face covered with short soft white hairs: space round the eyes dusky: neck very short; its sides of a dirty yellow: hind part of the neck and the back covered with hair above two inches long; soft, but uneven; the bottoms of a yellowish white, middle part black, ends whitish: sides covered with dirty and dusky hairs; belly, with soft, woolly, dirty white hair: legs and thighs black: feet dusky: claws white: base of the tail clothed with long hairs, like those on the back; rest of the tail covered

* This species has eight cutting teeth in each jaw. *Tygn.*

with



Virginian Opossum — N. 217.



with small scales; the half next the body black, the rest white: it has a disagreeable appearance, looking like the body of a snake, and has the same prehensile quality as that of some monkeys: body round, and very thick: legs short: on the lower part of the belly of the female is a large pouch, in which the teats are lodged, and where the young shelter as soon as they are born.

The usual length of the animal is, from the tip of the nose to the base of the tail, about twenty inches; of the tail twelve inches.

Inhabits *Virginia, Louisiana, Mexico, Brazil, and Peru*: is very destructive to poultry, and sucks the blood without eating the flesh: feeds also on roots and wild fruits: is very active in climbing trees: will hang suspended from the branches by its tail, and, by swinging its body, fling itself among the boughs of the neighbouring trees: continues frequently hanging with its head downwards: hunts eagerly after birds and their nests: walks very slow: when pursued and overtaken, will feign itself dead: not easily killed, being as tenacious of life as a cat: when the female is about to bring forth, she makes a thick nest of dry grass in some close bush at the foot of a tree, and brings four, five, or six young at a time.

As soon as the young are brought forth, they take shelter in the pouch, or false belly, and fasten so closely to the teats, as not to be separated without difficulty: they are blind, naked, and very small when new-born, and resemble *fetuses*: it is therefore necessary that they should continue there till they attain a perfect shape, strength, sight, and hair; and are prepared to undergo what may be called a second birth: after which, they run into this pouch as into an asylum, in time of danger; and the parent carries them about with her. During the time of this second

SIZE.

PLACE.

MANNERS.

FALSE BELLY.

gestation, the female shews an excessive attachment to her young, and will suffer any torture rather than permit this receptacle to be opened; for she has power of opening or closing it by the assistance of some very strong muscles.

The flesh of the old animals is very good, like that of a sucking pig: the hair is dyed by the *Indian* women, and wove into garters and girdles: the skin is very foetid.

M. de *Euffon* seems not to be acquainted with this animal, but has compiled an account of its manners, and collected the synonyms of it. The figures * which he has given belong to the following species, as does the description.

218. *MOLUCCA.*Carigue, ou Saragoy. *De Laet*, 485.Carigueya. *Maregrave*, 223.Mus Marfupialis, *Beutel ratze. Klein*
quail, 59.Vulpes major putoria cauda tereti & glabra? *Barroet France Equin*, 166.Philander orientalis fœmina. *Sch. Mus.*

i. 61. tab. xxxvi. fig. 1. 2. xxxviii.

fig. 1.

Sarigue, ou l'Opossum. *De B ffor*, 311.x. tab. lkv. lxvi. *Schreber*, cxlvi. A.B. *LEV. MUS.*

○ with long, oval, and naked ears: mouth very wide: over each eye is an oblong spot of white: lower side of the upper jaw, throat, and belly, of a whitish ash-color: rest of the hair of a cinereous brown, tipped with tawny, darkest on the back: tail long as the body; near the base covered with hair, the rest naked: claws hooked.

On the belly of the female is a pouch, in which the young (like those of the former) shelter. *Maregrave* found six young within the pouch of the *Carigueya*, which I consider as the same animal. It had ten cutting teeth above, and eight below.

* The figure in the first edition was very indifferent, I have therefore changed it for the very faithful one in the *Phil. Transf.*

Length from nose to tail, ten inches. The tail exceeds the length of head and body. Its whole figure is of a much more slender and elegant make than the former.

Size.

The tail pulverised, and taken in a glass of water, is reckoned in *New Spain* a sovereign remedy against the gravel, colic, and several other disorders.

This genus is not confined to *America*, as *M. de Buffon* supposes; who combats the opinion of other naturalists on this subject with much warmth: but the authority of *Pisô*, *Valentyn*, and of *Le Bruyn*, who have seen it both in *Java* and in the *Molucca Isles*, and of numbers of collectors in *Holland*, who receive it frequently from those places. This and N^o 219 are proofs of what I advance. It is also met with in *New Holland*.

Place.

This species is found in great numbers in *Aroe* and *Solor*: It is called in the *Indies*, *Pelandor Aroe*, or the *Aroe Rabbet*. They are reckoned very delicate eating; and are very common at the tables of the Great, who rear the young in the same places in which they keep their rabbits. It inhabits also *Surinam*, and the hot parts of *America*.

Seba figures and describes, in his 1st vol. 64. tab. xxxix. an *Opossum* under the name of *Philander maximus orientalis fœmina*. It has a pouch like the former: is much larger: seems to have a longer and more slender tail: has broader ears; has a dusky spot over each eye, and is of a darker color. It feeds on fruits: was brought from *Amboina*, where it is called *Coes Coes*.*

α. GREATER.

* *In Indiis orientalibus, idque solum, quantum hæterus constat, in Amboina, similis Bestia (Carigueya) frequens ad felis magnitudinem accedens, maculata ab incolis comeditur, si rite preparetur, nam alias facit. Nomen illi Cous Cous inditum. Pifô India, 323.*

I am

I am unacquainted with this species, so leave these two conjoined till I receive fuller information.

Much is wanted to complete the natural history of this genus.

219. JAVAN.

Filander. *Le Bruyn voy. East Indies*, ii. 101. tab. ccxiii. Ed. Angl.

O. (according to *Le Bruyn's* figure) with a narrow fox-like head: upright pointed ears: a brown stripe passing through the eyes: fore legs very short: five toes on the fore feet; three only on the hind, two of which are very strong; the outmost slender and weak; and found on dissection to consist internally of two bones, closely united, with two weak claws bursting out of the skin*: tail thick, shorter than the body.

In the upper jaw are six cutting teeth; in the lower two, which are formed like those of squirrels: no canine teeth†.

On the belly is a complete pouch, like the *Virginian* kind: hair on the body rude: face seemingly that of a hare.

PLACE.

Discovered first by Mr. *Le Bruyn*, who saw in *Java* several in an inclosure along with rabbits: they burrowed like them; leaped in their pace; preserved their young in the pouch, which would often peep out when the old ones were still.

The fidelity of *Le Bruyn's* figure has been since confirmed by the specimens sent from *Java* into *Holland*.

* *Pallas* in act. acad. *Petrop. pars* ii. 229. tab. ix*.

† The same.

- Mus sylvestris Americanus Scalpes dictus.* *Seb. Myf. i. 46. tab. xxxi. fig. 1. 2.*
Philander saturate spadiceus in dorso, in ventre dilatè flavus, pedibus albi-
cantibus. *Briffon quad. 211.*
Didelphis murina. D. cauda semipilosa, mammis fenis. *Lin. syst. 72.*
La Marmose. De Buffon, x. 336. tab. lii. liii. Schreber, cxlix.

220. MURINE.

O with long broad ears, rounded at the end, thin and naked: eyes encompassed with black: face, head, and upper part of the body, of a tawny color: the belly yellowish white: the feet covered with short whitish hair: toes formed like those of the *Virginian*: tail slender, covered with minute scales, from the tip to within two inches of the base, which are cloathed with hair. Length from nose to tail, about eight inches; tail of the same length: the female wants the false belly of the former; but, on the lower part, the skin forms on each side a fold, between which the teats are lodged.

This species varies in color: I have seen one from *Guiana*, brown above, white beneath.

Inhabits the hot parts of *South America*: agrees with the others in its food, manners, and the prehensile powers of its tail: it brings from ten to fourteen young at a time; at first, in some species, there are that number of teats: the young affix themselves to the teats as soon as they are born, and remain attached, like so many inanimate things, 'till they attain growth and vigor to shift a little for themselves.

221. MEXICAN.

Cayopollin. *Hernandez Nov. Hisp.* 10.Animal caudimanum. *Nieremberg*, 158.*Mus Africanus* Kayopollin dictus, mas.*Seb. Mus.* tab. xxxi. fig. 3.

Philander saturatè spadiceus in dorso, in

ventre ex albo flavicans, cauda ex saturatè spadiceo maculata. *Briffon quad.* 212. *Schreber*, cxlviii.Le Cayopollin. *De Buffon*, x. 350. tab. lv. *LEV. MUS.*

O with large, angular, naked, and transparent ears: nose thicker than that of the former kind: whiskers very large: a slight border of black furrounds the eyes: face of a dirty white, with a dark line running down the middle: the hairs on the head, and upper part of the body, ash-colored at the roots; of a deep tawny brown at the tips: legs dusky: claws white: belly dull cinereous: tail long, and pretty thick, varied with brown and yellow: is hairy near an inch from its origin; the rest naked: length, from nose to tail, about nine inches; the tail the length of the body and head.

Inhabits the mountains of *Mexico*: lives in trees, where it brings forth its young: when in any fright, they embrace their parent closely: the tail is prehensile, and serves instead of a hand.

222. CAYENNE.

Le Crabier. *De Buffon, Supplem.* iii. 272.*Canis ferus major, Cancofus* vulgo dictus. *Koupara. Barrere France Equinoæ.* 149.

O with a long slender face: ears erect, pointed, and short: the coat woolly, mixed with very coarse hairs, three inches long, of a dirty white from the roots to the middle; from thence to the ends of a deep brown: sides and belly of a pale yellow: legs of a dusky brown: thumb on each foot distinct: on the toes

of

of the fore feet, and thumb of the hind, are nails; on the toes of the hind feet crooked claws: tail very long, taper, naked, and scaly.

Length seventeen *French* inches: of the tail fifteen and a half. The subject measured was young.

Inhabits *Cayenne*: very active in climbing trees, on which it lives the whole day. In marshy places, feeds on crabs, which, when it cannot draw out of their holes with its feet, hooks them by means of its long tail. If the crab pinches its tail, the animal sets up a loud cry, which may be heard afar: its common voice is a grunt like a young pig. It is well furnished with teeth, and will defend itself stoutly against dogs: brings forth four or five young, which it secures in some hollow tree. The natives eat these animals, and say their flesh resembles a hare. They are easily tamed, and will then refuse no kind of food.

O with the upper part of the head, and the back and sides, covered with long, soft, glossy hairs, of a dark cinereous color at the bottoms, and of a rusty brown towards the ends: belly of a dirty white.

Tail taper, covered with short brown hairs, except for four inches and a half of the end, which was white, and naked underneath: toes like the former.

The skin I examined had lost part of the face: the length from the head to the tail was thirteen inches: the tail the same.

This was found near *Endeavour* river, on the eastern coast of

VOL. II

E

New

223. New-York

New Holland, with two young ones *. It lodges in the grafts, but is not common.

224. VULPINE.

Stockdale's Det. Bay, 150.

O. with very long whiskers: ears erect, and pointed: upper parts of the body greyish, mixed with dusky and white hairs, tinged with rufous: the last predominates about the shoulders: all the under side of the neck and body of a tawny buff: about a quarter of the tail, next to the body, of the same color with the back; the rest black: length from the tip of the nose to the tail, two feet two inches: the tail fifteen.

Inhabits *New Holland*.

225. SHORT-
TAILED.

Mus sylvestris Americana, fœmina. *Seb.*

Mus. i. 50. tab. xxxi.

Philander obscure rufus in dorso, in ven-

tre helvus, cauda brevi crassa, *Briffon*
quad. 213. *Schreber*, cli.

O. with naked ears: the back of a dull red; belly of a paler: tail scarce half the length of the body; thick at the base, lessening towards the end: no false belly.

Inhabits *South America*: the young adhere to the teats as soon as born. *Seba* says it lives in woods, and brings from nine to twelve young at a time.

* *Cook's voy.* iii 586.

Philander ex rufo luteus in dorso, in ventre
ex flavo albicans, capite crasso. *Briffon*
quad. 213. *S. b. Mus.* i. 50. *tab.* xxxi.

fig. 8. *Klein quad.* 58.
Le Phalanger. *De Buffon*, xiii. 92. *tab.*
x. xi. *Schreder*, clii.

226. PHALANGER.

O. with a thick nose: short ears, covered with hair: eight cutting teeth in the upper jaw; two in the lower: hair on the upper part of the body reddish, mixed with light ash-color, and yellow: the hind part of the head, and middle of the back, marked with a black line: the throat, belly, legs, and part of the tail, of a dirty yellowish white; the rest of the tail brown and yellow: the body of the female marked with white: the first and second toes of the hind feet closely united: the claws large: the thumb on the hind feet distinct, like that of the other species: the bottom of the tail is covered with hair, for near two inches and a half; the rest naked: the length, from nose to tail, near nine inches; the tail ten.

This species inhabits the *East Indian* islands, as I am informed by Doctor *Pallas*; nor is it found in *Surinam*, as *M. de Buffon* conjectures.

PLACE.

De zak, of Beurs Rot. *Merian insect. Surinam.* 66. *tab.* lxvi.

Mus sylvestris Americana. *Seb. Mus.* i. 49. *tab.* xxxi. *fig.* 5.

Philander ex rufo helvus in dorso, in ventre ex flavo albicans. *Briffon quad.* 212.

Mus sylvestris Americanus, catulos in dorso gerens. *Klein quad.* 58.

Didelphis dorfigera. D. cauda basi pilosa corpore longiore, digitis manuum muticis, *Lin. syst.* 72.

Le Philandre de Surinam. *De Buffon*, xv. 157. *Mus.* Lev.

227. MERIAN *.

O. with long, sharp-pointed, naked ears: head, and upper part of the body, of a yellowish brown color: the belly white,

* From *Merian*, a German paintress, who first discovered the species at *Surinam*.

E 2

tinged

tinged with yellow: the fore feet divided into five fingers; the hind into four, and a thumb, each furnished with flat nails: tail very long, slender, and, except at the base, quite naked.

SIZE.

The length, from nose to tail, is ten inches. The tail exceeds the length of the body and head.

Inhabits *Surinam*: burrows under ground: brings five or six young at a time, which follow their parent: on any apprehension of danger, they all jump on her back, and twisting their tails round her's, she immediately runs with them into her hole.

* Flying.

228. FLYING.

Flying Opossum. *Stockdale's Bot. Bay*, 297. *White's*, 288.

O. with large ears: whole upper part of the body covered with a rich fur of a glossy black, mixed with grey. On each hip is a tan-color'd spot; all the under side white: tail at the base light color'd; increasing to black as it advances towards the tip: along the middle of the back from the head to tail, is a black line: on the fore feet are five toes; on the hind only three, with a thumb without any nail. From the fore to the hind feet, is a large membrane like the flying squirrel's: length from nose to tail, twenty inches: of the tail twenty two.

Inhabits *New Holland*. The fur exquisitely fine.

Kangaroo.



*P. Magell. Sculp.**• Kangaroo — . 1. 229.*

** Gerboid.

Kangaroo. *Cook's voy.* iii. 577. *tab.* xx.
 Yerboa gigantea. *Zimmerman*, 526.

229. KANGARU.

O. with a small head, neck, and shoulders: body increasing in thickness to the rump.

The head oblong, formed like that of a fawn, and tapering from the eyes to the nose: end of the nose naked and black: upper lip divided.

Nostrils wide and open: lower jaw shorter than the upper: aperture of the mouth small: whiskers on both jaws: those on the upper longest: strong hairs above and below the eyes.

Eyes not large; irides dusky; pupil of a blueish black.

Ears erect, oblongly ovated, rounded at the ends, and thin, covered with short hairs; four inches long.

No canine teeth: four broad cutting teeth in the upper jaw: two long lanceolated teeth in the lower, pointing forward: four grinding teeth in each jaw, remote from the others. This animal has the very singular power of separating the lower incisors, and of bringing them again close to each other.

Belly convex and great.

Fore legs very short, scarcely reaching to the nose; useless for walking.

Hind legs almost as long as the body: the thighs very thick: on the fore feet are five toes, with long conic and strong claws; on the hind feet only three: the middle toe very long and thick, like that of an ostrich; and extends far beyond the two others, which

TEXTU.

LEGS.

which are placed very distinct from it, and are small: the claws short, thick, and blunt: the inner toe of the hind feet is singularly distinguished by having on it two small claws: the bottom of the feet, and hind part, black, naked, and tuberculated, as the animal rests often on them.

TAIL. Tail very long, extending as far as the ears; thick at the base, tapering to a point: the end is black; at the extremity is a strong hard nail: the hair on all parts short and rather hard.

Scrotum large and pendulous, and is placed before the penis.

The female has on the belly an oblong pouch of a vast depth. The receptacle of its young.

Hair on the whole animal soft, and of an ash-color; lightest on the lower parts.

SIZE. Length of the largest skin I examined, three feet three inches from the nose to the tail: of the tail, two feet nine.

Weight of the largest which was shot, was eighty four pounds; but this, on examination of the grinding teeth, had not attained its full growth*. Later accounts inform us, they grow to the weight of a hundred and forty pounds: to the length of six feet to the base of the tail: the tail itself, according to Mr. *Stockdale's* publication, only two feet one.

PLACE. Inhabits the western side of *New Holland*, and has as yet been discovered in no other part of the world. The natives call it *Kangaru*. It lurks among the grass: feeds on vegetables: drinks by lapping: goes chiefly on its hind legs, making use of the fore feet only for digging, or bringing its food to its mouth. The dung is like that of a deer. It is very timid: at the sight of men

* *Cook's voy.* iii. 586.

flies from them by amazing leaps, springing over bushes seven or eight feet high; and going progressively from rock to rock. It carries its tail quite at right angles with its body when it is in motion; and when it alights often looks back: is much too swift for gre-hounds: is very good eating, according to our first navigators; but the old ones, according to the report of the more recent voyagers, were lean, coarse, and tough.

The weapon of defence was its tail, with which it would beat away the strongest dog.

In the spring of the present year I had opportunity of observing the manners of one brought into the capital alive. It was in full health, very active, and very mild and good natured: on first coming out of its place of confinement, it for a little time went on all fours, but soon assumed an upright attitude. It would sport with its keeper in a very singular manner: it first placed its tail in a perpendicular manner, erected its body on it as on a prop, and then raising its whole body, darted its hind legs on the breast of the man. It was capable of striking with great force if provoked: and it could scratch violently with its fore claws.

This is a very anomalous animal: but has more relation to this genus than any other; and in form of its legs comes very near to the *Javan*. N° 219.

230. LESSER
KANGARU.Kangaroo rat. *Stockdale*, 277. *White*, 286.

O. with the visage of a rat; with two sharp pointed cutting teeth in the upper; two larger in the lower, with truncated ends: fore feet very short, furnished with four toes: hind legs and tail resembling the great species. Three toes on each hind foot; the middle greatly exceeding the other two in length: on the belly is a pouch; within which were four nipples. The color above is of a pale brown, lighter on the belly: in size double to that of the *Brown rat*.

MANNERS.

From the form of its parts, the manners probably the same with those of the former: one was shewn in *London* in 1790, but so shy as to elude a perfect description, continually concealing itself in the straw of the box.

231. SPOTTED
KANGARU.*Stockdale's Bot. Bay*, 147.

O. with a long canine visage: upright sharp ears: head and body black; the first plain: the body and thighs marked with large spots of white, thinly dispersed: tail covered with short hairs at the base; the rest very bushy, covered with very long black hairs. Fore legs covered with short hairs for a small space next to the body; the remaining part naked: the feet furnished with five toes; the hind feet with four and a thumb, with a claw. Length from the nose to the tail twenty-five inches: tail about nine.

Inhabits *New Holland*.

Six cutting teeth, two canine teeth, in each jaw.

Sharp nose: slender bodies.

Five toes before; five behind.

XXIII. WEESEL.

- Mustela. Agricola An. Subter.* 485. *Gef-*
ner quad. 752. *Sno-mus. Faun. Suec.* N° 18.
Mustela supra rutila, infra alba. Bris-
son quad. 173.
La Belette. De Buffon, vii. 225. tab.
xxix.
Weefel. Br. Zool. i. 82. Br. Zool. il-
lustr. tab. ci. Schreber, cxxxviii.
LEV. Mus.
Wesfel or Weefel, mustela vulgaris;
in Yorkshire, the Fitchet, or Fou-
mart. Raii syn. quad. 195.
The Whitred. Sib. Scot. iii. 11.
Wiesel. Klein quad. 62.
Mustela nivalis. Lin. syst. 69.

232. COMMON.

W with small rounded ears: whole upper part of the head and body of a pale tawny brown; under side entirely white: a brown spot beneath the corners of the mouth: length, from nose to tail, between six and seven inches: tail two and a half.

Inhabits most parts of *Europe*; is common in *Siberia*, as far as *Kamtschatka*; is met with in *N. America*, even as high as *Hudson's Bay*; found also in *Barbary**. Is very destructive to chickens, birds, and young rabbits; a great devourer of eggs: does not eat its prey on the spot; but after killing it, by a bite near the head, carries it off to its retreat: is a great destroyer of field mice; a gentleman informed me he found eighty-five, newly killed, in one hole, which he believed belonged to this animal:

* *Shaw's Travels*, 249.

very active, runs up the sides of walls with great ease; no place is secure from its ravages: frequents outhouses, barns, and granaries: is a great enemy to rats and mice, and soon clears its haunts from those pernicious animals: brings four or five young at a time: its skin and excrements intolerably fœtid. In *Norway*, *Sweden*, *Russia*, and *Siberia*, it always changes to white at approach of winter. In *Siberia* it is called *Lasmitska*: their skins are sold to the *Chinese* for three or four rubles the hundred.

233. TOUAN.

Le Touan de la *Cepede*, &c. vi. 252. *tab.* lxi.

W with the upper part of head and body blackish; sides of the body, head, and legs, of a bright ferruginous: the lower part of the neck and body of a more pure white: the length from the nose to tail is rather more than five inches: the tail is rather more than two inches long, and tapers to a point.

Inhabits *Cayenne*: lives in hollow trees: lives on worms and insects, and brings two young at a time, which it carries on its back.

Mustela.

- Mustela. *Gesner quad.* 753.
 Wiesel. *Kramer Austr.* 312. *Meyer's An.*
 ii. tab. 23, 24.
 Mustela erminea. M. plantis fiffis,
 caudæ apice albo. *Lin. syst.* 68.
 Wefla. *Faun. suec.* No. 17.
- Mustela hyeme alba, æstate supra ru-
 tila infra alba, caudæ apice nigro.
Briffon quad. 176.
 Le Roselet. *De Buffon*, vii. 240. tab.
 xxix. *Schreiber*, cxxxvii. A.
 Stoat. *Br. Zool.* i. 84. *LEV. Mus.*
- β. ERMINE, when white. Mus Pon-
 ticus. *Plinii lb.* viii. c. 37. *Agri-*
cola An. Subter. 484.
 Armelinus, Hermelein. *Gesner quad.*
 754.
 Gornoftay. *Rzacznfski Poln.* 235.
 Mustela candida, animal ermineum.
- Raii syn. quad.* 198.
 L'Hermine. *De Buffon*, vii. 240. tab.
 xxix. fg. 2. *Briffon quad.* 176. *Schre-*
iber, cxxxvii. B.
 Ermine. *Hfft. Kamtschatka*, 99. *Pontop.*
Norway. ii. 25. *Br. Zool.* i. 84.
LEV. Mus.

234. STOAT.

W with the upper part of the body pale tawny brown: edges of the ears, and ends of the toes, of a yellowish white: throat, breast, and belly, white: end of the tail black: length, from nose to tail, ten inches; tail five and a half: in the N. of Europe and Asia, and in the Highlands of Scotland, it becomes entirely white at the approach of winter, the end of the tail excepted: resumes its brown color in the spring: sometimes found white in England: one was brought to me in a former winter, mottled with brown and white, the season not having been severe enough to effect a total change*; but in February 1780, I saw in my grounds two others in the state of most perfect and beautiful ermines. In the mountains of Southern Asia and Persia, it retains its brown color the whole year†.

* *Br. Zool. illustr. tab.* ci.† *Pallas.*

Inhabits, in great abundance, the N. of *Europe*, and of *Asia*; in *Kamtschatka* and the *Kurile* islands: is met with in *Newfoundland* and *Canada* *: the skins a great article of commerce in *Norway* and *Siberia*: is found in the last place in plenty, in birch forests, but none in those of fir or pine: the skins are sold on the spot, from two to three pounds *sterling* per hundred †: taken in *Norway* in traps, baited with flesh; in *Siberia* ‡, either shot with blunt arrows, or taken in a trap made of two flat stones, propped by a stick, to which is fastened a baited string, which, on the left touch of the animal, falls down and kills it; its manners and food the same with the former; but does not frequent houses: its haunts are woods and hedges, especially such as border on some brook.

235. QUIQUI.

Mustela Quiqui. *Molina Chili*. 273.

W. with a cuneiform nose: ears short and round, with a white spot in the middle: general color brown: legs and tail short: feet like those of a lizard: length from nose to tail thirteen inches.

Inhabits *Chili*: is fierce and irritable: lives under ground: feeds on mice.

236. CUJA.

Mustela Cuja. *Molina Chili*. 272.

W. with black eyes: nose turned up at the end: hair black; very thick, but soft: tail as long as the body, well furnished with hair: very like the ferret in size, shape, and teeth.

* *Charlevoix hist. Nouv. France*, v. 197. † *Muller Ruff. Samlung*. 516.‡ *Bell's travels*, i. 199. *Pontop. Norway*, ii. 25.

Inhabits *Chili* : lives on mice : breeds twice a year, and brings three or four at a time.

La Fouine de la Guiane. *De Buffon*, Suppl. iii. 161. tab. xxiii.

237. S. AMERICAN FITCHET.

W with a long sharp nose : that, the cheeks, throat, and sides of the neck, black : forehead and sides of the head, to the ears, white : ears short, round, and edged with white : from each ear, a narrow white stripe extends along the sides of the neck : the body covered with coarse hairs, grey at their bases, black and white at the ends : legs and feet black, tinged with red : the toes not unlike those of a rat.

Length from nose to tail near twenty-one inches and a half : tail full of hair, of a bright chestnut, mixed with white ; is rather shorter in proportion than that of the *English* Fitchet, to which it has a great resemblance.

SIZE.

Inhabits *Guiana*.

PLACE.

Putorius. *Gesner quad.* 767.

Yltis. *Agricola An. Subter.* 485.

Pole-cat, or Fitchet. *Raii syn. quad.* 196.

Tchorz. *Rzeczynski Polon.* 236.

Mustela foetida. Ittis. Teuffels kind.

Klein quad.

Mustela putorius. M. pedibus fissis, corpore flavo nigricante; ore auri-

culisque albis. *Lin. Syst.* 67. Iller.

Faun. suv. No. 16.

Mustela pilis in exortu ex cinereo albidis, colore nigricante terminatis, oris circumferentia alba. *Briffon quad.* 186.

Le Putois. *De Buffon*, vii. 199. tab. xxiii. *Schreber*, cxxxi.

Pole-cat. *Br. Zool.* i. 77. *Mus. LEV.*

238. FITCHET.

W with the space round the mouth white ; the tips of the ears of the same color : head, body, and legs, of a chocolate-color,

color, almost black: on the sides the hairs are of a tawny cast: tail black: length seventeen inches; tail six.

Inhabits most parts of *Europe*; is common in the temperate parts of *Russia*, but grows scarcer in *Siberia*, except in the desert of *Baraba*, and beyond the lake *Baikal*. None are found north of those places: they are usually met with, in the places just cited, with white or yellowish rumps, bounded with black.

The Fitchet burrows under ground, forming a shallow retreat, about two yards in length, generally terminating under the roots of some large tree; sometimes forms its lodge under haystacks, and in barns: brings five or six young at a time: preys on poultry, game, and rabbits: in winter frequents houses, and will rob the dairy of milk. This animal is excessively foetid; yet the skin is dressed with the hair on, and used as other furs, for tippets, &c.; and is also sent abroad to line cloaths.

239. SARMATIAN.

Mustela farmatica, *Ruffis* Perugina. *Pal-las*, *Itin.* i. 453. *Gueldenstaedt*, in *Nov. Com. Petrop.* xiv. 441. *tab. x. Zim-*

merman, 486. *Schreber*, cxxxii. *Przewiajska*, or the girdled weasel? *Rzac-zinski*, *auct. hist. Polon.* 328.

W with broad, short, round ears, edged with long white hairs: mouth surrounded with white: head, feet, and under side of the body, of a full black: head crossed beyond each eye with a white band, passing beneath the ears along the sides of the neck, and down to the throat: from the hind part of the head, another of yellow passes on each side obliquely towards
the

the shoulders; above, is a third: the upper part of the body is of a brownish black, striped and spotted irregularly with obscure yellow: tail dusky, full of hairs, intermixed with white ones longer than the rest: the end wholly black.

Length, from the tip of the nose, about fourteen inches; of the tail six.

SIZE.

Inhabits only *Poland*, and the southern provinces of *Russia*, between the *Dnieper* and *Volga*; and in *Asia*, the *Caucasian* mountains, and *Georgia*; and by report, *Bucharia*.

PLACE.

It is a most voracious animal, feeding on the marmots, mice, and other lesser animals that inhabit with it the vast plains of the *Russian* empire. Seizes on its prey, and first sucks out the blood; does not meddle with eggs: lives usually in holes made by other beasts, but is not without the power of burrowing: preys by night: sleeps little: very fierce and untameable: its eyes flaming: its smell fœtid, especially when it erects its tail, which it does in anger: is very active: it moves by frequent jumps: copulates in the spring: goes two months, and brings four or eight young, according to the report of the natives.

MANNERS.

Mustela Siberica, Kolonnok, *Russii*. *Pallas* Itin. 701.

240. SIBERIAN.

W. with the face black, whitish about the nostrils, and spotted towards the eyes; the rest of the animal of a deep yellow, nearly approaching to fox or orange color; with the throat sometimes spotted with white: tail very full of hair, and of a deeper

deeper color than the body: hair in general loose and long: the soles of the feet thickly covered with fur.

SIZE.

Its body more slender than the Fitchet, coming nearer to the form of the Stoat: length to the tail twelve inches; of the tail fix.

PLACE.

Begins to appear in the *Altaic* mountains, between the *Ob* and the *Irtish*, from whence it is common, in wooded mountains, to the *Amur* and lake *Baikal*. It has great resemblance in its manners, haunts, and food with the *fable*; but does not extend so far north.

241. FERRET.

Viverra. Plinii lib. viii. c. 55. Agricola An. Subter. 486.

Mustela rustica, viverra, Furo, Ictis.

Gesner quad. 762. Raii syn. quad. 198.

Fret. Klein quad. 63. Schreber, cxxxiii.

Viverra pilis subflavis, longioribus, casta-

neo colore terminatis (masf.) M. pilis ex albo subflavis vestita. (fæm.) Brisson quad. 177.

Mustela Furo. M. pedibus fissis, oculis rubicundis. Lin. syst. 68. Mus. Lev.

W with a very sharp nose: red and fiery eyes: round ears: color of the whole body a very pale yellow: length about fourteen inches; tail five.

Inhabits, in its wild state, *Africa**; from whence it was originally brought into *Spain*†, in order to free that country from the multitudes of rabbits, with which the kingdom was overrun; from thence the rest of *Europe* was supplied with it: is a lively active animal: the natural enemy of rabbits: sucks the

* *Shaw's travels*, 249.

† *Kai γαλες εἰπας ἄς ἡ λυγρὴ εἴη Strabo, lib. iii.*

blood of its prey, seldom tears it: breeds in our climate: and brings five, six, or nine at a time: but is apt to degenerate, and lose its savage nature: warreners* are therefore obliged to procure an intercourse between the female and a pole-cat, by leaving it near the haunts of the last: the produce is a breed of a much darker color than the ferret, partaking more of that of the pole-cat. The ferret has the same disagreeable smell with that animal.

Martes gutturo albo. Agricola An. Subter.
485. *Gesner quad.* 764.
Stein-marter. *Klein quad.* 64.
Martes, alias Foyna, Martin, or Martlet.
Raii syn. quad. 200.
Kuna. *Rzacinski Polon.* 222.
Mustela pilis in exortu albidis castaneo
colore terminatis vestita, gutturo albo.

Briffon quad. 178.
Mustela martes. M. pedibus fassis, cor-
pore fulvo nigricante, gula pallida. *Lin.*
syft. 67. *Mard. Faun. Suec.* No. 15.
La Fouine. *De Buffon.* vii. 186. *tab. xviii.*
Schreber. cxxix.
Martin. *Br. Zool.* i. 79. *LEV. MUS.*

242. MARTIN.

W with broad rounded ears: lively eyes: head brown,
• with a tinge of red: body, sides, and legs, covered
with hair, ash-colored at the bottoms, bright chestnut in the
middle, black at the tips: throat and breast white: belly deep
brown: tail full of hair, and of a dusky color: feet broad, co-
vered at bottom with thick down: claws white: length eighteen
inches: tail ten.

Inhabits most parts of *Europe*, even to the warmer parts of
Russia, but does not extend far east in that empire: is a most
elegant lively animal: capable of being tamed: is very good-
natured and sportive: lives in woods; and breeds in the hollow

PLACE.

* *Br. Zool.* i. 78. ii. 498.

of trees; and often, during winter, shelters in magpies nests: brings from four to six young at a time: destroys poultry, game, &c. and will eat rats, mice, and moles: the skin and excrements have a musky smell: the fur is of some value, and used to line the robes of magistrates.

243. GREY-
HEADED.

Le grande Marte de Guianne. *de La Cépède. de Buffon*, Suppl. vi, 250. tab. lx.

W. with the head and upper part of the sides of the neck greyish: throat and under side of the neck white: all the rest of the body, limbs, and tail, black: length from the tip of the nose to the tail above two feet: of the tail (which is full of hair) eighteen inches.

PLACE.

Inhabits *Guiana*.

244. PINE.

Martes gutture luteo. Agricola An. Subter. 485.

Martes sylvestris. Gesner quad. 765.

Martes abietum. Raii syn. quad. 200.

Baum-Marter. *Klein quad.* 64.

Mustela pilis in exortu ex cinereo al-

bidis castaneo colore terminatis, gutture flavo. Brisson quad. 179.

La Marte. *De Buffon*, vii. 186. *tab.* xxii. *Schreber*, cxxx.

Yellow-breasted Martin. *Br. Zool.* i. 81. *Faunul. Sinent. LEV. Mus.*

W. with a yellow breast and throat: the hair of a dark chefnut-color, and of far superior fineness to the former; in other respects agreeing with it.

Inhabits

Inhabits the N. of *Europe*, *Asia*, and *America*: found also in *Great Britain* *: are not found about the river *Oby*, nor in any part of *Siberia*: inhabits large forests, especially those of pines: never lodges near houses, as the other species is said† to do: *M. de Buffon* says, that it brings but two or three young at a time: its prey is the same with the former; its fur of far greater value. The peninsula of *Kamtchatka*, and *North America*, abound with them: their skins a prodigious article of commerce. Those found about Mount *Caucasus*, with an orange throat, are esteemed the finest in the furriers shops.

Zobela, *Agricola An. Subter.* 485.

Mustela Sobella. Gesner quad. 768.

Mustela Zibellina, the Sable. *Raii syn. quad.* 201. *Klein quad.* 64.

Mustela Zibellina, Aristotele Satherius, Nipho Cebalus. Alciato Mus Samar-

ticus et Scythicus. Charleton Ex. 20.

Mustela Zibellina. M. pedibus fissis,

corpore obscurè fulvo, fronte exalbida, gutture cinereo. *Lin. syst.* 68.

Mustela Zibellina. Nov. Com. Petrop. v. 330. tab. vi.

Martes Zibellina. Mustela obscurè fulvo, gutture cinereo. Brisson quad.

180.

La Zibeline. De Buffon, xiii. 309.

245. SABLE.

W with long whiskers: rounded ears: large feet: white claws: long and bushy tail; color of the hair black at the tips, cinereous at the bottom: chin cinereous, sometimes white, yellow, or spotted: the edges of the ears yellowish: sometimes the hair has a tawny cast; for in spring, after shedding

* *M. de Buffon* says, we have none of these animals in *England*, *Parce qu'il n'y a pas de bois*. That gentleman never did our kingdom the honour of making a progress through it.

† All foreign writers agree in this; but those which inhabit my neighborhood always keep in the woods, except in their nocturnal excursions.

the coat, the color varies: there are instances of their being found of a snowy whiteness*.

The size is equal to that of the Martin, to which it has a great resemblance in form: but this specific distinction must be noted—the tail of the *martin* is much longer than the hind legs, when extended: that of the Sable shorter.

PLACE.

Inhabits *Siberia*, *Kamtschatka*, and some of the *Kurile* isles, which lie between *Kamtschatka* and *Japan*. Notwithstanding what Mr. *Scheffer* says†, it is certain there are none to be found west of the *Uralian* mountains, from whence they increase in numbers, in proportion as you advance eastward.

Sables live in holes in the earth, or beneath the roots of trees: sometimes, like the martin, form nests in the trees, and will skip with great agility from one to the other: are very lively, and much in motion during night: sleep much in the day: one that was kept tame would, on sight of a cat, sit up on its hind legs: excrements most excessively foetid: prey, during summer, on ermines, weefels, and squirrels, but above all on hares; in winter, on birds; in autumn on hurtleberries, cranberries, and the berries of the service-tree: but during that season their skins are at the worst, that diet causing them to itch, and to rub off their fur against the trees: they bring forth at the end of *March*, or beginning of *April*, and have from three to five at a time, which they suckle for four or five weeks‡.

Their chase was, in the more barbarous times of the *Russian* empire, the employ, or rather the tasks, of the unhappy exiles into *Siberia*: as that country is now become more populous, the

* *Strahlenberg* *hist. Russia*, 442.

† *Scheffer* *Lapl.* 136.

‡ *Hist. Kamtschatka*, 109, &c.

fables have in great measure quitted it, and retired farther *North* and *East*, to live in desert forests and mountains. They live near the banks of rivers, or in the little islands in them*: on this account they have, by some, been supposed to be the Σαῦρος of *Aristotle*, *Hist. An. lib. viii. c. 5*; which he classes with the animals conversant among waters.

At present the hunters of fables form themselves into troops, from 5 to 40 each; the last subdivide into lesser parties, and each chuses a leader, but there is one that directs the whole: a small covered boat is provided for each party, loaden with provision, a dog and net for every two men, and a vessel to bake their bread in: each party also has an interpreter for the country they penetrate into: every party then sets out according to the course their leader points out: they go against the stream of the rivers, drawing their boats up, till they arrive in the hunting-country; there they stop, build huts, and wait till the waters are frozen, and the season commences. Before they begin the chase their leader assembles them, they unite in a prayer to the Almighty for success, and then separate: the first fable they take is called God's fable, and is dedicated to the church.

They then penetrate into the woods, mark the trees as they advance, that they may know their way back; and in their hunting-quarters form huts of trees, and bank up the snow round them: near these lay their traps, then advance farther, and lay more traps, still building new huts in every quarter, and return successively to every old one, to visit the traps, and take out the game to skin it, which none but the chief of

* *Aquil's Travels*, 140.

the party must do: during this time they are supplied with provisions by persons who are employed to bring it on sledges, from the places on the road where they are obliged to form magazines, by reason of the impracticability of bringing quantities thro' the rough country they must pass. The traps are a sort of pit-fall, with a loose board placed over it, baited with fish or flesh: when fables grow scarce, the hunters trace them in the new-fallen snow, to their holes, place their nets at the entrance, and sometimes wait, watching two or three days for the coming out of the animal: it has happened, that these poor people have, by the failure of their provisions, been so pinched with hunger, that, to prevent the cravings of appetite, they have been reduced to take two thin boards, one of which they apply to the pit of the stomach, the other to the back, drawing them tight together by cords placed at the ends*: such are the hardships our fellow-creatures undergo, to supply the wantonness of luxury.

The season of chace being finished, the hunters re-assemble; make a report to their leader of the number of fables each has taken; make complaints of offenders against their regulations; punish delinquents; share the booty; then continue at the headquarters 'till the rivers are clear of ice; return home and give to every church the dedicated furs.

COMMERCIAL
HISTORY.

The following is the commercial history of this fur-trade, which Dr. J. R. Foster was so obliging as to translate for me, from *Müller's Sammlung Russ. Geschichte*, iii. 495 to 515, being an abstract from above 20 pages.

* *Bell's Travels*, i. 245.

“ SABLE, SOBOL in *Russian*; ZOBEL in *German*: their price varies, from 1 l. to 10 l. sterling, and above: fine and middling sable skins are without bellies, and the coarse ones are with them: forty skins make a collection called *Zimmer*: the finest sables are sold in pairs, perfectly similar, and are dearer than single ones of the same goodness; for the *Russians* want those in pairs for facing caps, cloaks, tippets, &c. The blackest are reputed the best. Sables are in season from *November* to *February*; for those caught at any other time of the year are short haired, and then called *Nedofoboli*. The hair of sables differs in length and quality: the long hairs, which reach far beyond the inferior ones, are called *Os*; the more a skin has of such long hairs, the blacker they are, and the more valuable is the fur; the very best have no other but those long and black hairs. *Motchkka* is a technical term in the *Russian* fur-trade, expressing the lower part of the long hairs; and sometimes it comprehends likewise the lower and shorter hairs: the above mentioned best sable furs are said to have a black *Motchkka*. Below the long hairs are, in the greater parts of sable furs, some shorter hairs, called *Podofie*, i. e. *Under-Os*: the more *Podofie* a fur has, the less valuable: in the better kind of sables the *Podofie* has black tips, and a grey or rusty *Motchkka*: the first kind of *Motchkka* makes the middling kind of sable furs; the red one the worst, especially if it has but few *Os*: between the *Os* and *Podofie* is a low woolly kind of hair, called *Podfada*; the more *Podfada* a fur has, the less valuable, for the long hair will, in such case, take no other direction than the natural one; for the character of sables is, that notwithstanding the hair naturally lies from the head towards the tail, yet will
it

it lie equally in any direction, as you strike your hand over it: the various combinations of these characters, in regard to *Os*, *Motchkka*, *Podofie*, and *Podjada*, make many special divisions of the goodness of furs: besides this, the furriers attend to the size, preferring always, *ceteris paribus*, the biggest, and those that have the greatest gloss: the size depends upon the animal being a male or female, the latter being always smaller. The gloss vanishes in old furs: the fresh ones have a kind of *bloomy* appearance, as they express it; the old ones are said to have done *blooming*: the dyed fables always lose their gloss, become less uniform, whether the lower hairs have taken the dye or not, and commonly the hairs are somewhat twisted or crisped, and not so strait as in the natural ones: some fumigate the skins, to make them look blacker; but the smell, and the crisped condition of the long hair, betrays the cheat; and both ways are detected, by rubbing the fur with a moist linen cloth, which grows black in such cases.

“ The *Chinese* have a way of dying the fables, so that the color not only lasts, (which the *Russian* cheats cannot do) but the fur keeps its gloss, and the crisped hairs only discover it: this is the reason that all the fables, which are of the best kind, either in pairs or separate, are carried to *Russia*; the rest go to *China*: the very best fables come from the environs of *Nertchinsk* and *Yakutsk*; and in this latter district, the country about the river *Ud* affords sometimes fables, of whom one single fur is often sold at the rate of 60 or 70 rubles (12 or 14l.) The *bellies* of fables, which are sold in pairs, are about two fingers breadth, and are tied together by forty pieces, which are sold from 1 to 2l. sterling. *Tails* are

are sold by the hundred; the very best sable furs must have their tails, but ordinary sables are often cropped, and a hundred sold from 4 to 8 l. sterling: the *legs* or *feet* of sables are seldom sold separately. White sables are rare, and no common merchandize, but bought only as curiosities: some are yellowish, and are bleached in the spring on the snow."

The common sables are scarcely better in hair and color than the martin.

The sable is found again in *North America*. The *Russians* have often discovered the skins mixed with those of martins, in the fur-dresses which the *Ichutki* get from the *Americans* by way of exchange. Their fur is more glossy than that of the *Siberian* sable, and of a bright chefnut-color; but of a coarser quality. It is to be observed, that no sables are found N. E. of the river *Anadyr*, the country of the *Ichutki*.*

The information I received from Doctor *Pallas*, respecting the character of this animal, obliges me to lay aside my notion of its being found in the new world, under the name of *The Fisher*; yet I have reason to suppose I have recovered it on that continent, by seeing the skin of another quadruped highly resembling it, in the cabinet of Mrs. *Blackburn*, sent from *Canada*; which I describe under the name of the *American*.

Its length, from nose to tail, was twenty inches. The trunk of the tail only five inches: but from the rump to the end of the hairs eight. The ears more pointed than those of the *Asiatic* sable: feet very large, hairy above and below: five toes, with white claws on each foot.

AMERICAN.

* Doctor *Pallas*.

COLOR.

Color of the head and ears whitish: whiskers short and black: whole body of a light tawny: feet brown. This seems to have been one of the bleached kind before mentioned.

246. FISHER.

W• with a black nose: strong and stiff whiskers: six small weasel-like teeth above and below: six large canine teeth: four grinding teeth in each upper jaw; three sharp-pointed, the fourth flat: in the lower jaws six; the last flatted, the next tridentated; the next to those bidentated: ears broad and round, dusky on their outsides, edged with white: face and sides of the neck pale brown, or cinereous, mixed with black: hairs on the back, belly, legs, and tail, black; brownish at their base: sides brown: the feet very broad; covered with hair even on their soles: five toes on the fore feet; generally four, but sometimes five, on the hind feet; with sharp, strong, and crooked white claws: fore legs shorter than those behind: tail full and bushy, smallest at the end, seventeen inches long: length, from nose to tail, twenty-eight inches.

Inhabits *North America*: notwithstanding its name, is not amphibious: preys on all sorts of lesser quadrupeds*: by the number of skins imported, is not an uncommon animal; not less than 580 being brought in one season from *New York* and *Pennsylvania*: seems to be the animal called by *Jesselyn*†, the *SABLE*; which, he says, is perfectly black. I have seen many of the skins, which vary in color. *LEV. MUS.*

* By a letter from Mr. *Peter Collinson*, who received the account from *Eartram*, of *Pennsylvania*.

† *Jesselyn's voy.* 87.

Le Vanfire. *De Buffon*, xiii. 167. *tab. xxi. de la Cetele, de Buffon*, Suppl. vii. 249. *tab. lix.*

247. MADAGASCAR.

W with short ears: the hair on the whole body brown at the roots, and barred above with black, and ferruginous: the tail of the same color: the length from nose to tail, about fourteen inches; the tail, to the tip of the hairs at the end, near ten.

Inhabits *Madagascar*.

Le Pekan. *De Buffon*, xiii. 304. *tab. xlii. Schreber*, cxxxiv. *LEV. MUS.*

248. PEKAN.

W with very long and strong whiskers: ears a little pointed: hair on the head, back, and belly, cinereous at the roots, of a bright bay at the ends; very soft and glossy: on the sides is a tinge of grey: between the fore-legs a white spot: legs and tail black: toes covered with thick hair, above and below: claws sharp.

In form like a martin: its length, from nose to tail, one foot seven inches: the length of the trunk of the tail above ten; and the hairs extend an inch beyond.

Inhabits *North America*: described from a skin.

Le Vison. *De Buffon*, xiii. 308. *tab. xliii.*

249. VISON.

W with rounded ears: color of the hair brown, tinged with tawny, very bright and glossy: beneath is a thick down, cinereous tipped with rust color: legs very short: tail dusky.

H 2

Length

SIZE. Length to the tail above seventeen inches: tail, to the extremity of the hairs, nine.

PLACE. Inhabits *North America*: described from the stuffed skins, in 1765, in the cabinet of M. *Aubry*, curate of *Saint Louis*, in *Paris*. A fuller account of this and the preceding animal is desired.

250. WHITE-CHEEKED.

W. with rounded ears: broad and blunt nose: dusky irides: head flat: face, crown, legs, rump, and tail, black: chin and cheeks white: throat of a rich yellow: back and belly of a pale yellow, intimately mixed with cinereous.

SIZE. Length, from nose to tail, eighteen inches: tail of the same length, covered with long hair.

Described from the living animal at Mr. *Brooks's*, April 1774. Place unknown.

251. GRISON.

Le Grifon. *De Buffon*, xvi. 169. tab. xxv. *Allamand*, v. 65. tab. viii. *Schreber*, cxxiv.

W. with large head and eyes: short but broad ears: upper part of the body of a deep brown, each hair tipped with white, which gives it a grey or hoary look: from each side of the forehead extends a broad white line, passing over the eyes, and reaching as far as the shoulders: the nose, throat, and whole under side of the body, thighs, and legs, black.

SIZE.

Length, from the tip of the nose to the origin of the tail, seven inches. By the figure, the tail is little more than half the length of the body.

Inhabits *Surinam*, but is a very scarce animal: first described by Mr. *Allamand*.

Galera, subfusca, cauda elongata, auribus subnudis appressis. *Browne's Jamaica*, 485. tab. xlix. Le Tayra, ou le Galera. *De Buffon*, xv. 252. GUINEA. 155. *Schreber*, cxxxv.

W with the upper jaw much longer than the lower: eyes placed mid-way between the ears and tip of the nose: ears like the human: tongue rough: tail declining downwards, lessening towards the point: feet strong, and formed for digging: shape of the body like that of a rat: size of a small rabbit: of a dusky color: the hair rough.

Inhabits *Guinea*: common about the negro settlements: burrows like a rabbit: very fierce; if drove to necessity will fly at man or beast: very destructive to poultry: seems to be the *Kokeboe* of *Bosman* *, which only differs in color, being red.

PLACE.

Mustela barbara. M. pedibus fissis, atra, collo subtus macula alba triloba. *Lin.* *Mustela maxima* atra moscum redolens. *Tayra*, grosse Belette. *Barrere France* *Æquin*, 155. 253. GUIANA.

W with round ears, covered with down: an ash-colored space between the eyes: a trilobated spot on the lower part of the neck: size of a martin: color black: hairs coarse.

Inhabits *Brazil* and *Guiana*: when it rubs itself against trees, leaves behind an unctuous matter, that scents of musk.

PLACE.

* *Hist. Guinea*, 239.

254. WOOLLY.

La petite Fouine de la Guiane. *De Buffon, Suppl.* iii. 162. *tab.* xxiv.

W• with a long slender nose: upper jaw longer than the lower: ears very short and round: body covered with woolly hair: tail taper, ending in a point, between eight and nine inches long: body, from nose to tail, between fifteen and sixteen.

M. *de Buffon* does not mention the color; by his figure the belly seems white. He says it inhabits *Guiana*. I am doubtful whether it is not one of the above species.

255. ICHNEUMON.

Ichneumon. *Aristot. hist. An. lib.* ix. c. 6.
Oppian Cynege. iii. 407.

Ichneumon. *Linii lib.* viii. c. 24.

L'Ichneumon, que les *Egyptiens* nomment Rat* de Pharaon. *Belinobj.* 95.
Portraits. 66. *Prss. Alp.* i. 234. *Gesner quad.* 566. *Raii syn. quad.* 202.
Sbaro's Travels. 24. 376.

Musella Aegyptiaca. *Klein quad.* 64.

β. INDIA. Quil, vel Quirpele. *Garcia Arom.* 214. *Raii syn. quad.* 197.

Viverra Mungo. *Kampfer Amoen.* 574.

De Mongkos. *Valentin Ambon.* iii.

Serpenticida five Moncus. *Rumf. herb. Amboin. App.* 69. *tab.* xxviii

Indian Ichneumon. *Edw.* 199.

Ichneumon seu vulpecula Ceilonica.

Meles Ichneumon digitis mediis longioribus, lateralibus aequalibus, unguibus subuniforribus. *Hasselquist itin.* 191.

Ichneumon: Mus Pharaonis vulgo. *Brisson quad.* 181.

Viverra Ichneumon. V. cauda e basi incrassata sensim attenuata, pollicibus remotiusculis. *Lin. syst.* 63. *Schreber.* cxvi. A. LEV. MUS.

Seb. Mus. i. 66. *tab.* xli. *fig.* 1.

La Mangouste. *De Buffon.* xiii. 1; o. *tab.* xix. Le Nems, *tom.* xvi. 1. 4. *tab.* xxvii.

Viverra indica. V. ex griseo rufescens, *Brisson quad.* 177. *Raii syn. quad.* 198. *Schreber.* cxvi. LEV. MUS.

W• with bright flame-colored eyes: small rounded ears, almost naked: nose long and slender: body thicker than

* The *Egyptians* never style it *Phar*, or *Moufe*, but *Nems*, or *Ferret*, from its resemblance to that animal. *Hasselquist*, 196. This *Forsk.* confirms, p. 111.

that of others of this genus: tail very thick at the base, tapering to a point: legs short: the hair is hard and coarse: color various in different animals, from different countries; in some alternately barred with dull yellowish brown and white; in others, pale brown and mouse-coloured; so that the animal appears mottled: throat and belly of a uniform brown: beneath the tail is an orifice not unlike that of a badger.

The specimen in the *Asiatick Museum* was thirteen inches and a half long to the origin of the tail; the tail eleven: the *Ægyptian* variety is the largest. Some are forty-two inches long from the nose to the extremity of the tail. M. de Buffon gives the figure of one in the xxvith plate of his Supplement, vol. iii. under the name of *La Grande Mangouste*: the tail is longer, and more slender than that of the common kind, and the hair universally more broken and coarser.

Inhabits *Ægypt*, *Barbary*, *India*, and its *islands*: a most useful animal; being an inveterate enemy to the serpents and other noxious reptiles which infest the torrid zone: attacks without dread that most fatal of serpents the *Naja*, or *Cobra di Capello*; and should it receive a wound in the combat, instantly retires; and is said to obtain * an antidote from a certain herb; after
which

* A fact, as yet, not well established: Botanists are not yet agreed about the species of this sanative plant, whose use, it is pretended, this weasel pointed out to mankind: those who have seen the combats between the *Ichneumon* and *Naia*, never could discover it: *Kampfer*, a writer of the first authority, who visited *India*, and who had a tame *Ichneumon*, and been witness to its battles with the serpent, says no more than that it retired and eat the roots of any herb it met with. It is from the *Indians* he received the account of the root, whose veracity
he

which it returns to the attack, and seldom fails of victory. Is a great destroyer of the eggs of crocodiles, which it digs out of the sand; and even kills multitudes of the young of those terrible reptiles: it was not therefore without reason, that the antient *Ægyptians* ranked the *Ichneumon* among their Deities: is at present domesticated, and kept in houses in *India* and in *Ægypt*; for it is more useful than a cat, in destroying rats and mice: grows very tame: is very active; springs with great agility on its prey; will glide along the ground like a serpent, and seems as if without feet: sits up like a squirrel, and eats with its fore feet: catches any thing that is flung to it: is a great enemy to poultry: will feign itself dead till they come within reach: loves fish: draws its prey, after sucking the blood, to its hole: its excrements very fœtid: when it sleeps, brings its head and tail under its belly, appearing like a round ball, with two legs sticking out. *Rumphius* observes how skilfully it seizes the serpents by the throat, so as to avoid receiving any injury: and *Lucan* beautifully de-

he speaks most contemptuously of. *Amæn. Exot.* 576. *Rumphius* never saw the plant growing; but describes it from a specimen sent him from *Java*; for he says the *Indians* would persuade him that it had no leaves. Vide *Herb. Amboin. App.* 71. All that seems certain is, that the *Indians* have a plant, of whose alexipharmic virtues they have a high opinion, and are said to use it with success against the dreadful *macassar* poison, and the bite of serpents. *Kæmpfer* says he had good success with one species, in putrid fevers, and found it infallible for the bite of a mad dog. As there is no doubt but a most useful plant of this nature does exist in the *Indes*, it is to be hoped that strict enquiry will be made after it. In order to direct their searches, they are referred to

Garcia ab Horto Hist. Aromaticum in Clus. Exot. 214.

Kæmpfer Amæn. Exot. 573. *Rumph. Herb. Amboin. App.* 29.

Amæn. Acad. ii. 89. *Flora Zeylanica*, 46. 190, 239.

scribes

cribes the same address of this animal, in conquering the *Ægyptian Asp.*

*Aspidas ut Pharias cauda solertior hostis
Ludit, et iratas incerta provocat umbra :
Obliquansque caput vanae serpentis in auras
Effusæ toto comprehendit guttura morsu
Lestiferam citra saniem : tunc irrita pestis
Exprimitur, faucesque fluunt pereunte veneno.* Lib. iv. 724.

Thus oft' th' *Ichneumon*, on the banks of Nile,
Invades the deadly *Aspic* by a wile ;
While artfully his slender tail is play'd,
The serpent darts upon the dancing shade ;
Then turning on the foe with swift surprize,
Full on the throat the nimble feizer flies :
The gaping snake expires beneath the wound,
His gushing jaws with poisonous floods abound,
And shed the fruitless mischief on the ground.

Rowe.

Gm. Lin. 85.

256. *CAFRE.*

W with short hairy ears: hairs on the body shining, rude,
• mixed with yellow, black, and brown: tail grows gradually
more slender from the base, tip black.

Inhabits the *Cape of Good Hope*.

Le Surikate. De Buffon, xiii. 72. tab. viii. *Schreber*, cxvii. *Miller's plates*, xx.

257. *FOUR-TOED.*

W with a very sharp-pointed nose: head depressed: cheeks
• inflated: upper jaw much longer than the lower; tip
VOL. II. **I** **black:**

black: whiskers black, arising from warty tubera: irides dusky: region about the eyes black: ears small, rounded, black, lying close to the head.

Tongue oblong, blunt, aculeated backwards.

Six small incisors; two long canine in each jaw, and five grinders on each side.

Back very broad, and a little convex: belly broad and flat.


Legs short: feet small, naked at the bottom; four toes on each: the claws on the fore feet long, like those of the badger; on the hind feet short.

Color of the hairs brown near the bottom; black near the ends, and hoary at the points; those on the back undulated: inside of the legs yellowish brown: tail tufted with black.

Length from nose to tail eleven inches; of tail eight: the last thick at the base, ending pretty abrupt.

Inhabits the *Cape of Good Hope*, where it is called *Meer-rat*: feeds on flesh; preys on mice; is a great enemy to *Blatte*: is always making a grunting noise: is much in motion: sits quite erect, dropping its fore legs on its breast, and moving its head with great ease, as if on a pivot, and appearing as if it listened, or had just spied something new. When pleased, it makes a rattling noise with its tail, for which reason the *Dutch* at the *Cape* call it *Klapper-maus**. It is also found in *Java*, where the *Javanese* style it *Jupe*; the *Dutch*, *Suracatje**. The animal which I examined was brought alive from the *Cape*. Well engraven in *Miller's* plates, tab. xx.

* *Pallas* Miscel. Zool. 59, 60.



Yellow Weasel . N.º 258.



Yellow maucauco. *Syn. quad.* No. 108. *Viverra caudivolvola. Schreber, tab. xlii.* 258. YELLOW.

W with a short dusky nose: small eyes: ears short, broad, and flapping, and placed at a great distance from each other: head flat and broad: cheeks swelling out: tongue very long: legs and thighs short, and very thick: five toes to each foot, separated and standing all forward: claws large, a little hooked, and of a flesh-color.

The hairs short, soft, glossy, closely set together: on the head, back, and sides a mixture of yellow and black: cheeks, inside of the legs, and the belly, yellow: half way down the middle of the belly is a broad dusky list, ending at the tail; and another from the head along the middle of the back to the tail: tail of a bright tawny, mixed with black; is round, and has the same prehensile faculty as some of the monkies have: length from the nose to the tail nineteen inches; of the tail seventeen.

MANNERS.

It was very good-natured and sportive; would catch hold of any thing with its tail, and suspend itself: lay with its head under its legs and belly.

PLACE.

Shewn about twelve years ago in *London*: its keeper said it came from the mountains of *Jamaica*, and called it a *Potto*, the name given by some writers to a species of *Sloth* found in *Guinea*.
LEV. MUS.

Le Kinkajou. De Buffon, xvi. 244. tab. 1.

259. MEXICAN.

W with a short dusky nose: tongue of a vast length: small eyes, encircled with dusky: ears short and rounded, and placed very distant: the hairs short; on the head, upper part of the body, and the tail, the colors are yellow, grey, and black intermixed: the sides of the throat, and under side, and the insides of the legs, of a lively yellow: the belly of a dirty white, tinged with yellow.

SIZE.

The toes separated: the claws crooked, white, guttered beneath. The length from head to tail two feet five (*French*); of the tail, one foot three: the tail is taper, covered with hair, except beneath, near the end, which is naked, and of a fine flesh-color. It is extremely like the former; but larger in all its parts.


MANNERS.

Like the former, it has a prehensile tail, and is naturally very good-natured: goes to sleep at approach of day; wakes towards night, and becomes very lively: makes use of its feet to catch at any thing: has many of the actions of a monkey: eats like a squirrel, holding the food in its hands: has variety of cries during night; one like the low barking of a dog: its plaintive note is cooing; its menacing, hissing; its angry, confused.

Is very fond of sugar, and all sweet things: eats fruits, and all kinds of vegetables: will fly at poultry, catch them under the wing, suck the blood, and leave them without tearing them: prefers a duck to a pullet; yet hates the water.

M. de Buffon calls this animal *le Kinkajou*, after a description
(given





Brazilian Weasel —. N.º 60.

(given by M. Dennis) of one of that name found in *N. America*, described also by *Charlevoix*, under the name of *Carcajou*; both which, in fact, are the same as my *Puma*, N° 189. M. Dennis gives it the same manners; adds, that it climbs trees, watches the approach of the moose, falls on, and soon destroys it. He says, he lost a heifer by one of those animals, which at once eat through its neck; but the quadruped in question never could have the powers attributed to so ferocious a creature. This therefore is new, and by form and manners a proper concomitant of the animal last described.

This animal was brought to *Paris* from *New Spain*, and lived there two or three years. It is a very distinct species from the former, of which M. de *Buffon* gives a very indifferent figure, taken from the animal I describe.

Coati. *Marcgrave Brasil.* 228. *De Laet*, 486. *Raii syn. quad.* 180. *Klein quad.* 72.

Vulpes minor, rostro superiore longiusculo, cauda annulatim ex nigro et rufo variegatâ. *Quachy. Barrere France Equin.* 167.

Viverra nasua. V. rufa, cauda albo annu-

lata. *Lin. syst.* 64.

Ursus naso producto et mobili, cauda annulatim variegata. Brisson quad. 190.

Le Coati brun. *De Buffon*, viii. 358. *tab.* xlviii. *Schreber*, cxviii.

Badger of *Guiana. Bancroft*, 141. *LEV. Mus.*

260. BRASILIAN.

W with the upper jaw lengthened into a pliant, moveable • *proboscis*, much longer than the lower jaw: ears rounded: eyes small: nose dusky: hair on the body smooth, soft, and glossy, of a bright bay color: tail annulated with dusky and bay: breast whitish: length, from nose to tail, eighteen inches; tail, thirteen.

β. DUSKY. Nose and ears formed like the preceding: above and beneath the eye two spots of white: hair on the back and sides dusky at the roots, black in the middle, and tipped with yellow: chin, throat, sides of the cheeks, and belly, yellowish: feet black; tail annulated with black and white; sometimes the tail is of an uniform dusky color *. *Le Coati noiatre* of M. de Buffon, tab. xlvii. Schreber, cxix. The *Coati-mondi* of Marcgrave.

Inhabits *Brazil* and *Guiana*: feeds on fruits, eggs, and poultry: runs up trees very nimbly: eats like a dog, holding its food between its fore-legs: is easily made tame: is very good-natured: makes a sort of whistling noise: seems much inclined to sleep in the day. *Marcgrave* observes, that they are very subject to gnaw their own tails.

261. STIFLING.

Yzquiepatl. *Hernandez Mex.* 332. *Raii*
syn. quad. 181. *Klein quad.* 72.
Meles Surinamenfis. *Briffon quad.* 185.
Ichneumon de Yzquiepatl. *Seb. Mus.* i.

tab. xlii.

Le Coate. *De Buffon?* xiii. 288, tab.
xxxviii. *Schreber*, cxx.

W with a short slender nose: short ears and legs: black body, full of hair: tail long, of a black and white color: length, from nose to tail, about eighteen inches.

Inhabits *Mexico*, and perhaps other parts of *America*. This, and the four following species, remarkable for the pestiferous,

* Described as a distinct species by *Linnaeus*, under the title of *viverra Narica*. *V. subfusca, cauda unicolore*, 64. and by M. *Briffon*, under that of *Ursus naso producto et mobili, cauda unicolore*, 190.

suffocating

suffocating and most fœtid vapour they emit from behind, when attacked, pursued, or frightened: it is their only means of defence: some turn* their tail to their enemy, and keep them at a distance by a frequent *crepitus*; and others ejaculate their urine, tainted with the horrid effluvia, to the distance of eighteen feet: the pursuers are stopped by the terrible stench: should any of this liquid fall into the eyes, it almost occasions blindness; if on the cloaths, the smell will remain for several days, in spite of all washing; they must even be buried in fresh soil, in order to be sweetened. Dogs that are not true bred, run back as soon as they perceive the smell; those that have been used to it, will kill the animal; but are often obliged to relieve themselves by thrusting their noses into the ground. There is no bearing the company of a dog that has killed one, for several days.

Professor *Kalm* was one night in great danger of being suffocated by one that was pursued into a house where he slept; and it affected the cattle so, that they bellowed through pain. Another, which was killed by a maid-servant in a cellar, so affected her with its stench, that she lay ill for several days: all the provisions that were in the place were so tainted, that the owner was obliged to throw them away.

Notwithstanding this, the flesh is reckoned good meat, and not unlike that of a pig: but it must be skinned as soon as killed, and the bladder taken carefully out. The *Virginian* species,

* *Wood's voy.* in *Dampier*, iv. 96; the rest of the account is taken from *Cateby* and *Kalm*.

or *skunk*, is capable of being tamed, and will follow its master like a dog: it never emits its vapour, except terrified.

It breeds in hollow trees, or holes under ground, or in clefts of rocks: climbs trees with great agility: kills poultry, eats eggs, and destroys young birds.

262. STRIATED.

Pole-cat, or Skunk. *Lawson Carolina*. *Viverra putorius*. *V. fusca lineis quatuor dorsalibus parallelis albis*. *Lin.*
 Pole-cat. *Catesby Carolina*, ii.
Mustela Americana foetida. *Klein quad.* *syft.* 64.
 64.
Mustela nigra tæniis in dorso albis. *Le Cœpate. De Buffon*, xiii. 288. *tab.*
Briffon quad. 181. *xl. Schreter*, cxxii.

W with rounded ears: head, neck, belly, legs, and tail, black: the back and sides marked with five parallel white lines: one on the top of the back; the others on each side: the second extends some way up the tail, which is long and bushy towards the end: size of an *European* Pole-cat; the back more arched: varies in the disposition of the stripes.

Inhabits *N. America*: when attacked, bristles up its hair, and flings its body into a round form: its vapour horrid. *Du Pratz* says, that the male of the *Pole-cat*, or *Skunk*, is of a shining black: perhaps the *Coase* of *M. de Buffon* is the male; for his description does not agree with the *Tzquitpatl*, which he makes synonymous.

Chinche:

Chinche. *Feuille obs. Peru*, 1714, p. 272.
 Skunk, Fiskatta. *Kalm's voy. Forster's*
tr. i. 273. *tab.* ii. *Jeffelyn's voy.* 85.
 Enfant du Diable, Bete puante. *Charles-*

voix Nouv. France, v. 196.
 Le Chinche. *De Buffon*, xiii. 294. *tab.*
 xxxix. *Schreber*, cxxi. *LEV. MUS.*

263. SKUNK.

W with short rounded ears: black cheeks: a white stripe
 • from the nose, between the ears, to the back: upper
 part of the neck, and the whole back, white; divided at the bot-
 tom by a black line, commencing at the tail, and passing a little
 way up the back: belly and legs black: tail very full of long
 coarse hair; generally black, sometimes tipped with white, and
 sometimes wholly white*, that figured by *M. de Buffon* entirely
 white: nails on all the feet very long, like those on the fore-feet
 of a badger. Rather less than the former.

Inhabits *Peru*, and *N. America*, as far as *Canada*: of the same
 manners and stench with the others.

Viverra Cinghe. Molina Chili. 269.

264. CINGHE.

W with black hair, changeable into blue: along the back a
 • bed of white round spots from head to tail: head long:
 ears large, well covered with hair, and pendulous: hind legs longer
 than the fore.

Inhabits *Chili*: carries its head low: back arched; which it

* *De la Cupides de Buffon*, Suppl. tom. vii. p. 233. *tab.* lvii.

generally covers with its bushy tail, like the squirrel: digs holes in the ground, in which it hides its young.

MANNERS.

In manners and food agrees with the *Stifling*; and its dreadful stench. *Molina* denies that the smell comes from the urine, but from a greenish oil coming from a bladder seated near the anus, from which it ejects the fetid liquor. The *Indians* value the skins highly, and use them as coverlets for their beds.

265. ZORRINA. *Annas* of the Indians, *Zorrinas* of the Mariputa, Mafutiliqui. *Gumilla* *Organi-
Spaniards. Garcilasso de la Vega,* *noque*, iii. 240. *De Buffon, Schreber,*
331. *cxxiii.*

W with the back and sides marked with short stripes of black and white; the last tinged with yellow: tail long and bushy; part white, part black: legs and belly black. Less than the preceding.

Inhabits *Peru*, and other parts of *S. America*: its pestilential vapour overcomes even the panther of *America*, and stupefies that formidable enemy.

266. RATEL. *Viverra* Ratel. *Sparman* *Stock. Westf.* *Stink-bingsem. Kolben*, ii. 133.
Hondl. 1777, 148. tab. iv. *Blaireau puant. Voy. de la Caille*, 182.

W with a blunt black nose: no external ears; in their place, only a small rim round the orifice: tongue rough: legs short: claws very long: strait, like those of a badger, and guttered beneath: color of the forehead, crown, and whole upper part of the body, of a cinereous grey: cheeks, and space round the

the ears, throat, breast, belly, and limbs, black: from each ear to the tail extends along the sides a dusky line, leaving beneath another of grey.

Length from nose to tail forty inches: of the tail, twelve: fore claws, an inch and three quarters long: hind claws one inch.

SIZE.

PLACE.

MANNERS.

Inhabits the Cape of *Good Hope*; lives on honey, and is a great enemy to bees, which in that country usually inhabit the deserted burrows of the *Æthiopian* boar, the porcupine, jackals, and other animals: preys in the evening: ascends to the highest parts of the deserts to look about, and will then put one foot before its eyes, to prevent the dazzling of the sun. The reason of its going to an eminence, is for the sake of seeing or hearing the *honey-guide* cuckoo*, which lives on bees, and, as it were, conducts it to their haunts: the *Hottentots* profit of the same guide. This animal cannot climb; but when he finds the bees lodged in trees, through rage at the disappointment, will bite the bark from the bottoms: by this sign also, the *Hottentots* know that there is a nest of bees above.

The hair is very stiff, and the hide so tough, probably formed so by nature, as a defence against the sting of bees, that it is not easily killed. It makes a stout resistance by biting and scratching, and the dogs cannot fasten on its skin. A pack which could tear a middle-sized lion to pieces, can make no impression on the

* A new species, very fond of honey, which by its noise directs men, as well as this beast, to the bees nest. *Sparman*, in *Phil. Trans.* lxxvii. 43.

hide of this beast : by worrying, they will leave it for dead, yet without inflicting on it any wounds.

This seems to be the *Stink-bingssem* of Kolben, and *Blaireau-puant* of *La Caille*, which they brand for the horrible stench which it emits from behind, by breaking wind; but the *Abbé* says, it quickly discharges the noisome air. Mr. *Sparman* is silent in respect to this circumstance. The *Hottentots* call it *Ratel*.

267. MARIPUTO.

Viverra Mariputo. Gm. Lin. 88.

W of a black color, with a white bed, reaching from the forehead to the middle of the back: no ears: length twenty inches; tail nine.

Observed by *Mutis* in *New Spain*, about the mines of *Pampluna*: sleeps in the day: forms deep boroughs: wanders about in the night: feeds on worms and insects: is very swift.

268. CEYLON:

Gm. Lin. 89.

W above grey, mixed with dusky: below white. Size of the martin.

Inhabits the *Philippine* isles and *Ceylon*.

Gm. Lin. 90.

269. HERMA-
PHRODITE.

W with three dusky lines along the back: tail longer than the body, with the tip black.

Inhabits *Barbary*. Described by PALLAS.

Cook's first voy. iii. 626. Martin-cat. Stockdale's Bot. Bay, 176.

270. QUOLL.

W with short rounded ears: color black; marked with oblong spots on the body, neck, and tail; belly of a pure white: length from the tip of the nose to the base of the tail, eighteen inches: tail tapers elegantly to a point, and is about the same length as the body.

Inhabits the *Western* side of *New Holland*.

White's Bot. Bay, 181.

271. TAPOA
TAPA.

W with long ears erect: color brown; lightest on the tail: tail about the length of the body, covered with long hairs, and ending in a point: size of a rat.

Inhabits *New Holland*. According to Mr. *White's* description the teeth are so anomalous as to render it difficult to reduce this animal to any genus.

White's

272. SPOTTED
TATA.*White's Bot. Bay, 185.*

THIS, according to Mr. *White's* account and figure, differs from the former only in having the body and sides marked with irregular white spots: tail plain.

273. MUSKY.

W• with nose, lower part of the cheeks, legs, and end of the tail, black: on the middle of the cheeks is a white spot: body cinereous, dashed with yellow: some obscure dusky lines and spots mark the body and lower part of the tail.

Inhabits *Bengal*: smells of musk. Sir *Elijah Impey*.

274. CIVET.

La Civette qu'on nommoit anciennement

Hyæna. Belon obs. 94.

Zibettus. Cail opusc. 43.

Felis Zibethus. Gesner quad. 837.

Animal Zibethicum, masc. et fœm. Hernandez Mex. 580, 581.

Civet Cat. Raii syn. quad. 178.

Coati Civetta vulgo. Klein quad. 73.

Meles fasciis et maculis albis nigris et rufescentibus variegata. Brisson quad. 186.

Viverra Zibetha. V. cauda annulata, dorso cinereo nigroque undatim striato. Lin. syst. 65.

La Civette. De Buffon, ix. 299. tab. xxxiv. Schreber, cxi. LEV. MUS.

W• with short rounded ears: sky-blue eyes: sharp nose; the tip black: sides of the face, chin, breast, legs, and feet black; the rest of the face, and part of the sides of the neck, white, tinged with yellow: from each ear are three black stripes,

stripes, ending at the throat and shoulders: the back and sides cinereous, tinged with yellow, marked with large dusky spots disposed in rows: the hair coarse; that on the top of the body longest, standing up like a mane: the tail sometimes wholly black; sometimes spotted near the base: length, from nose to tail, about two feet three inches; the tail fourteen inches: the body pretty thick.

Inhabits *India**, the *Philippine* isles†, *Guinea*‡, *Æthiopia*||, and *Madagascar*§: the famous drug *musk*, or civet, is produced from an aperture between the privities and the anus, in both sexes, secreted from certain glands. The persons who keep them, procure the musk by scraping the inside of this bag twice a week with an iron *spatula*, and get about a dram each time; but it is seldom sold pure, being generally mixed with suet or oil, to make it more weighty: the males yield the most; especially when they are previously irritated. They are fed, when young, with pap made of millet, with a little flesh or fish; when old, with raw flesh: in a wild state prey on fowl.

PLACE.

* *Dellon's voy.* 82. † *Argensola*, iii. ‡ *Bosman*, 238, *Barbot.* 114.

§ *Rauwolf's Travels*, ii. 482. § *Flacourt's Madagascar*, 154; where it is called *Falanouc*.

275. β ZIBETH.

Animal Zibethicum Americanum. *Hernandez Mex.* 538. Le Zibet. *De Buffon*, 299. tab. xxxi.
Schreber, cxii.
 Felis Zibethus. *Gesner quad.* 836.

W• with short rounded ears: sharp long nose: pale cinereous face: head, and lower part of the neck, mixed with dirty white, brown, and black; sides of the neck marked with stripes of black, beginning near the ears, and ending at the breast and shoulders: from the middle of the neck, along the ridge of the back, extends a black line, reaching some way up the tail: on each side are two others: the sides spotted with ash-color and black: the tail barred with black and white; the black bars broader on the upper side than the lower.

A variety first distinguished from the other by *M. de Buffon*; but figured long before by *Hernandez* and *Gesner*: unknown in *Mexico**, till introduced there from the *Philippine* isles. These animals seem not to be known to the antients.

276. MUSK.

W• with the upper part of the body cinereous, dashed with yellow, and marked with some obscure dusky lines: nose, part of the cheeks, legs, and end of the tail, black; on the middle of the cheeks is a white spot.

PLACE.

Inhabits *Bengal*: has a very strong musky scent: described from a drawing in Sir *Elijah Impey*'s collection.

* *Hernandez Nov. Hist.* ii.

Sonnerat, voy. ii. 144. tab. xci.

277. MALACCA.
CIVET.

W with a long nose: short erect ears: the ground-color of the whole animal perlaceous grey: face black: above each eye four black spots: from the hind part of the head are three black lines; one passes down the hind part of the neck and one down each side of the neck, and over part of the shoulders: from the breast another extends along the middle of the belly; three others begin at the small of the back, and reach to the tail: on the body and thighs are forty-one round black spots: the tail annulated with black and grey: legs and feet black: size of a common cat.

This animal lives by the chase: leaps with great agility from tree to tree: is very fierce: emits a strong musky smell, produced from a liquor which exudes from an orifice above the parts of generation. The *Malayes* collect it, and pretend that it strengthens the stomach, and excites to love. The *Chinese* esteem it highly on account of the last quality; and buy it from the *Malayes*. Inhabits the peninsula of *Malacca*.

MANNERS.

278. GENET.

La Genette. *Belon obs.* 74.Genetha. *Gesner quad.* 549, 550.Genetta vel Ginetta. *Raii syn. quad.* 201.Coati, ginetta Hispanis. *Klein quad.* 73.Mustela cauda ex annulis alternatim albidis et nigris variegata. *Briffon quad.*

186.

Viverra Genetta. V. cauda annulata, corpore fulvo-nigricante maculato. *Lin. Syst.* 65.La Genette. *De Buffon*, ix. 343. *tab.* xxxvi. *Schreber*, cxlii. *Lev. Mus.*

W with ears a little pointed: slender body: very long tail: color of the body a pale tawny, spotted with black; and the ridge of the back marked with a black line: the tail annulated with black and tawny: feet black: sometimes the ground color of the hair inclines to grey: about the size of a martin; but the fur is shorter.

PLACE.

Inhabits *Turky*, *Syria*, and *Spain*; frequents the banks of rivers; smells of musk, and, like the civet, has an orifice beneath the tail: is kept tame in the houses at *Constantinople*, and is as useful as a cat in catching mice.

279. PILOSELLO.

La Genette de la France, de Buffon, Suppl. iii. tab. xlvii. p. 236.

W with nose of a deep brown: face and chin cinereous: a dark line up the forehead: under side of the neck cinereous, mixed with rust: back and whole body of the same color, varied with irregular black spots: outside of the hind legs and thighs dusky: soles of the feet and upper part down to the claws, clothed with down: tail tawny, annulated with black. Lesser than the common ferret.

Inhabits



Josephine — v. 280.



Inhabits the rock of *Gibraltar*, and the mountains of *Ronda*: called by the *Spaniards* *Pilosello*; found also in *France*. After the famous victory near *Tours*, gained over the *Saracens* in 726 by *Charles Martel*, such quantities of rich garments, made of the skins of these animals, were found, as to give occasion to the hero to establish an order of knighthood called *L'Ordre de la Genette*. On the first institution there were sixteen knights; among them were the most illustrious princes of the time. *Martel* himself was the sovereign. The collar consisted of the chains of gold, mixed with enamelled roses of red; pendent was a genet of gold, enamelled with black and red. The order continued during the second race of kings. It is said to have given way afterwards to the *Order of the Star*.

La Fossane. *De Buffon*, xiii. 163. tab. xx. *Schreber*, cxiv. *LEV. MUS.*

280. FOSSANE.

W with a slender body: rounded ears: black eyes: body and legs covered with cinereous hair, mixed with tawny: from the hind part of the head, towards the back and shoulders, extend four black lines: the whole under side of the body of a dirty white: tail semi-annulated.

Inhabits *Madagascar*, and *Guinea*, *Bengal*, *Cochin-china*, and the *Philippine* isles: is fierce, and hard to be tamed: in *Guinea* is called *Berbe*; by the *Europeans*, Wine-bibber, being very greedy of *Palm wine**: destroys poultry: is, when young, reckoned very good to eat†.

PLACE.

* *Fosman*, 239.

† *Flacourt hist. Madagascar*, 512; where it is called *Fossa*.

The specimen in the *Leverian Museum* differed in so many respects, that it is necessary to give a full description of it.

W. with a white spot on each side of the nose, and another beneath each eye: the rest of the nose, cheeks, and throat, black: ears very large, upright, rounded, thin, naked, and black: forehead, sides, ~~thighs~~, rump, and upper part of the legs, cinereous: on the back are many long black hairs: on the shoulders, sides, and rump are dispersed some black spots: tail black towards the end; near the base mixed with tawny, and slightly annulated with black: feet black: claws white.

Size of the Genet, to which it bears a great resemblance: tail of the length of the body.

Six cutting teeth, two canine, in each jaw.

XXIV. OTTER.

Five toes on each foot; each toe connected by a strong web.

Lutra. Agricole An. Subter. 482. *Gesner*
quad. 687. *Raii syn quad* 187.
Wydra. Rzeczinski Polen. 221.

Otter. Klein quad. 91.

Mustela Lutra. M. plantis palmatis nudis,
cauda corpore dimidio brevior. Lin.

syf. 66. *Utter. Faun. suec.* No. 12.

Lutra castanei coloris. B.iffon quad. 201.

Le Loutre. Belon Aquat. 26. *Dr Buffon,*

vii. 134. tab. xi. Schreber, cxxvi. A. B.

Otter. Br. Zool. i. N° 19. Br. Zool. illustr.
tab. c. LEV. Mus.

281. GREATER.

O with short ears: eyes placed near the nose: lips thick: whiskers large: the color a deep brown, except two small spots each side the nose, and another beneath the chin: the throat and breast cinereous: legs short and thick, and loosely joined to the body; capable of being brought on a line with the body, and performing the part of fins; each toe connected to the other by a broad strong web.

The usual length, from the tip of the nose to the base of the tail, is twenty-three inches; of the tail sixteen: the weight of the male otter, from eighteen to twenty-six pounds; of the female, from thirteen to twenty-two. Mr. *Ives* says that the otters of the *Euphrates* are no larger than the common cat.

SIZE.

Inhabits all parts of *Europe*, N. and N. E. of *Asia*, even as far as *Kamtschatka*; is found in none of the *Aleutian* or *Fox Islands*, except in the easternmost, which are supposed to be near to the new world: is found in *Chili**: abounds in *North America*, particularly

PLACE.

* *Molina*, 253.

in *Canada*, where the most valuable furs of this kind are produced: dwells in the banks of rivers; burrows, forming the entrance of its hole beneath the water; works upwards towards the surface of the earth, and makes a small orifice, or air-hole, in the midst of some bush: is a cleanly animal, and deposits its excrements in only one place: swims and dives with great ease: very destructive to fish; if they fail, makes excursions on land, and preys on lambs and poultry. Sometimes breeds in sinks and drains; brings four or five young at a time: hunts its prey against the stream: frequents not only fresh waters, but sometimes preys in the sea; but not remote from shore: will give a sort of loud whistle by way of signal to one another*: is a fierce animal; its bite hard and dangerous: is capable of being tamed, to follow its master like a dog, and even to fish for him, and return with its prey.

The *Latax* of *Aristotle*†; possibly a large variety of Otter ‡.

Siya

* *Leonard Baldner*, iii. 139, fig. This was the person whom Mr. *Willughby* calls a fisherman on the *Rhine*, of whom, on his travels in 1663, he bought a most beautiful and accurate collection of drawings of birds, fish, and a few beasts, frequenting that great river about *Strasbourg*, of which city *Leonard* styles himself, fisherman and burgher. The work is dated in 1653. If I may judge from the elegance of his dress, in the portrait prefixed to the first volume, it should appear that he was a person of considerable wealth. A German MS. description is placed opposite to each drawing. This valuable work is now in the possession of *EDWARD KING, Esq*; and had been bought by a relation of his out of the collection of Dr. *MEAD*.

† *Hist. An. lib. viii. c. 5. vide Br. Zool. i. 86. 4to.*

‡ Sir *Joseph Banks*, on his return from *Newfoundland*, was so obliging as to communicate to me the following account of some animals seen by a gentleman who
went

Siya & Cariguibeiu. *Marcgrave Brasl.* Lutra Brasiliensis. *Raii syn. quad.* 189. 282. BRASILIAN.
 234. *Des Marchais*, iii. 306. *Brisson quad.* 202.

O. with a round head like that of a cat: teeth feline: eyes small, round, and black: large whiskers: ears round: feet in form of those of a monkey, with five toes; the inner the shortest: claws sharp: tail reaching no lower than the feet; flat and naked*.

Hair soft, and not long; entirely black, excepting the head, which is dusky; and the throat, which is yellow.

Bulk of a middling dog. If the same with the otters of *Guiana*, mentioned by *M. de Buffon*, it weighs from forty to a hundred pounds†.

SIZE.

Inhabits *Brasyl*, *Guiana*, and the borders of the *Oronoko*, pro-

PLACE.

went on that voyage; which I take the liberty of inserting here, as they bear some relation to the Otter in their way of life. He observed, sitting on a rock, near the mouth of a river, five animals, shaped like *Italian* grehounds, bigger than a fox, of a shining black color, with long legs, and long taper tail. They often leaped into the water and brought up trouts, which they gave to their young which were sitting with them. On his appearing, they all took to the water and swam a little way from shore, kept their heads out of the water, and looked at him. An old Furrier said, that he remembered the skin of one sold for five guineas; and that the *French* often see them in *Hart-Bay*.

* *Barrere Fr. Equin.* 155.

† *Suppl.* iii. 158, 159.

vided

vided the *Guacki* of *Gumilla* be the same *. *Maregrave* says that it is an amphibious animal; lives on fish, and crustaceous animals, such as cray-fish; and is very dextrous in robbing the nets and wheels of what it finds in them: makes a noise like a young puppy. The flesh is reckoned delicate eating, and does not taste fishy, notwithstanding its food.

If this is the *Guacki*, as probably it is, it burrows on the banks of rivers, and lives in society: are extremely cleanly, and carry to a distance the bones and reliques of the fish they have been eating. They go in troops; are very fierce, and make a strong defence against the dogs; but if taken young are soon tamed.

233. LESSER.

Noerza. *Agricola An. Subter.* 485. *Gesner*

quad. 768.

Latax. *Germ.* Nurtz. nobis Nurek. *Rzac-*zinski *Polon.* 218.

Mustela Lutreola. M. plantis palmatis,

hirsutis ore albo. *Lin. syst.* 66. *Fennis,*Tichurt; *Succis,* Mænk. *Faun. suec.*


Nº 13.

Norka. *Ritchkoff's orenb. Topogr.* i. 295.*Schreber,* cxxvi.

O. with roundish ears: white chin: top of the head hoary;
in some tawny: body tawny and dusky; the short hairs
being yellowish; the long hairs black: the feet broad, webbed,

* *Hist. de l'Orenoque*, iii. 239. *Gumilla* calls them also *Loups ou Chiens d'Eau*, and says they are as large as a setting-dog. There is a great disagreement in the form of the feet, with others of the Otter kind. The writers who have had opportunity of examining it, are silent about the webs, the character of the genus. Till that point is settled, I must remain doubtful whether it be the *Saricourienne* of *Andrew Thevet*, as *M. de Buffon* conjectures. The size of the latter is another objection, which will apologize for my making a separate article of that animal till this point is settled.

and



Lepus Otter—. 1. 283.



and covered with hair: tail dusky, and ends in a point: of the form of an otter, but thrice as small.

Inhabits *Poland*, and the north of *Europe*; and is found on the banks of all the rivers in the country north of the *Taïk*. None are found beyond the lake *Baikal*, or in the north-east parts of *Siberia*. Lives on fish, frogs, and water-insects: its fur very valuable; next in beauty to that of the sable. Caught in *Baskiria* with dogs and traps: is most excessively fœtid.

PLACE.

The *Minx* of *North America* is the same animal with this. The late worthy Mr. *Peter Collinson* * favored me with the following account he received from Mr. *John Bartram*, of *Pensylvania*: ‘The *Minx*,’ (says he) ‘frequents the water like the Otter, and very much resembles it in shape and color, but is less; will abide longer under water than the musk quash, musk rat, or little beaver: yet it will leave its watery haunts to come and rob our hen-roosts; bites off their heads and sucks their blood: when vexed, it has a strong loathsome smell; so may be called the *Water Pole Cat*: its length, from nose to tail, twenty inches; the tail four: is of a fine shining dark brown color.’

From the conformity between the names this animal goes by, in *America* and *Sweden* (*Minx* and *Menk*) it seems as if some

* By letter dated June 14, 1764. *Lawson* also gives some account of it, p. 122, *Hist. Carolina*: He says it is a great enemy to the Tortoises; whose eggs it scrapes out of the sand and devours: eats fresh-water muscles, whose shells are found in great abundance at the mouth of their holes, high up in the rivers, in whose banks they live: may be made domestic: is a great destroyer of rats and mice. *La Hontan*, i. 232, seems to mean the same animal, by his *Foutersaux*, an amphibious sort of little Pole-cats.

Swedish colonist, who had seen it in his own country, first bestowed the name it now goes by, a little changed from the original: the skins are often brought over to *England*.

284. CHINCHIMEN.

Molina Chili, 265.

O with head, whiskers, ears, eyes, shape, and length of the tail, exactly resembling the domestic cat: feet furnished with five toes, palmated, and with strong and crooked claws: body covered with two sorts of hair, one very short and fine, the other long and rude: length from nose to tail twenty inches.

MANNERS.

Inhabits the sea of *Chili*, and very seldom quits that element: goes always in pairs: loves to bask in the sun: creeps to the summit of the rocks, where it is taken in traps: has a hoarse voice, and all the fierceness of the wild cat.

285. SARICOVIERNE.

O of the size of a cat, with a fur fine as velvet, grey and black: web footed.

Lives more in the water than on land: the flesh very delicate, and good to eat.

This appears to me to be the very same with *La petite Loutre d'eau douce de Cayenne*, described and figured by M. de Buffon*, probably from a young animal.

* *Suppl.* iii. 159. tab. xxii.

SIZE.

The body, says he, is seven inches (*French*) in length: the tail six inches and seven lines; slender, taper, tuberculated, convex above, flat beneath: ears rounded, and longer than usual with otters: head, cheeks, and back, dusky; and the sides marked regularly with the same colors, issuing from the back, extending almost to the belly; the spaces between of a yellowish grey: above each eye is a white spot: the throat, and whole under side of the body, of the same color: the toes before are divided; those behind webbed.

M. *de la Borde*, as quoted by M. *de Buffon*, mentions another species of Otter frequent in the rivers of *Guiana*, weighing from twenty to twenty-five pounds, and of a yellowish color.

Mustela Lutris. M. *plantis palmatis pilosis*, *caudae corpore quadruplo brevior* ii. 367. *tab. xvi.*
Lin. syst. 66. *Schreber*, cxxviii. Sea Otter. *Hist. Kamtschatka*, 122. *Müller's voy.* 57, 58;
Lutra marina, *Kalan. Nov. Com. Petrop.*

286. SEA.

O. with a black nose: upper jaw longer and broader than the lower: long white whiskers: irides hazel: ears small, erect, conic: in the upper jaw are six cutting teeth; in the lower four: the grinders broad, adapted for breaking and comminuting crustaceous animals, and shell-fish: skin thick: hair thick and long, excessively black and glossy: beneath that a soft down: color sometimes varies to silvery: legs thick and short: toes covered with hair, and joined by a web: the hind feet exactly like those of a seal, and have a membrane skirting the out-

SIZE.

side of the exterior toe, like that of a goose. Length from nose to tail is usually above three feet; but there have been instances of some being a foot longer: the tail thirteen inches and a half; flat, fullest of hair in the middle; sharp-pointed. The biggest of these animals weigh seventy or eighty pounds.

PLACE.

Inhabits, in vast abundance, *Bering's* island, *Kamtchatka*, the *Alentian* and the *Fox islands* between *Asia* and *America*, and in the interior sea as far as has been discovered to the east of *De Fuca's* freights. They are sometimes seen in troops of hundreds, and a hundred leagues from land. They are entirely confined between lat. 49. and 60 north; and between east long. from *London* 126 to 150. During winter they are brought in great numbers by the eastern winds from the *American* to the *Kurilian* islands.

MANNERS.

Are most harmless animals: most affectionate to their young; will pine to death at the loss of them, and die on the very spot where they have been taken from them: before the young can swim, they carry them in their paws, lying in the water on their backs: run very swiftly; swim often on their back, their sides, and even in a perpendicular posture: are very sportive; embrace each other, and even kiss: inhabit the shallows, or such which abound with sea-weeds: feed on lobsters, fish, *Sepiæ*, and shell-fish: breed once a year; bring but one young at a time; suckle it a year, bring it on shore: are dull sighted, but quick scented: hunted for their skins, which are of great value; sold to the *Chinese* for seventy or a hundred *rubles* apiece: each skin weighs three pounds and a half. The young are reckoned very delicate meat, scarcely to be distinguished from a sucking lamb.

LENGTH from nose to tip of tail four feet four inches : of the tail about thirteen inches : diameter of body scarcely more than five inches and a half : fore legs about three inches and a half long : hind legs about four inches : head small, eyes small, ears most extremely small, scarce visible : fore feet webbed ; hind feet more strongly so : color of the whole animal a rich very deep chestnut or dark brown, rather paler beneath : cheeks and throat paler than the other parts, or more inclining to whitish.

Inhabits *Staten-Land*.

287. **SLENDER.**

PLACE.

DIV.

DIV. II. SECT. III,

DIGITATED QUADRUPEDS.

Without canine teeth; and with two cutting teeth in each jaw.

Generally herbivorous, or frugivorous.

DIV. II. SECT. III. Digitated Quadrupeds.

XXV. CAVY.

Two cutting teeth in each jaw.

Generally four toes on the fore feet, three behind.

Short ears: no tail, or a very short one.

Pace creeping; and slow: numerous breeders: short-lived.

288. CAPIBARA.

Caby-bara. *Marcgrave Brasil*, 230. *Piso**Erasl*, 99. *Rair syn. quad.* 126.River hog. *W'after in Dampier*, iii. 400.Cochon d'Eau. *Des Marchais*, iii. 314.*Sus maximus palustris*. Cabiari, cabionora.*Barrere France Æquin*, 160.Capivard. *Froger's voy.* 99.*Sus hydrochæris*. *S. plantis tridactylis cauda nulla*. *Lin. syst.* 103.*Hydrochærus*, Le Cabiari. *Briffon quad.*80. *De Buffon*, xii. 384. *tab. xlix.**Irabubos*. *Gumilla Orenoque*, iii. 238.

C with a very large and thick head and nose: small rounded ears: large black eyes: upper jaw longer than the lower: two strong and great cutting teeth in each jaw: eight grinders in each jaw; and each of those grinders form on their surface seemingly three teeth, each flat at their ends*: legs short: toes long, connected near their bottoms by a small web; their ends guarded by a small hoof: no tail: hair on the body short, rough, and

* *M. de Buffon* denies this: his description was taken from a young subject; but *Marcgrave* and *Des Marchais*, who had opportunities of examining these animals in their native country, agree in this singular construction of the teeth.

brown

brown; on the nose, long and hard whiskers: grows to the size of a hog of two years old.

Inhabits the country from the Isthmus of *Darien* to the *Brazils*, and even to *Paraguay*; lives in the fenny parts, not remote from the banks of great rivers, such as the *Oronoque*, *Amazons*, and *Rio de la Plata*: runs slowly: swims and dives remarkably well, and keeps for a long time under water: feeds on fruits and vegetables: is very dextrous in catching fish, which it brings on shore, and eats at its ease: it sits up, and holds its prey with its fore feet, feeding like an ape: feeds in the night, and commits great ravages in gardens: keeps in large herds, and makes an horrible noise like the braying of an ass: grows very fat: the flesh is eaten, is tender, but has an oily and fishy taste: is easily made tame *, and soon grows very familiar.

PLACE.

Cuniculus vel Porcellus Indicus. *Gesner quad.* 367.

Cavia Cobaya. *Marcgrave Brasl.* 224. *Piso Brasl.* 102.

Mus seu cuniculus *Americanus et Guineensis*, Porcelli pilis et voce, Cavia Cobaya. *Raii syn. quad.* 223.

Cavia Cobaya *Brasli.* quibusdam mus Pharaonis. Tatu pilosus. *Klein quad.*

49.

Mus porcellus. M. cauda nulla, palmis tetradactylis, plantis tridactylis. *Lin. syst.* 79. *Amœn. Acad.* iv. 190. *tab.* ii.

Cuniculus ecaudatus, auritus albus, aut rufus, aut ex utroque variegatus. *Brisson quad.* 102.

Le Cochon d'Inde. *De Buffon*, viii. 1. *tab.* i. *LEV. Mus.*

289. RESTLESS.

C with the upper lip half divided: ears very large, broad, and rounded at the sides: hair erect, not unlike that of a young pig; color white, or white varied with orange and black, in irregular blotches: no tail: four toes on the fore feet; three on the hind.

* *Muratori hist. Paraguay*, 258.

VOL. II.

N

Inhabits

PLACE.

Inhabits *Brasil*: no mention made by writers of its manners in a wild state: domesticated in *Europe*: a restless, grunting, little animal; perpetually running from corner to corner: feeds on bread, grains, and vegetables: breeds when two months old: brings from four to twelve at a time; and breeds every two months: would be innumerable, but numbers of the young are eaten by cats, others killed by the males: are very tender, multitudes of young and old perishing with cold: are called in *England*, *Guinea Pigs*, being supposed to come from that country. Rats are said to avoid their haunts.

290. ROCK.

Aperca. Brasiliensis, nobis Veldratte, Cuniculus ecaudatus auritus, ex cinereo
vel Boschratte. *Marcgrave Brasil.* 223. rufus. *Briffen quad.* 103.
Piso Brasil. 103. *Rau syn. quad.* 206. L'Aperca. *De Buffon*, xv. 160. *LEV.*
Cavia Aperca. Klein quad. 50. *Mus.*

C with divided upper lip: short ears: four toes on the fore feet; three on the hind: no tail: color of the upper part of the body black, mottled with tawny: throat and belly white: length one foot.

SIZE.

PLACE.

Inhabits *Brasil*: lives in the holes of rocks: is driven out, and taken by little dogs: is superior in goodness to our rabbits: its paces like those of a hare.





Patagonian Cavy—. 1. 291.

Narborough's voy. 33. *LEV. MUS.*

291. PATAGONIAN.

C. with long ears, much dilated near the bottom: upper lip divided: on each side of the nose tufts of soft hairs, and long whiskers: tip of the nose black: face, back, and fore part of the legs, cinereous and rust-colored: breast and sides tawny: belly of a dirty white: on each thigh a white patch: rump black: legs very long: claws long, strait, and black; four on the fore feet; three on the hind: tail a mere naked stump.

This animal is found of the weight of six-and twenty pounds*.

Is found in plenty about *Port Desire*, in *Patagonia*: lives in holes of the earth, like the rabbit: the flesh of a snowy whiteness, and excellent flavor†.

Sir *John Narborough*, and other voyagers, call it a hare.

Paca. Marcgrave Brasl. 224. *Pife Brasl.* 101. *De Laet*, 484.

Mus Brasiliensis magnus, porcelli pilis et voce, *Paca* dictus. *Raii syn. quad.* 226.

Cavia Paca. Klein quad. 50.

Cuniculus major, palustris, fasciis albis notatus. *Paca Marcgrave. Barrere France Equin.* 152.

Mus Paca. M. cauda abbreviata, pedibus

pentadactylis, lateribus flavescenti-lineatis. *Lin. syst.* 81.

Cuniculus caudatus, auritus, pilis obscure fulvis, rigidis, lineis ex albo flavescitibus ad latera distinctis. *Briffon quad.* 99.

Le Paca. De Buffon, x. 269. *tab. xliii. Supplem. iii.* 203. *tab. xliii. LEV. MUS.*

292. SPOTTED.

C. with the upper jaw longer than the lower: nostrils large: whiskers long: ears short and naked: neck thick: hairs short and hard: color of the upper part of the body dark

* *Byren's voy.* 18.

† The same, 19.

brown; the lower part, or sides, marked lengthways with lines of grey spots: the belly white; in some, perhaps young ones, the sides and spots are of a pale yellow: five toes on each foot: only the meer rudiment of a tail: length about ten inches: is made like a pig, and in some parts is called the *Hog-Rabbit**.

Inhabits *Brazil*, and *Guiana*: lives in fenny places: burrows under ground: grows very fat: is esteemed in *Brazil* a great delicacy: grunts like a pig: eats its meat on the ground, not sitting up, as some others of this genus do: are discovered by little dogs, who point out the places they lie in: the master digs over them, and when he comes near transfixes them with a knife; otherwise they are apt to escape: will bite dreadfully. There is a variety quite white, found on the banks of the river *St. Francis*†.

293. BRISTLY.


Agnus filiorum Israel. Prosp. Alp. Ægypt. Ashnoko. *Bruce's travels*, v. 139.
 1. 232. *Hirax Syriacus. Gmel. Lin. Syst.* 167.
Daman Israel. De Buffon Suppl. vi. 276. *Schreber, tab. cccxi. B.*
tab. xlii.

C. with short oval ears, covered within and without with hair: color of the whole animal above grey and ferruginous: from the chin to the extremity of the belly white: on the upper a strong bristly musticho, three inches five eighths long; above the eyes another tuft, two inches and two eighths long; all over the body are scattered similar bristles, two inches and a quarter in length: the toes are fleshy; the lower part naked, the upper covered with

* *Waser's voy. in Dampier*, iii. 401.

† *De Laet*, 484.

black



Bristly Cavy — N. 293.



black hairs : the claws somewhat resemble nails, and are ill adapted for burrowing : no tail : the length of the whole animal is about seventeen inches.

This species was first taken notice of by *Prosper Alpinus*, who calls it *Agnus filiorum Israel* ; the *Daman Israel* of the *Arabs*. He says it is larger than a rabbit, an object of the chase, and that the flesh is sweeter than that of the rabbit.

Inhabits, according to Mr. *Bruce*, mount *Libanus*, the mountain of the Sun in *Abyssinia*, and in great numbers Cape *Makomet*, on the *Arabian* gulph, not far to the east of *Suez*. By *Alpinus* we find they are also inhabitants of *Ægypt*. They are gregarious, and sit by dozens on the great stones to bask in the sun, before the mouth of caves, or clefts in the rocks, their places of refuge at the sight of man. They are justly supposed by Mr. *Bruce* to have been the *Sapben* (mistranslated the coney) of HOLY WRIT. *Solomon* says, ‘ The hills are the refuge for the wild goats, and the rocks for the *Conies*. See his *Sapben*. ‘ The *Sapben*, adds he, are but a feeble ‘ folk, yet make they their houses in the rocks*.’ They retire into the depths of the clefts, and there make themselves a house ; i. e. a nest of straw. Neither the *Christians* of *Abyssinia* and the *Mahometans*, eat the flesh of these animals. The *Arabs* of mount *Libanus* and of *Arabia Petræa* use them as a food. The flesh is as white as a chicken, and free from any rankness.

Mr. *Bruce* supposes that Doctor *Shaw* intended this animal by his *Jird*† ; but, as our learned countryman expressly says that his

PLACE.

* *Proverbs*, ch. xxx. v. 24, 26.

† *Shaw's Travels*, p. 248.

animal has a tail, and that only a little shorter than that of the common rat, we must have recourse to some other species, perhaps genus, for the *Fird of Barbary*.

294. LONG-NOSE.

Aguti vel Acuti. *Margrave Brasl.* 224.

Pise Brasl. 102.

Acuti ou Agoutis. *De Laet*, 484. *Rochefort Antilles*, i. 287.

Mus sylvestris Americanus cuniculi magnitudine, pilis et voce Porcelli, Aguti. *Ratij syn. quad.* 226.

Cavia Aguti. M. cauda abbreviata, palmis tetradactylis, plantis tridactylis, abdomine flavescente. *Lin. syst.* 80.

Cuniculus caudatus, auribus, pilis ex rufoet fusco mixtis rigidis vestitus. *Brisson quad.* 98.

L'Agouti. *De Buffon*, viii. 375. *tab.* 1.

Small Indian Coney. *Brown's Jamaica*, 484.

Long-nosed Rabbet. *Waser's vey. in Dampier*, iii. 401.

Cuniculus omnium vulgatissimus, Aguti vulgo. *Barrere France Equin.* 153*.

C with a long nose: divided upper lip: short rounded ears: black eyes: hard and shining; on the body mixed with red, brown, and black; on the rump, of a bright orange-color: belly yellow: legs almost naked, slender, and black: four toes on the fore feet; three on the hind; tail short, and naked: size of a rabbet.

Inhabits *Brasyl*, *Guiana*, &c. Grunts like a pig: is very voracious: sits on its hind legs, and holds its food with the fore feet when it eats: hides what it cannot consume: hops like a hare: goes very fast: when pursued, takes shelter in hollow trees: is capable of being tamed: when angry, sets up the hair on its

* The animal described by *Seba* under the name of *Cuniculus Americanus*, i. 67. *tab.* xli. seems the same with this, notwithstanding he says, that the hind feet are tetradactylous.

back,

back, and strikes the ground with its feet: is eaten by the inhabitants of *South America*.

Cuniculus minor caudatus, olivaceus, L'Akouchy. *De Buffon*, xv. 258. *Suppl.* liii. 295. OLIVE.
Akouchy. *Barrere France Equin.* 153. 211. tab. xxxvi.
Des Marchais, iii. 303.

A Species of *Aguti*, less than the former, and of an olive-color: which is the whole account left us by M. *Barrere*. *Des Marchais* says, it is more delicate food than the other.

Inhabits *Guiana*, and the islands of *St. Lucia* and *Grenada*: inhabits the woods: lives on fruits: is excellent meat: its flesh is white: easily made tame: makes a cry (but very rarely) like the *restless cavy*: abhors water.

Java hare. *Cateby Carolina, App. tab.* mixto. *Briffon quad.* 98. 296. JAVAN.
xviii. *Mus leporinus. Lin. syst.* 80.
Cavia Javenfis. Klein quad. 50. *Cuniculu Americanus. Seb. Mus.* i. 67.
Cuniculus caudatus auritus, rufesco ad- tab. xlii. fig. 2.

C. with a slender small head: prominent naked ears, rounded at the tops: hairs very stiff like bristles, especially on the back: color of the upper part of the body reddish: breast and belly white: legs long: hind parts large: four toes on the fore feet; three on the hind: tail short: size of a hare.

Inhabits *Surinam* and the hotter parts of *South America*, where it is a common food: the flesh is white, but dry. It is not found

found in *Java* or *Sumatra*, as *Catesby* asserts. Governor *Loten* assures me, that he made the most diligent enquiry after it in most parts of *Java*, but could never find the least traces of any such animal.

297. CAPE.

Cavia capensis. *Pallas Miscel. Zool.* 30.tab. ii. *Spicill.* 16. tab. ii.*Africaansch basterd-mormeldier. Pösmacr**Monogr. De Buffon, Supplem.* iii. 177.
tab. xxix.

C. with a thick head, and full cheeks: ears oval, half hid in the fur: head of the color of a hare: along the top of the back dusky, mixed with grey; sides and belly of a whitish grey: four toes on the fore feet, three behind: tail scarce visible: size of a rabbit, but the shape of the body thick and clumsy.

Inhabits in great abundance the rocky mountains near the *Cape of Good Hope*, where it is called *Kaapsche Dafs*, *Klip Dafs**, or *Cape Badger*: burrows under ground: has a slow creeping pace; a sharp voice, often repeated: is esteemed very good meat.

* *Kolben*, *Dutch edition*, as quoted by *Dr. Pallas*. *La Caille* mentions this species under the name of *Marmot*.

Les Rats musqués, Pilonis. *Rocheport Antilles*, i. 288. *Du Tertre hist. Antilles*, ii. 302. *De Buffon*, x. 2.

298. MUSK.

C of a black or tan color on the upper part of its body: white on the belly: tail very short *: almost as big as a rabbit.

Inhabits *Martinico* and the rest of the *Antilles*: burrows like a rabbit: smells so strong of musk, that its retreat may be traced by the perfume: an obscure species, never examined by a naturalist.

* *Nouv. voy. aux isles de l'Amerique*, i. 438.

XXVI. HARE.

Two cutting teeth in each jaw.

Short tail: or none.

Five toes before; four behind.

299. COMMON.

*Lepus. Plinii lib. viii. c. 55. Gesner quad.*605. *Raii syn. quad.* 204.*Hafse, Klein quad.* 51.*Lepus timidus. L. cauda abbreviata auriculis apice nigris? Lin. syst.* 77. *Hafse, Faun. suec.* No. 25.*Lepus caudatus ex cinereo rufus. Brisson**quad.* 94.*Le Lievre. De Buffon, vi. 246. tab. xxxviii.**Br. Zool.* i. N^o 20.*Arnæb. Forsskal. iv. LEV. Mus.* in which are several curious varieties of colored hares.

H. with ears tip with black: eyes very large and prominent: chin white: long white whiskers: hair on the face, back, and sides, white at the bottom, black in the middle, and tip with tawny red: throat and breast red: belly white: tail black above, white beneath: feet covered with hair even at the bottom: a large hare weighs eight pounds and a half. I am informed, that in the *Isle of Man* some have been known to weigh twelve: its length, from the nose to the tail, two feet.

Inhabits all parts of *Europe*, most parts of *Asia*, *Japan*, *Ceylon**, *Agypt*†, and *Barbary*‡: a watchful, timid animal: always lean: swifter in running up hill than on even ground: when started, immediately endeavours to run up hill: escapes the hounds by various artful doubles: lies the whole day on its seat: feeds by night: returns to its form by the same road that it had

* *Kæmpfer Japan*, i. 126. *Knox Ceylon*, 20.† *Prosp. Alp.* i. 232.‡ *Shaw's Travels*, 249.

taken

taken in leaving it: does not pair: the rutting-season is in *February* or *March*, when the male pursues the female by the sagacity of its nose: breeds often in the year; brings three or four at a time: are very subject to fleas: the *Dalecarlians* make a cloth of the fur, which preserves the wearer from their attacks: the fur is of great use in the hat manufacture: feeds on vegetables: fond of the bark of young trees: a great lover of birch, parfly and pinks: was a forbidden food among the *Britons*: the *Romans*, on the contrary, held it in great esteem.

Inter quadrupedes gloria prima lepus,

was the opinion of *Martial*; and *Horace*, who was likewise a *Bon vivant*, says, that every man of taste must prefer the wing

Fœcundi leporis sapiens seclabitur armos.

There have been several instances of what may be called monsters in this species, *horned hares*, excrescences growing out of their heads, likest to the horns of the roe-buck. Such are those figured in *Gesner's* history of quadrupeds, p. 634; in the *Museum Regium* HAFNIE, No. 48. tab. iv; and in *Klein's* history of quadrupeds, 32. tab. iii; and again described in *Wormius's Museum*, p. 321, and in *Grew's Museum* of the Royal Society. These instances have occurred in *Saxony*, and I think in *Denmark*, to which may be added another near *Afracan**.

HORNED HARES.

A farther account of two straw-colored animals like dogs, which run like hares, and were of the same size, seen by the late navigators in *New Holland*†, will, I fear, be a long desideratum among naturalists.

* *Pallas.*

† *Cook's voy.* iii. 565.
O 2

300. VARYING. *Lepus hieme albus. Forster hist. nat. Vol. G.E. Pb. Transf. lvii. 343.* *Lepus variabilis. Pallas. nov. sp. i. Lev. Mus.*
 Alpine hare. *Br. Zool. i. N^o 20.*

H. with soft hair, in summer grey, with a slight mixture of black and tawny: with shorter ears, and more slender legs, than the common hare: tail entirely white, even in summer: the feet most closely and warmly furred. In winter, the whole animal changes to a snowy whiteness, except the tips and edges of the ears, which remain black, as are the soles of the feet, on which, *in Siberia*, the fur is doubly thick, and yellow. Less than the common species.


PLACE.

Inhabits the highest *Scottish Alps*, *Norway*, *Lapland*, *Russia*, *Siberia**, *Kamtschatka*, and the banks of the *Volga*, and *Hudson's Bay*. In *Scotland*, keeps on the tops of the highest hills; never descends into the vales; never mixes with the common hare, which is common in its neighborhood: does not run fast: apt to take shelter in clefts of rocks: is easily tamed: full of frolic: fond of honey and carraway comfits: eats its own dung before a storm: changes its color in *September*: resumes its grey coat in *April*: in the extreme cold of *Greenland* only, is always† white. Both kinds of hares are common in *Siberia*, on the banks of the *Volga*, and in the *Orenburg* government. The one never changes color: the other, native of the same place, con-

* Vide *Pontop. Norway*, ii. 9. *Scheffer Lapland*, 137. *Sirablenberg Russia*, 370. *Ritchkoff Orenberg Topog.* i. 287.

† *Egede, Greenl.* 62. *Cranitz Greenl.* i. 70.

stantly



1. *Varying Hare* —. N. 300 — 11. *Hooded Rabbit* — p. 104.



stantly assumes the whiteness of the snow during winter. This it does, not only in the open air, and in a state of liberty: but, as experiment has proved, even when kept tame, and preserved in houses in the stove-warmed apartments; in which it experiences the same changes of colors as if it had dwelt on the snowy plains*.

They collect together, and are seen in troops of five or six hundred, migrating in spring, and returning in autumn†. They are compelled to this by the want of subsistence, quitting in the winter the lofty hills, the southern boundaries of *Siberia*, and seek the plains and northern wooded parts, where vegetables abound; and towards spring seek again the mountainous quarters‡. Mr. *Muller* says, he once saw two black hares, in *Siberia*, of a wonderful fine gloss, and of as full a black as jet. Near *Casan* was taken another in the middle of the winter 1768. These specimens were much larger than the common kind.

MIGRATIONS.

BLACK HARES.

In the southern and western provinces of *Russia* is a mixed breed of hares, between this and the common species. It sustains, during winter only, a partial loss of color: the sides, and more exposed parts of the ears and legs, in that season, become white; the other parts retain their colors. This variety is unknown beyond the *Urallian* chain. It is called by the *Russians*, *Russak*; they take them in great numbers in snares, and export their skins to *England* and other places, for the manufacture of hats||. The *Russians* and *Tartars*, like the *Britons* of old, hold

α. SPURIOUS:

* *Pallas* nov. sp. fasc. i. p. 7.
nov. sp. fasc. i. p. 15.

† *Bell's Travels*, i. 238.

|| The same, p. 6.

‡ *Pallas*

the

the flesh of hares in detestation, esteeming it impure: that of the VARIABLE, in its white state, is excessively insipid.

301. AMERICAN.

Hare, hedge Coney. *Lawson*, 122. *Catesby*, App. xxviii.

H. with the ears tip with grey: upper part of the tail black; lower white: neck and body mixed with cinereous, rust-color, and black: legs of a pale ferruginous: belly white; fore legs shorter, hind legs longer, in proportion, than those of the common hare.

Length eighteen inches: weighs from three to four pounds and a half.

Inhabits all parts of *North America*. In *New Jersey*, and the colonies south of that province, it retains its color the whole year. In *New England**, *Canada*, and about *Hudson's Bay*, at approach of winter, it changes its short summer's fur for one very long, silky, and silvery, even to the roots of the hairs; the edges of the ears only preserving their color: at that time it is in the highest season for the table†; and is of vast use to those who winter in *Hudson's Bay*, where they are taken in vast abundance, in springes made of brass wire, to which the animals are led by a hedge made for that purpose, with holes left before the snares for the rabbits to pass through.

They breed once or twice a year, and have from five to seven at a time: they do not migrate, like the preceding, but always haunt the same places: they do not burrow, but lodge under

* *Jesslyn's Rarities*, 22. † *Clerk Californ.* i. 156.

fallen

fallen timber, and in hollow trees: they breed in the grafs; but in fpring shelter their young in the trees, to which they alfo run when purfued; from which, in the fouthern colonies, the hunters * force them by means of a hooked flick, or by making a fire, and driving them out by the fmoke. I have had an opportunity of examining this fpecies in its brown drefs from *Penfyvania*, and its winter's drefs from *Hudfon's Bay*.

- Cuniculus. *Plinii, lib. viii. c. 55. Gefner quad. 362. Agricola An. Subt. 482*
 Rabbet, or Coney. *Raii fyn. quad. 205.*
 Lepusculus, cuniculus terram fodiens, Kaninchen. *Klein quad. 52.*
 Lepus cuniculus. L. cauda abbreviata, auriculis nudatis. *Lin. fyft. 77.*
 Kanin. *Faun. suec. No. 26. Br. Zool. i. N° 22.*
 Lepus caudatus, obfcure cinereus. *Briffon quad. 95.*
 Le Lapin. *De Buffon, vi. 303. tab. l. li. Lev. Mus.*

302. RABBET.

H. with ears almoft naked: color of the fur, in a wild ftate, brown; tail black above, white beneath: in a tame ftate, varies to black, pied, and quite white: the eyes of the laft of a fine red.

Inhabits, in a wild ftate, the temperate and hot parts of *Europe*, and the hotteft parts of *Asia* and *Africa*: not originally *Britifh*; but fucceeds here admirably: will not live in *Sweden*, or the northern countries, except in houfes. *Strabo* † tells us, that they were firft imported into *Italy* from *Spain*. Not natives of *America*; but encrease greatly in *S. America*.

Moft prolific animals: breed feven times in a year: produce eight young at a time: fuppofing that to happen regularly, one

* *Kalm, ii. 45.*† *Lib. ii¹.*

pair may bring in four years 1,274,840. In warrens, keep in their holes in the middle of the day; come out morning and night: the males apt to destroy the young: the skins a great article of commerce; numbers exported to *China*: the fur of great use in the hat-manufacture.

β. ANGORA RABBIT. With hair long, waved, and of a silky fineness, like that of the goat of *Angora*, vol. i. p. 61. and the *Cat*, vol. i. p. 296.

γ. HOODED RABBIT. With a double skin over the back, into which it can withdraw its head: another under the throat, in which it can place its fore feet: has small holes in the loose skin on the back, to admit light to the eyes: color of the body cinereous: head and ears brown.

Described from a drawing, and manuscript account, by Mr. G. Edwards, preserved in the *Museum*; inscribed “A Russian Rabbit;” but I find that it is unknown in that empire.

303. BAIKAL.

Cuniculus insigniter caudatus, coloris Leporini. *Nov. Com. Petrop.* v. 357. tab. xi.

Lepus cauda in supina parte nigra in prona alba. *Brisson quad.* 97. Le Tolai. *De Buffon*, xv. 138.

H. with a tail longer than that of a rabbit: ears longer in the male, in proportion, than those of the *varying* hare: fur of the color of the common hare: red about the neck and feet: tail

tail black above, white beneath : size between that of the common and the *varying* hare.

Inhabits the country beyond lake *Baikal*, and extends through the great *Gobée*, even to *Thibet*. The *Tanguts* call it *Rangwo*, and consecrate it among the spots of the moon*: agrees with the common rabbit in color of the flesh; but does not burrow, running instantly (without taking a ring as the common hare does) for shelter, when pursued, into holes of rocks; so agrees in nature with neither that nor the rabbit. Called by the *Mongols*, *Tolai*. The fur is bad, and of no use in commerce.

Lepus Capensis. L. cauda longitudine capitis, pedibus rubris. *Lin. syst.* 78.

304. CAPR.

H. with long ears dilated in the middle: the outsides naked, and of a rose-color: inside and edges covered with short grey hairs: crown and back dusky, mixed with tawny: cheeks and sides cinereous: breast, belly, and legs, rust-colored: tail bushy, carried upwards; of a pale ferruginous color.

Size of a rabbit.

Inhabits the country three days north of the *Cape of Good Hope*. Is called there the *Mountain Hare*, for it lives only in the rocky mountains; does not burrow. It is difficult to shoot it, as it instantly, on the sight of any one, runs into the fissures of the rocks.

The same species probably extends as high as *Senegal*. M. *Adanson* (44) observes, that the hares of that country are not so large

* *Pallas nov. sp.* i. 20.

as those of *France*; their color between that of the *European* kind and a rabbit; and their flesh white.

305. VISCACCIA.

Lepus viscaccia. Molina Chili, 289. Acosta Peru, 1725, p. 32. Garcilasso de la Vega 331.
in Purchas's Pilgrims, iii. 966. Feuillie

H. with the appearance of a rabbit, excepting the tail; in that part and color like a fox: the tail is long, and turned up, and covered with coarse hair: the rest of the hair soft: size superior to that of a rabbit.

MANNERS.

Inhabits *Peru* and *Chili*: lives under ground, and forms two boroughs one above the other; in the one it keeps its provisions, in the other sleeps: goes out only in the night: its flesh is white and tender. The antient *Peruvians* make stuffs of the hair, which were so fine as to be worn only by the nobility. In *Chili* it goes into the hat-manufactory: its tail is its weapon of defence.

306. CUY.

Lepus pusillus. Molina Chili, 288.

H. with a conoid body: ears small, pointed, and covered with hair: nose long: tail so short as scarcely to be seen: is domesticated and varies in color to white, brown, and spotted with divers colors: fur very fine: size of a field mouse.

Inhabits *Chili*: breeds every month, and brings from six to eight young: is delicate eating.

•• Without

** Without a tail.

Tapeti. *Marcgrave Brasl.* 223. *Piso* *syll.* 78.
Brasl. 102. *Lepus ecaudatus. Brisson quad.* 97.
Cuniculus Brasiliensis Tapeti dictus. Raii *Le Tapeti. De Buffon, xv.* 162.
syn. quad. 205. *Collar'd Rabbet. Wafer's voy. in Dampier,*
Lepus Brasiliensis. L. cauda nulla. Lin. *iii.* 401.

307. BRASILIAN.

H. with very large ears, like the common kind: a white ring round the neck: face of a reddish color: chin white: black eyes: color of the body like the common hare, only darker: belly whitish: no tail: some want the white ring round the neck.

Inhabit *Brasyl*: live in woods: do not burrow: are very prolific: very good meat: found also in *Mexico**, where they are called *Citli*.

Lepus Alpinus. Pallas, nov. sp. fasc. i. 52. *tab. ii. Ilin. ii.* 701. *tab. A. Zimmerman.*

308. ALPINE.

H. with short, broad, rounded ears: head long: very long whiskers: two very long hairs above each eye: color of the fur at the bottom dusky, towards the ends of a bright ferruginous; the tips white; intermixed are several long dusky hairs; but on first inspection the whole seems of a bright bay.

Length of that I saw was nine inches.

* *Hernandez An. Nov. Hisp.* 2.

PLACE.

These animals are first seen on the *Altaic* chain, and extend to lake *Baikal*; and from thence to *Kamtschatka*; and, as it is said, in the new-discovered *Fox* or *Aleutian* islands. They inhabit always the middle region of the snowy mountains, in the rudest places, wooded and abounding with herbs and moisture.

VOICE.

They sometimes form burrows between the rocks, and oftener lodge in the crevices; and are found in pairs, or more, according to conveniency: in cloudy weather they collect together, and lie on the rocks, and give a keen whistle, so like that of a sparrow, as to deceive the hearer. On the report of a gun, they run into their holes; but soon come out again, supposing it to be a clap of thunder, to which they are so much used in their lofty habitations.

By wonderful instinct they make a provision against the rigorous season in their inclement seats. A company of them, towards autumn, collect together vast heaps of choice herbs and grasses, nicely dried, which they place either beneath the overhanging rocks, or between the chafms, or round the trunk of some tree. The way to these heaps is marked by a worn path. In many places the herbs appeared scattered, as if to be dried in the sun and harvested properly. The heaps are formed like round or conoid ricks; and are of various sizes, according to the number of the society employed in forming them. They are sometimes of a man's height, and many feet in diameter, but usually about three feet.

Thus they wisely provide their winter's stock, otherwise they must perish, being prevented by the depth of snow to quit their retreats in quest of food.

They select the best of vegetables, and crop them when in the fullest vigor, which they make into the best and greenest hay by the judicious manner in which they dry it. These ricks are the origin of fertility amidst the rocks; for the reliques, mixed with the dung of the animals, rot in the barren chasms, and create a soil productive of vegetables.

These ricks are also of great service to that branch of mankind who devote themselves to the laborious employ of fable-hunting: for being obliged to go far from home, their horses would often perish for want, if they had not the provision of these industrious little animals to support them; which is easily to be discovered by their height and form, even when covered with snow. It is for this reason that this little beast has a name among every *Siberian* and *Tartarian* nation, which otherwise would have been overlooked and despised. The people of *Yakutz* are said to feed both their horses and cattle with the reliques of the winter stock of these hares.

These animals are neglected as a food by mankind, but are the prey of *sables* and the *Siberian* weasel, which are joint inhabitants of the mountains. They are likewise greatly infested by a sort of *gadfly*, which lodges its egg in their skin in *August* and *September*, which often proves destructive to them.

Lepus Ogotona. Pallas Nov. sp. fasc. i. 59. tab. iii.

309. OGOTONA.

H. with oblong oval ears, a little pointed: shorter whiskers than the former: hairs long and smooth: color of those
on

on the body, brown at the roots, light grey in the middle, white at the ends, intermixed with a very few dusky hairs: a yellowish spot on the nose: space about the rump of the same color: outside of the limbs yellowish: belly white.

Length about six inches: weight of a male, from six ounces and a half to seven and a quarter; of the female, from four to four and three quarters.

Inhabits only the country beyond lake *Baikal*, and from thence common in all parts of the *Mongolian* desert, and the vast desert of *Gobee*, which extends on the back of *China* and *Tibet*, even to *India*. It lives in the open vallies, and on gravelly or rocky naked mountains. These little creatures are called by the *Mongols*, *Ogotona*: are found in vast abundance: live under heaps of stones, or burrow in the sandy soil, leaving two or three entrances. Their holes run obliquely: in these they make a nest of soft grass. The old females make for security many of these burrows near each other, that they may, if disturbed, retreat from one to the other.

They wander out chiefly in the night. Their voice is excessively shrill, a note like that of a sparrow, twice or thrice repeated; but very easily to be distinguished from that of the *Alpine* hare.

They live in the vallies, principally on the tender bark of a sort of *Service* and the dwarf elm; in the spring on different herbs. Before the approach of severe cold, in the early spring, they collect great quantities of herbs, and fill their holes with them, which the inhabitants of the country consider as a sure sign of change of weather.

Directed by the same instinct with the former, they form in
autumn

autumn their ricks of hay of a hemispherical shape; about a foot high and wide: in the spring these elegant heaps disappear, and nothing but the reliques are seen.

They copulate in the spring, and about the latter end of *June* their young are observed to be full grown.

They are the prey of hawks, magpies, and owls: but the Cat *Manul* makes the greatest havock among them: and the ermine and fitchet is equally their enemy.

Lepus pusillus. Pallas Nov. sp. i. 31. tab. i. Nov. Com. Petrop. xiii. 531. tab. xiv. 310. CALLING.
Zimmerman.

H. with a head longer than usual with hares, and thickly covered with fur, even to the tip of the nose: numerous hairs in the whiskers: ears large and rounded: legs very short: soles furred beneath: its whole coat very soft, long, and smooth, with a thick long fine down beneath, of a brownish lead-color: the hairs of the same color; towards the ends of a light grey, and tipped with black: the lower part of the body hoary: the sides and ends of the fur yellowish.

Length about six inches: weight from three ounces and a quarter to four and a half; in winter scarcely two and a half.

Inhabits the south-east parts of *Russia*, and about all the ridge of hills spreading southward from the *Urallian* chain; also about the *Irtish*, and in the west part of the *Altaic* chain; but no where

SIZE.

PLACZ.

where in the east beyond the *Oby*. They delight in the most funny vallies, and herby hills, especially near the edges of woods, to which they run on any alarm.

MANNERS.

They live so concealed a life as very rarely to be seen: but are often taken in winter, in the snares laid for the ermines; so are well known to the hunters. About the *Volga* they are called *Semlanoi Saetsbik*, or *Ground Hare*: the *Tartars*, from their voice, style them *Tschotsebot* or *Itstiskan*, or the *Barking Mouse*; the *Kalmucs* call them *Rusla*.

They chuse for their burrows a dry spot, amidst bushes covered with a firm sod, preferring the western sides of the hills; in these they burrow, leaving a very small hole for the entrance; and forming long galleries, in which they make their nests: but those of the old ones, and females, are numerous and intricate: their place would be scarcely known but for their extreaments, and even those they drop, by a wise instinct, under some bush, lest their dwelling should be discovered by their enemies among the animal creation.

VOICE.

It is their voice alone that betrays their abode: it is like the piping of a quail, but deeper, and so loud as to be heard at the distance of half a *German* mile. It is repeated by just intervals thrice, four times, and often six. This is wonderful, as this little animal does not appear to be particularly organized for the purpose. The voice is emitted at night and morning: seldom in the day, except in rainy and cloudy weather. It is common to both sexes; but the female is silent for some time after parturition, which is about the beginning of *May*, N. S. They bring forth six at a time, blind, and naked; which she suckles often, and covers carefully with the materials of her nest.

YOUNG.

These



1. Callung Hare —, V. 310.
 2. Alpine Hare —, V. 308.
 3. Cygona Hare —, V. 309.



These most harmless and inoffensive animals never go from their holes: feed and make their little excursions by night: drink often: sleep little: are easily made tame: will scarcely bite when handled. The males in confinement are observed to attack one another, and express their anger by a grunting noise.

XXVII. BEAVER.

Two cutting teeth in each jaw.

Five toes on each foot.

Tail compressed, and covered with scales.

311. CASTOR.

Καστωγ. *Arist. hist. An. lib. viii. c. 5.**Oppian. Halieut. i. 398.*Fiber. *Plinii lib. viii. c. 30. Agricola An.**Subt. 482. Delon Aquat. 25.*Castor. *Gesner quad. 309. Rondel. 236.**Scheneveld Icib. 34.*Beaver. *Rail syn. quod. 209.*Bobr. *Rzaczynski Polon. 215.*Biber. *Klein quad. 91. Kramer Austr. 315.*

Castor castaneæ coloris, cauda horizonta-

liter plana. *Briffon quad. 90.*Castor Fiber. C. cauda ovata plana. *Lin.**lyst. 78.*Batwer, Biur. *Faun. suec. N° 27.*Le Castor, ou Le Bievre. *De Buffon, viii.**282. tab. xxxvi.*Beaver. *Br. Zool. i. Pl. 9. LEV. Mus.*

B. with strong cutting teeth: short ears, hid in the fur: blunt nose: hair of a deep chestnut brown: tail broad, almost oval, compressed horizontally, covered with scales: the fore feet small; the hind large: length from nose to tail, about three feet: tail eleven inches long, three broad.

PLACE.

Inhabits *Europe*, from *Lapland* to *Languedoc**: in great plenty in the *North*: a few are yet found in the *Rhone*†, the *Gardon*, the *Danube*, the *Rhine*, and the *Vistula*. I have an instance of two old and six young being taken in 1742, at *Gornichem*, in *Holland*; another in 1757 in the *Yffel*, in *Guelderland*; and another in 1770 in the *Maas*, near the village *Hedel*, not far from *Bois le duc*: this last weighed forty pounds, and had two bags of *castoreum*, weigh-

* *De Buffon*, viii. 286.† *Ibid.*



Castor Beaver—V. 311.



ing four ounces, and of excellent quality. It had inhabited the river for some years, and done much damage to the willow-trees, with whose bark its stomach was found full. They are much more frequent in the *Lippe*, above *Wesel*, from which river they might descend into those of *Holland* *.

Abound in the *Asiatic* part of the *Russian* empire; are found in companies, or associated, about the *Konda*, and other rivers which flow into the *Oby*. They are met with dispersed, or in the state of *Terriers*, in the wooded parts of independent *Tartary*, and in the chains of mountains which border upon *Siberia*. None are to be seen in *Kamtschatka*, by reason of the interruption of the woods beyond the river *Kozyma*; nor yet in the new-discovered islands west of that country: only in the isle of *Kadjak*, the nearest to *America*, some skins have been procured by the *Russians*, which probably were got by the natives from *America*, in whose northern parts they are found in prodigious abundance.

The most industrious of animals: nothing equals the art with which they construct their dwellings. They chuse a level piece of ground, with a small rivulet running through it. This they form into a pond, by making a dam across; first by driving into the ground stakes five or six feet long, placed in rows, wattling each row with pliant twigs, and filling the interstices with clay, ramming it down close. The side nearest to the water is sloped; the other perpendicular. The bottom is from ten to twelve feet thick; but the thickness gradually diminishes to the top, which is about two or three. The length of these dams is sometimes not less than a hundred feet.

MANNERS.

* *Martine's Katechism. Natur.* ii. 143.

Their houses are made in the water collected by means of the dam, and are placed near the edge of the shore. They are built on piles; are either round or oval; but the tops are vaulted; so that their inside resembles an oven, the top a dome. The walls are two feet thick; made of earth, stones, and sticks, most artificially laid together; and the walls within as neatly plaistered as if with a trowel. In each house are two openings; one into the water, the other towards the land. The height of these houses above the water is eight feet. They often make two or three stories in each dwelling, for the convenience of change, in case of floods. Each house contains from two to thirty beavers; and the number of houses in each pond is from ten to twenty-five. Each beaver forms its bed of moss; and each family forms its magazine of winter provision, which consists of bark and boughs of trees. This they lodge under water, and fetch it into their apartments as their wants require. *Larsson* says they are fondest of the *jassafra*, ash, and sweet-gum. Their summer food is leaves, fruits, and sometimes crabs and craw-fish; but they are not fond of fish.

To effect these works, a community of two or three hundred assembles; each bears his share in the labor: some fall, by gnawing with their teeth, trees of great size, to form beams or piles; these are gnawed all round in as regular a manner as a cutter cuts in felling a tree, bringing the bottom of the wood to a point*: others roll the pieces along to the water; others dive, and with their feet scrape holes, in order to place them in; while others

* This will be best understood by inspecting the specimens in the LEVERIAN MUSEUM.

exert their efforts to rear them in their proper places: another party is employed in collecting twigs, to wattle the piles with; a third, in collecting earth, stones, and clay; a fourth is busied in beating and tempering the mortar; others, in carrying it on their broad tails to proper places, and with the same instrument ram it between the piles, or plaister the inside of their houses. A certain number of smart strokes with their tail, is a signal given by the overseer, for repairing to such or such places, either for mending any defects, or at the approach of an enemy; and the whole society attend to it with the utmost assiduity. Their time of building is early in the summer; for in winter they never stir but to their magazines of provisions, and during that season are very fat. They breed once a year, and bring forth, the latter end of the winter, two or three young at a birth.

Besides these associated beavers, is another sort, which are called *Terriers*; which either want industry or sagacity to form houses like the others. They burrow in the banks of rivers, making their holes beneath the freezing depth of the water, and work up for a great number of feet. These also form their winter stock of provision.

Beavers vary in their colors: the finest are black; but the general color is a chestnut brown, more or less dark: some have been found, but very rarely, white; others spotted: both varieties are preserved in the LEVERIAN MUSEUM. The skins are a prodigious article of trade; being the foundation of the hat-manufactory. In 1763 were sold, in a single sale of the *Hudson's Bay Company*, 54,670 skins. They are distinguished by different names. *Coat Beaver* is what has been worn as coverlets by the *Indians*. *Parchment Beaver*, because the lower side resembles it.

Stage

Stage Beaver is the worst, and is that which the *Indians* kill out of season, on their stages or journies. The valuable drug *Castoreum** is taken from the inguinal glands of these animals. The antients had a notion it was lodged in the testicles, and that the animal, when hard pressed, would bite them off, and leave them to its pursuers, as if conscious of what they wanted to destroy him for.

*Imitatus Castora, qui se
Eunuchum ipse facit, cupiens evadere damno
Testicularum.* JUVENAL, xii. 34.

Just as the BEAVER, that wise thinking brute,
Who, when hard hunted on a close pursuit,
Bites off the parts, the cause of all the strife,
And leaves them as a ransom for his life. DRYDEN.

* The *Russian Castoreum* is so much better than the *American*, that we give two guineas a pound for that, and only 8s. 6d. for the last; the first being less waxy, and pulverises readier. Notwithstanding we take this drug from *Russia*, we export there vast numbers of Beaver skins. The flesh is reckoned good eating, being preserved, after the bones are taken out, by drying it in the smoke.

MS. hist. Hudson's Bay.

Muffasco,

Musfascus. *Smith's Virginia*, 27.

Musquash. *Jessely's Voy. New England*, 86.

Musk Rat. *Larson Carolina*, 120.

Castor Zibethicus. C. cauda longa compresso-lanceolata, pedibus fissis. *Lin. f.*

Castor cauda verticaliter plana, digitis omnibus a se invicem separatis. *Briffon quad.* 93.

L'Ondatra. *De Buffon*, x. i. tab. i.

Rat Musqué. *Charlevoix Nouv. France*, v. 157. *Lescarlot N. Fr.* 350. *LEV. Mus.*

312. MUSK.

B. with a thick blunt nose: ears short, and almost hid in the fur: eyes large: toes on each foot separated; those behind fringed on each side with strong hairs, closely set together: tail compressed sideways, and very thin at the edges, covered with small scales, intermixed with a few hairs: color of the head and body a reddish brown: breast and belly ash-color, tinged with red: the fur very fine: length, from nose to tail, one foot; of the tail, nine inches: in the form of its body, exactly resembles a beaver.

Inhabits *North America*: breeds three or four times in a year *, and brings from three to six young at a time: during summer, the male and female consort together: at approach of winter, unite in families, and retire into small round edifices, covered with a dome, formed of herbs and reeds cemented with clay: at the bottom are several pipes, through which they pass in search of food; for they do not form magazines like the beavers: during winter, their habitations are covered many feet deep with snow and ice; but they creep out and feed on the roots that lie

* *MS. hist. Hudson's Bay.*

beneath: they quit their old habitations annually, and build new ones. The fur is soft, and much esteemed: the whole animal, during summer, has a most exquisite musky smell: which it loses in winter: perhaps the scent is derived from the *Calamus Aromaticus*, a favorite food of this animal. *Lescarbot* says they are very good to eat.

§13. GUILLINO.

Castor Huidobrius. *Molina Chili*, 266.

B. with a square head: short and round ears: small eyes: color grey; dark on the back, whitish on the belly. It has two sorts of hair, like the common beaver: one short and fine, and susceptible of any dye; the other species of hair long and hard: the toes of the fore feet bordered with a membrane; the hind feet webbed: the back very broad: the tail long and hairy, and length from the nose to the tail three feet; height two feet.

MANNERS:

Inhabits the deepest rivers and lakes of *Chili*: has the *foramen ovale* half closed: can live long under water: feeds on fishes and crabs: is fierce and bold, and will seize its prey in sight of mankind: is killed by the hunters when it comes to discharge its excrements, which it does always in the same place: most beautiful stuffs are made of the fur, resembling velvet; it is also of great use in the manufacture of hats.

M. *Molina* calls it *Huidobrius*, from the family name of his patron, the marquiss of *Casa Reale*.

M. *Molina*

M. *Molina* was one of the *Jesuits* whom the *Spaniards* expelled out of *South America*. They robbed him of all his effects and manuscripts: by a singular fortune he found in *Italy* the manuscript which furnishes us with the valuable natural history of *Chili*.

XXVIII.
PORCUPINE.

Two cutting teeth in each jaw.

Body covered with long, hard, and sharp quills.

Upper lip divided.

314. CRESTED.

Υ51. 8. *Aristot. hist. An. lib. i. c. 6. Oppian**C. reg. iii. 391.**Hystrix. Plinii l. b. viii. c. 35. G. fner quad.**5⁶ 3. Rati syn. quad. 206.**Acanthion cristatus. Klein quad. 66.**Hystrix orientalis cristata. Seb. Mus. i. 79.**tab. l.**Hystrix cristata. H. palmis tetradactylis, plantis pentadactylis, capite cristato, cauda abbreviata. Lin. syst. 76. Hasselquist. itin. 290.**Hystrix capite cristato. Brisson quod. 85.**Le Porc-epic. De Buffon, xii. 402. tab. li. lii. Faunul. Sinens.*

P with a long crest on the top of the head, reclining backwards, formed of stiff bristles: the body covered with long quills; those on the hind part of the body nine inches in length, very sharp at the ends, varied with black and white; between the quills a few hairs: the head, belly, and legs, are covered with strong bristles, terminated with soft hair, of a dusky color: the whiskers long: ears like the human: four toes before, five behind: tail short, and covered with quills: length, from nose to tail, two feet; tail, four inches.

Inhabits *India*, the sand-hills on the S. W. of the *Caspian sea*, southern *Tartary*, *Persia*, and *Palestine*, and all parts of *Africa*: is found wild in *Italy*; but is not originally a native of * *Europe*: is brought into the markets of *Rome*, where it is eat †. The *Italian* porcupines have shorter quills, and a lesser crest, than those of *Asia* and *Africa*: is an harmless animal: lives on fruits, roots, and vegetables: sleeps by day, feeds by night: the report of its

* *Agricola An. Subt. 486.*† *Ray's Travels, i. 311. Pb. Tr. abridg. v. 147.*

darting





Long-tailed Porcupine . N. 36.

darting its quills fabulous: when angry, retires and runs its nose into a corner, erects its spines, and opposes them to its assailant: makes a snorting noise.

These animals produce a *Bezoar*; but, according to *Seba*, only those which inhabit *Java*, *Sumatra*, and *Malacca*. These *Bezoars* were very highly valued, and have been sold for five hundred crowns apiece. It has also been pretended that a stone was procured from the head of this animal, infinitely more efficacious than other *Bezoars**; but this may be placed among the many impositions of oriental empirics.

Erinaceus Malacensis. Gm. Lin. 116. Seb. Mus. i. p. 81. tab. 41. fig. i.

315. MALACCA.

P with large pendulous ears: no crest: quills like the preceding, with the interstices filled with long hairs, resembling bristles: eyes large and bright: hair on the legs, and belly covered with short reddish prickly hairs: toes five in number, which might determine *Linnaeus* to place this animal among the hedge-hogs.

Inhabits the peninsula of *Malacca*.

Porcus aculeatus sylvestris, seu Hystrix orientalis singularis. Seb. Mus. i. 84. tab. lii.

Acanthion cauda prælonga, acutis pilis horrida, in exitu quasi panniculata. Klein quad. 67.

Hystrix cauda longissima, aculeis undique obfita, in extremo panniculata. Brisson quad. 89.

Hystrix macroura. H. pedibus pentadactylis, cauda longissima: aculeis clavatis. Lin. syst. 77.

316. LONG-
TAILED.

P with long whiskers: short naked ears: large bright eyes: body short and thick, covered with long stiff hairs as sharp

* *Tavernier, ii. 154.*

as needles, of different colors, according as the rays of light fall on them: feet divided into five toes; that which serves as a thumb turns backwards: the tail is as long as the body, very slender to the end, which consists of a thick tuft: the bristles appearing as if jointed; are thick in the middle, and rise one out of the other like grains of rice; are transparent, and of a silvery appearance.

Inhabits the isles of the *Indian Arcipelago*, and lives in the forests.

317. BRAZILIAN.

Tlaquatzin. *Hernandez, Mex.* 330.

Cuandu. *Brazilienfisus, Lufitanis.*

Ourico cachiro. *Marcgrave Brasil.* 233.

Piso Brasil. 99. 325.

Iron Pig. *Nieuboff,* 17.

Hystrix Americanus. Raii syn. quad. 208.

Hystrix prehensilis, H. pedibus tetradactylis, cauda elongata prehensili seminuda. Lin. fssl. 76.

H. cauda longissima, tenui, medietate extrema aculeorum experte, 87.

H. Americanus major, 88.

Hystrix longius caudatus, brevioribus aculeis. Barrere France Equin. 153.

Hystrix minor leucophaeus, Gouandou. ibid.

Chat epineux. Des Marchais, iii. 303.

P with a short blunt nose: long white whiskers: beneath the nose a bed of small spines: top of the head, back, sides, and base of the tail, covered with spines; the longest, on the lower part of the back and tail, are three inches in length, very sharp, white, barred near their points with black; adhere closely to the skin, which is quite naked between them; are shorter and weaker as they approach the belly: on the breast, belly, and lower part of the legs, are converted into dark-brown bristles: feet divided into four toes: claws very long; on the place of the thumb a great protuberance: tail eighteen inches long, slender, and taper towards the end; the last ten inches is almost naked, having only



Brazilian Porcupine — 1317.



only a few hairs on it; has, for that length, a strong prehensile quality.

Inhabits *Mexico* and *Brazil*: and extends to *Chili*: lives in the woods: preys not only on fruits, but poultry: sleeps in the day, preys by night: makes a noise with its nostrils as if out of breath: grunts like a sow*: grows very fat: its flesh white, and very good: climbs trees, but very slowly; in descending, twists its tail round the branches, for fear of falling: is no more capable of shooting its quills than the first: may be tamed. *Piso* says there is a greater and lesser kind.

This species is very rarely brought into *Europe*. I had opportunity of describing it from a specimen some time in possession of Mr. *Greenwood*; who was so obliging as to permit me to have a drawing made of it, from which a very faithful figure is here given. *M. de Buffon*† has made mention of this animal in his work; but unjustly reproaches *Marcgrave* with confounding it with the *Mexican* species.

Hoitzlacuatzin, seu Tlacuatzin spinosus,
Hystrix novæ Hispaniæ. *Hernandez*
Mex. 322.
Hystrix novæ Hispaniæ. H. aculeis appa-

rentibus, cauda brevi et crasso. *Briffon*
quod. 86.
Le Coendou De Buffon, xii. 421. tab. liv.

318. MEXICAN.

P of a dusky color, with very long bristles intermixed with the down: the spines three inches long, slender, and varied with white and yellow; scarcely apparent, except on the tail,

* *Vocem edit ut Sus*, iii. *Marcgrave*, 233.

† Under the name of *Le Coendou*, xii. 421. tab. liv.

which

which is, according to *Hernandez*, thicker and shorter than that of the preceding species. He adds, that the tail, from the middle to the end, is free from spines.

SIZE. According to *Hernandez*, it grows to the bulk of a middle-sized dog. *M. de Buffon* says, its length is sixteen or seventeen inches from the nose to the tail; the tail nine, *French* measure, but taken from a mutilated skin.

PLACE. Inhabits the mountains of *Mexico*: lives on the summer fruits, and may be easily made tame. The *Indians* pulverise the quills, and say they are very efficacious in gravelly cases; and, applied whole to the forehead, will relieve the most violent head-ach. They adhere till filled with blood, and then drop off.

319. CANADA.

Porcupine from *Hudson's Bay*. *Edw.* 52.

Ellis's voy. 42. *Clerk's voy.* i. 177. 191.

Cavia Hudsonis. *Klein quad.* 51.

Hystrix dorsata. H. palmis tetradactylis, plantis pentadactylis, cauda mediocri,

dorso solo spinoso. *Lin. syst.* 76.

Hystrix aculeis sub pilis occultis, cauda brevi et crassa. *Briffon quad.* 87.

L'Urson. *De Buffon*, xii. 426. tab. Iv. *LEV. Mus.*

P with short ears, hid in the fur: head, body, legs, and upper part of the tail, covered with soft, long, dark brown hair: on the upper part of the head, back, body, and tail, numbers of sharp strong quills; the longest on the back, the left towards the head and sides; the longest three inches; but all are hid in the hair: intermixed, are some stiff straggling hairs, three inches longer than the rest, tipped with dirty white: under side of the tail white: four toes on the fore feet, five behind, each armed with long claws, hollowed on their under side: the form of the body is exactly that of a beaver; but is not half the size:

one, which Mr. *Banks* brought from *Newfoundland*, was about the size of a hare, but more compactly made: the tail about six inches long.

These animals vary in color. Sir *Astton Lever* had one, which is entirely white.

Inhabits *N. America*, as high as *Hudson's Bay*: makes its nest under the roots of great trees, and will also climb among the boughs, which the *Indians* cut down when one is in them, and kill the animal by striking it over the nose: are very plentiful near *Hudson's Bay*, and many of the trading *Indians* depend on them for food, esteeming them both wholesome and pleasant: feed on wild fruits and bark of trees, especially juniper: eat snow in winter, drink water in summer; but avoid going into it: when they cannot avoid their pursuer, will sidle towards him, in order to touch him with the quills, which seem but weak weapons of offence; for, on stroking the hair, they will come out of the skin, sticking to the hand. The *Indians* stick them in their noses and ears, to make holes for the placing their ear-rings and other finery: they also trim the edges of their deer-skin habits with fringes made of the quills, or cover with them their bark-boxes.

PLACE.

XXIX.
MARMOT.

Two cutting teeth in each jaw.

Four toes before, five behind.

Short ears, or none.

Tail covered with hair, and of a middling length; in some very short.

320. ALPINE.

Mus Alpinus. *Plinii lib.* viii. c. 37. *Agri-*
cola An. Suter. 484. *Gesner quad.* 743.*Raii syn. quad.* 221.Glis marmota. *Klein quad.* 56. *Hist. Mur.*
Alp. 230.Murmeltier. *Kramer Auflr.* 317.

Mus marmota. M. cauda abbreviata sub-

pilosa, auriculis-rotundatis, buccis gib-
bis. *Lin. syst.* 81.Glis pilis e fusco et flavicante mixtis ves-
titus. Glis flavicans, capite rufescente.
Brisson quad. 116, 117.La Marmotte. *De Buffon*, viii. 219. *tab.*
xxviii.

M. with short round ears, hid in the fur: cheeks large: color of the head and upper part of the body brownish ash, mixt with tawny: legs and lower part of the body reddish: tail pretty full of hair: length, from nose to tail, about sixteen inches; tail six: body thick.

PLACE.

Inhabits the loftiest summits of the *Alps* and *Pyrenean* mountains: feeds on insects, roots, and vegetables: while they are at food, place a centinel, who gives a whistle on seeing any sign of danger, on which they instantly retire into their holes: form holes under ground, with three chambers of the shape of a Y, with two entrances; line them well with moss and hay; retire into them about *Michaelmas*, and, stopping up the entrances with earth, continue in a torpid state till *April*: when taken out remain insensible, except brought before a fire, which revives them:

them: they lodge in society, from five to a dozen in a chamber: will walk on their hind feet: lift up their meat to their mouth with their fore feet, and eat it sitting up: bring three or four young at a time: are very playful: when angry, or before a storm, make a most strange noise; a whistle so loud and so acute, as quite to pierce the ear: grow very fat about the backs: are sometimes eaten; but generally taken in order to be shewn, especially by the *Savoyards*: grow very soon tame, and will then eat any thing: are very fond of milk, which they lap, making at the same time a murmuring noise, expressive of their satisfaction: very apt to gnaw any cloaths or linen they find: will bite very hard.

M. with a blunt nose: short rounded ears: cheeks puffed, and of a grey color: face dusky: nose black: hair on the back grey at bottom, black in the middle, and the tips whitish: belly and legs of an orange-color: toes black, naked, and quite divided; four, and the rudiments of another, on the fore feet; five behind: tail short, and of a dusky color: was rather larger than a rabbit.

321. QUEBEC.

Inhabits *Hudson's Bay* and *Canada*. Mr. *Brooks* had one alive a few years ago; it was very tame, and made a hissing noise: perhaps is the species which the *French* of *Canada* call *Siffleur*.

PLACE.

It has lately been described by Dr. *Pallas*, under the name of *Mus empetra* *.

* *Nov. sp. quadr. fasc. i. 75.*

VOL. II;

S

Bahama

322. MARYLAND.

Bahama Cony. *Catesby Carolina*, ii. 79.
 Monax, *Catesby Carolina App.* xxviii.
 Monax, or Marmotte of *America*. *Edw.*
 104.
 Glis Marmota, *Americanus*. *Klein quad.*
 56. *De Buffon*, Suppl. iii. 175.

Glis fuscus. Glis fuscus, rostro e cinereo
 cærulefcente. *Brisson quad.* 115.
 Mus Monax. M. cauda medio ocri pilosa,
 corpore cinereo, auriculis subrotundis,
 palmistetradactylis, plantis pentadacty-
 lis. *Lin. Syst.* 81.

M. with short rounded ears: black prominent eyes: nose sharper than that of the last: nose and cheeks of a blueish ash-color: back of a deep brown color: sides and belly paler: tail half the length of the body, covered with pretty long dusky hair: toes divided, and armed with sharp claws: four toes before, five behind: feet and legs black: is about the size of a rabbit.

PLACE.

Inhabits *Virginia* and *Pennsylvania*: during winter sleeps under the hollow roots of trees: is found also in the *Bahama* isles: lives on wild fruits and other vegetables: its flesh is very good, tasting like that of a pig: when surprized, retreats to holes in the rocks: whether it sleeps, during winter, in the climate of those isles, is not mentioned.

323. HOARY.

M. with the tip of the nose black: ears short and oval: cheeks whitish: crown dusky and tawny: hair universally rude and long: that on the back, sides, and belly, cinereous at the root, black in the middle, whitish at the tip, so that the animal has a hoary appearance: legs black: claws dusky; four before, five behind: tail black, mixed with rust-color.

About the size of the former.

Inhabits

Inhabits the northern parts of *North America*. Described from a specimen in the LEVERIAN MUSEUM.

Bobak Swiftch. *Rzaczynski Polon.* 233. tab. xviii.
 Bobak. *Beauplan hist. Ukrain, Churchill's* Sogur. *Rubruquis's Travels in Purchas.* iii.
coll. i. 600. *Forster hist. Velgæ, Phil.* 6.
Transf. lvi. 343. *De Buffon, xiii.* 136. *Arctomys. Pallas nov. sp. fasc. i. g.* tab. v.

324. BOBAK.

M with small oval thick ears, covered with greyish white down; with longish hairs on the edges: eyes small: whiskers small: color about the eyes and nose dusky brown; among the whiskers ferruginous: upper part of the body greyish, intermixed with long black or dusky hairs, tipped with grey: throat rust-colored: rest of the body, and the inside of the limbs, of a yellowish rust-color: four toes on the fore feet, with a short thumb furnished with a strong claw: five toes behind: tail short, slender, full of hair.

Length from nose to tail sixteen inches; of the trunk of the tail, about four: the hairs extend an inch beyond the end of the trunk.

SIZE.

Inhabits the high but milder and sunny sides of mountainous countries, which abound with fissile or free-stone rocks: seek dry situations, and such which are full of springs, woods, or sand. They are found in *Poland*, and the south of *Russia*, among the *Carpathian* hills; they swarm in the *Ukraine*, about the *Boristhenes*, especially between the *Sula* and *Supoy*; and again between the *Boristhenes* and the *Don*, and along the range of hills which extend to the *Volga*; they are found about the *Taik* and other

PLACE.

neighboring rivers. Inhabit the southern desert in *Great Tartary*, and the *Altaic* mountains east of the *Irtis*; cease to appear in *Siberia*, on account of its northern situation; but are found again beyond lake *Baikal*, and about the river *Argun* and lake *Dalay*; in the sunny mountains about the *Lena*; and very common in *Kamtschatka*, but rarely reach as high as *lat. 55*.

MANNERS.

They burrow extremely deep, and obliquely, to the depth of two, three, or four yards: they form numbers of galleries with one common entrance from the surface; each gallery ends in the nest of the inhabitant. Sometimes the burrows consist of only one passage. They are found in great abundance about the sepulchral tumuli, as they find they can penetrate with great facility in the soft dry earth; but they are very common in the rocky strata; and in the mineral part of the *Uralian* chain, often direct the miners to the veins of copper, by the fragments which appear at the mouth of their holes, flung out in the course of their labors. In very hard and rocky places, from twenty to forty of these animals join together to facilitate the work, and live in society, each with its nest at the end of its respective gallery; but the fewest galleries are found in the softest ground, and very frequently only a single one. In each nest they collect, especially towards autumn, the finest of hay, and in such plenty, that sufficient is found in one nest for a night's food for a horse.

During the middle and sunny part of the day they sport about the entrance of their holes, but seldom go far from them; on the sight of man they retire with a slow pace, and sit upright near the mouth, and give a frequent whistle, listening at the approach. In places where they live in large families, they al-

ways place a centinel to give notice of any danger, during the time the rest are feeding.

They are very fond of oleraceous plants: in a state of confinement eat cabbage and bread very greedily, and drink milk with great eagerness; but refuse water, and seem never affected with thirst: they are mild and good-natured; never quarrel or fight about their food in a wild state, and when confined, and placed with others, caught in distant parts, and strangers to them, grow instantly familiar with them: then very soon become tame, even when taken in full age; but the young immediately become familiar.

The number produced at a birth is not certainly known, probably at times eight; the females being furnished with that number of teats: they breed early, for in *June* the young are observed to be of half the size of the old.

They lie torpid during winter, except those which are kept tame in the stove-warmed rooms of the country; and even then, finding a defect of that warmth which the snug nest of their subterraneous retreat would afford, in cold nights creep for shelter into the very beds of the inhabitants. In that state they will not absolutely refuse food, but eat very little, and that with a seeming disgust; nature allotting for them, in the wild state, a long sleep and cessation from food, the result of plentitude previous to its commencement. They sometimes escape from confinement, find a retreat, and get their winter's sleep, and return to their master in the spring; but lose much of their gentle manners.


They grow very fat: the fat is used for softening of leather: the skins are used by the *Koreki*, people of *Yakutsk*, and the *Russians*,

flans, for cloathing. The *Calmucs* take them in small nets with large meshes, placed before their holes. The inhabitants of *Ukraine* catch them in *May* or *June*, by pouring water into the holes, which forces them into the nets. In *South Russia* they are destroyed by means of a log of wood with a weight at top; the end directed into a wooden box placed at the mouth of the hole, which falls as soon as the animal comes out, and oppresses it by the weight. Their flesh tastes like that of a hare, but is rank.

The *Calmucs* are very fond of the fat ones, and even esteem them medicinally: on the contrary, the *Mahometan Tartars* not only abstain from their flesh, but even give them protection; so that near the hords they are extremely numerous: these *Tartars* esteem a warren of *Bobaks* near them to be very fortunate, and think it a sin to kill one of them, a swallow, or a dove; but at the same time abominate the following animal.

In *Chinese Tartary* they are the propagators of *Rhubarb*, which grows among their burrows: the manure which they leave about the roots contributes to its increase; and the loose soil they fling up, proves a bed for the ripe seed; which, if scattered among the long grass, perishes without ever being able to reach the ground.





1. *Quebec* . Marmot — . V. 321.
 2. & 3. *Earless* . Marmot — . V. 326.

Mus Maulinus. *Molina Chili*, 284.

325. MAULINE.

M. with pointed ears: elongated nose: whiskers disposed in four rows: the tail longer than that of the common kind: five toes on each foot; an anomalous distinction: hair like the common: in size twice as large.

Discovered in the province of *Maule* in *Chili*, in 1764, and inhabits the woods: makes a stout defence against the dogs, which conquer it not without difficulty.

MANNERS.

Mus Noricus aut Citellus. *Agricola An. Subier.* 485. *Gesner quad.* 737. *Rat. syn. quad.* 220.

Ziesel. *Schwenkfeld. Theriograph.* 86.

Mus citellus. M. cauda abbreviata, corpore cinereo, auriculis nullis. *Lin. syst.* 80.

Tsitsjan. *Le Bruyn voy. Musc.* ii. 402 *.

Cuniculus caudatus, auriculis nullis, cinereus. *Briffon quad.* 101.

Le Ziesel. *De Buffon*, xv. 139.

Le Soufflik — 144. 195. *Suppl.* iii. 191. *tab.* xxxi.

Mus Marmotta. *Forster hist. nat. Volge.* Ph. Trans. lvii. 343.

Mus Citillus. *Pallas nov. sp. fasc.* i. 119. *tab.* vi. vii. B. *Nov. com. Petrop.* xiv. 549. *tab.* vii.

Earless Marmot. *Syn. quad.* 276. *Caspar M.* — 272.

326. EARLESS.

M. with a cinereous face: over each eye a white line: teeth yellow: whiskers black and long: no ears: hind part of

* Un chien courant que j'avois, y prit dans la plaine un petit animal nommé *Zits-jan*, qu'il m'apporta en vie, et un autre peu après, lesquels je fis éventrer, pour les conserver. C'est un espece de rat de campagne, de la grosseur d'un écureuil, qui a la queue courte, et le poil et la couleur d'un lapreau, aussi bien que la forme, hors qu'il a la tête plus grosse, et les deux dents de dessous la moitié plus longues que celles de dessus. Il a aussi les pattes de devant plus courtes que celles de derrière, avec quatre grifes, et une plus petite, et cinque à celles de derrière, ressemblant assez à celles d'un singe.

the head, and whole back, of a pale yellowish brown; often distinctly spotted with white; sometimes undulated with grey: under side of the body, and legs, of a yellowish white.

Tail covered with long hair; brown above, bordered with black on each side; each hair tipped with white: under part of a bright rust-color: three middle toes of the fore feet long: claws long and sharp: exterior and interior toes short; the last remote from the others: its claws short and blunt.

SIZE.

Length one foot; of the tail, to the end of the hairs, four inches and a half.

Inhabits *Bokemia*, *Austria*, *Hungary*, and from the banks of the *Volga* to *India* and *Persia*; through *Siberia*, and *Great Tartary*, to *Kamtschatka**; some of the intervening isles, such as *Kadjak*; and even the continent of *America* itself.

Burrows, and forms its magazine of corn, nuts, &c. for its winter food†: sits up like a squirrel while it eats: some inhabit the fields in *Siberia*, others penetrate into the granaries; the first form holes under ground, with a double entrance, where they sleep during winter: those which inhabit granaries, are in motion during the cold season. They couple the beginning of *May*, about the *Lena*, but about *Asracan* earlier, and bring from five to eight young, which they bring up in their burrows, and cover with hay: only one animal inhabits each burrow: the females are always separate from the males, except in the coupling season: whistle like the marmot: are very irascible; quarrelsome among themselves, and bite very hard: sit in multitudes near their holes: are very fond of salt: taken in numbers on board the barges

* *Yevraskha*, or *Marmotte minor*. *Gmelin voy. Siberia*, ii. 448.

† *Raii syn. quad.* 220.

which

which are laden with that commodity at *Solikamsky*, and fall down into the *Volga* below *Casan*.

Are both herbivorous and carnivorous; feed on plants, and destroy the young of small birds, and the lesser mice.

The *Bokemian* ladies were wont to make cloaks of the skins; we see them at this time made use of for linings, and appear very beautiful for that purpose, especially the spotted kind.

M. with truncated ears, the apertures large: short tail: upper fore teeth truncated; lower, slender and pointed: four toes on every foot, each furnished with claws: walks on the whole hind feet as far as the heel: color, testaceous red.

327. GUNDI.

Size of a small rabbit.

Inhabits *Barbary* towards *Mount Atlas*, near *Massifin*. Described by the late Mr. *Rohman*, a *Swede*. This account was communicated to me by Mr. *Zimmerman*. *Gundi* is its *Arabic* name, which I retain.

M. with short ears: head and body of a cinereous brown; the ends of the hairs white: two cutting teeth above; four below: no tail.

328. TAILLESS.

I communicated a drawing of this species to Mr. *Berwick*, who has given an engraving of it at p. 374 of his ingenious performance. Inhabits *Hudson's Bay*. In the LEVERIAN MUSEUM.

XXX.
SQUIRREL.

With two cutting teeth in each jaw.
Four toes before, five behind.
Long tail, cloathed with long hair.

329. COMMON.

- Sciurus. Gefner quad. 845. Raii syn. quad. 214.*
Więxiorka Rzeczynski Polon. 225.
Eichhorn. Klein quad. 53.
Sciurus vulgaris. Sc. auriculis apice barbatus, palmis tetradactylis, plantis pentadactylis, Lin. fyll. 86.
Ikorn, Gmelin Faun. succ. N° 37.
Sciurus rufus quandoque griseo admixto. Briffon quad. 104.
L'Ecuireuil. De Buffon, vii. 258. tab. xxxii. Br. Zool. i. 93. Lev. Mus.

S with ears terminated with long tufts of hair: large lively black eyes: head, body, legs, and tail, of a bright reddish brown: breast and belly white: hair on each side the tail lies flat. In *Sweden*, and *Lapland** changes in winter into grey. In many parts of *England* is a beautiful variety with milk-white tails.

Inhabits *Europe*; the northern and temperate parts of *Asia*; and a variety is even found as far *south* as the isle of *Ceylon*: is a neat, lively, active animal: lives always in woods: in the spring, the female is seen pursued from tree to tree by the males, feigning an escape from their embraces. Makes its nest of moss and dried leaves, between the fork of two branches: brings three or four young at a time: has two holes to its nest: stops up that on the side the wind blows, as *Pliny*† justly remarks: lays in a hoard of

* *Faun. Succ. and Scheffer Laph. 135.*† *Lib. viii. c. 38.*

winter provision, such as nuts, acorns, &c.; in summer, feeds on buds and young shoots: is particularly fond of those of fir, and the young cones: sits up to eat, and uses its fore-feet as hands: covers itself with its tail: leaps to a surprising distance: when disposed to cross a river, a piece of bark is its boat; its tail the sail*.

A large kind of grey squirrel is found about the upper parts of the river *Obi*, in the district of *Kuznetsk*, and is called *Teleutskaya Belka*, or the squirrel of the *Teleutian Tartars*: it is as large again as the common grey squirrels of those parts, and is preferred to them, on account of the silvery gloss of the skin. Few are sent into *Russia*, the greatest part being sent into *China*, and sell for 6*l.* or 7*l.* sterling per thousand†.

A white variety is found common in *Siberia*.

A beautiful black variety about lake *Baikal*. In the **LEVE-RIAN MUSEUM** is a most elegant specimen of this kind.

α. **WHITE-LEGGED SQUIRREL.** The head, whole upper part of the body, sides, and toes, of a reddish brown: face, nose, under side of the neck, belly, fore legs, inside of the ears and thighs, white: ears slightly tufted with black: tail long, covered with dusky hairs, much shorter than those in the *European* kind. *Br. Mus.*: by the catalogue, said to be brought from *Ceylon*.

* *Rzaczniski, Klein, Seheffer, Linnæus.*

† *Memorabilia Russ. Asiat. in Muller's Samlung. Russ. vii. 124.*

330. CEYLON.

Sciurus Zeylanicus, pilis in dorso nigricantibus, *Rukkaia* dictus a sono. *Raii* syn. quad. 215.

Sciurus macrourus, long-tailed Squirrel. *Ind. Zool. tab. i.*

S with ears tufted with black: nose flesh-colored: cheeks, legs, and belly, of a pale yellow: between the ears a yellow spot: forehead, back, sides, haunches, black: cheeks marked with a bifurcated stroke of black; under side red: tail twice as long as the body, of a light grey, and very bushy: the part next the body quite furrounded with hair: on the rest the hairs are separated, and lie flat. Is thrice the size of the *European* squirrel.

Inhabits *Ceylon*: is called there *Dandoelana*: also *Roekeda*, from the noise it makes.

331. ABYSSINIAN.

S with a round flesh-colored nose: hair on the upper part of the body of a rusty black: tail a foot and a half long: belly and fore feet grey: soles of the feet flesh-colored. Thrice the size of an *European* squirrel.

Described from *Thevenot**, who says it was bought at *Moco* from an *Abyssinian*; that it was very good-natured, and sportive like a squirrel; would eat any thing except flesh, and would crack the hardest almonds. A variety of the above?

* *Voyage des Indes Orientales*, v. 34.

Sciurus maximus. Gmelin Lin. i. 149. Grand Ecureuil. Sonnerat, voy. ii. 139.

332. MALABAR.

S. with short tufted ears: five toes to each foot: instead of a thumb to the hind foot, is a short excrescence, with a flat nail; all the other nails strong and crooked: tail very full of hair, and as long as the body: hair long, of a reddish color, reflecting gold; a beard of the same begins under each ear, and turns towards the body: all the hind part of the body and tail black: is of the size of a cat.

Inhabits the mountains of *Cardomone* which form part of the *Gauts*: is very fond of the milk of the coco nut, which it will pierce and suck out on the tree. Its cry is sharp and piercing.

PLACE.

Sonnerat voy. ii. 140.

333. GINGI.

S. of a dirty grey color; brightest on the belly: eyes encompassed with a white circle: on each side of the belly is a white line which extends along the shoulders and thighs: tail black: rather larger than the *European* kind.

Inhabits *Gingi*.

Sonnerat,

334. AYE. AYE.

Sonnerat, ii. 142. tab. lxxxviii.

S. with large broad ears, smooth, shining, and with several long hairs scattered over them: fur soft and fine; of a tawny white, intermixed with some long black hairs: the tail is very bushy, covered with long hairs, black at their ends, white at their bottoms: five toes to each foot: the two joints of the middle finger of the fore feet very slender; the thumb of the hind foot furnished with a flat nail.

SIZE. Length eighteen inches; tail of the same length: burrows under ground: goes out only in the night: the eyes fixed: is very slothful, and of gentle manners: very fearful: much inclined to sleep; and rests with its head between its legs.

PLACE. Inhabits *Madagascar*: is a very rare animal: takes its name from its cry, the note of astonishment of the natives of that island.

335. JAVAN.

S. black on the upper part of the body; of a light brown on the lower: end of the tail black: on the thumb a round nail.


This brief account leaves me uncertain whether this is not also a variety.

Inhabits *Java*: discovered by Mr. *Sparman*. *Memoirs society at Gottingen*. Dr. *PALLAS*.

S. with



No. 334.



S. with tufted ears: head, back, sides, upper part of the legs and thighs, and tail, of a dull purple: the lower part of the legs, and thighs, and the belly, yellow: end of the tail orange: length, from nose to tail, near sixteen inches; tail seventeen. 336. BOMBAY.

Inhabits *Bombay*. Described from a stuffed skin in Doctor *Hunter's* cabinet.

This species extends to *Balisère*, the opposite part of the peninsula of *Indoستان*.

M. de la Cépède * gives the description and figure of a large squirrel which agrees so much with this, that I suspect it to be only a variety. He says on one front of the face is a chestnut spot, surrounded with orange: in other respects, there is much agreement, only he makes no mention of the orange at the end of the tail.

Sciurus Erythræus. Pallas Nov. sp. fasc. i. 377. Miller's plates, tab. xlvi.

337. RUDDY.

S. with the ears slightly tufted: color above yellow, mixed with dusky: below of a blood red inclining to tawny: tail slender; of the same color, marked lengthways with a black stripe.

* See *M. de Buffon*, Suppl. vi. 254, tab. lxii.

Four toes on the fore feet; with a remarkable protuberance instead of a thumb: five toes on the hind.

Rather larger in size than a common squirrel.

Inhabits *India*.

338. GREY.

Grey Squirrel. *Jesselyn's voy. Catechy Carolina*, ii. 74. *Smith's voy.* 27. *Kalm's voy.* 95, 310.

Fox Squirrel. *Larson's Carolina*, 124. *Sciurus cinereus Virginianus major. Raii syn. quad.* 215.

Sciurus cinereus. Lin. syst. 86.

Sciurus cinereus. Auriculis ex albo flavicantibus. Brisson quad. 107.

Le Petit-Gris. *De Buffon*, x. 116. *tab. xxv. LEV. Mus.*

S. with plain ears: hair of a dull grey color, mixed with black, and often tinged with dirty yellow: belly and insides of the legs white: tail long, bushy, grey, and striped with black. Size of a half-grown rabbit.

Inhabits the woods of *North America*, *Peru**, and *Chili*†; are very numerous in *North America*; do incredible damage to the plantations of *Maya*; run up the stalks, and eat the young ears; descend in vast flocks from the mountains, and join those that inhabit the lower parts; are proscribed by the provinces, and a reward of three pence *per* head for every one that is killed; such a

* *Chinchillas* are small beasts, like squirrels, with wonderful smoothe and soft skins, which they wear as a healthfull thing to comfort the stomacke; they make coverings and rugs of the haire of these *Chinchillas*, which are found on the *Sierre of Peru. Acosta in Purchas's Pilg.* iii. 966.

† *Ovalle*, in his history of *Chile*, says, that the grey or ash-color'd squirrels, of the valley of *Guasco*, are valuable for the furs. *Clurebill's Coll.* vol. iii. 44.

number



1. Hudson's Bay Squirrel - N^o 341.
 2. Black - N^o 339. 3. Grey - N^o 338.



number was destroyed one year, that *Pensylvania* alone paid in rewards 8000*l.* of its currency.

Make their nests in hollow trees, with moss, straw, wool, &c. Feed on the mayz in the season, and on pine-cones, acorns, and masts of all kinds. Form holes under ground, and there deposit a large stock of winter provision. Descend from the trees and visit their magazines when in want of meat; are particularly busy at the approach of bad weather; during the cold season keep in their nests for several days together; seldom leap from tree to tree, only run up and down the bodies; their hoards often destroyed by swine; when covered with deep snow, the squirrels often perish for want of food; are not easily shot, nimbly changing their place, when they see the gun levelled; have the actions of the common squirrel; easily tamed; their flesh esteemed very delicate. The furs which are imported under the name of *petit-gris* are valuable, and used as linings to cloaks.

Quahtechalotl-thlitic. *Hernandez Mex.* L'Ecureuil noir. *Briffon quad.* 105.

582. *Hernandez Nov. Hisp.* 8.

Black Squirrel. *Catesby Car.* ii. 73.

Sciurus niger. *Lin. syst.* 86. *LEV. MUS.*

339. BLACK.

S. with plain ears: sometimes wholly black, but often marked with white on the nose, the neck, or end of the tail: the tail shorter than that of the former: the body equal.

Inhabits the *North of Asia*, *North America*, and *Mexico*. I should have placed it as a variety of the last species, did not Mr.

VOL. II.

U

Catesby

Catechy expressly say, that it breeds and associates in separate troops; is equally numerous with the former; commits as great ravages among the *Mays*; makes its nest in the same manner, and forms, like them, magazines for winter food.

A squirrel of a most beautiful shining black color, is found at the *Pulo Condore*, in lat. 8. 40.

β. SQUIRREL, with plain ears: coarse fur, mixed with dirty white, and black, but varies to white: throat, and inside of the legs and thighs, black: tail much shorter than those of squirrels usually are: of a dull yellow color, mixed with black: body of the size of the grey squirrel. LEV. MUS.

Inhabits *Virginia*; described from Mr. *Knaplan*'s collection, who told me that the planters called it the *Cat Squirrel*.

340. MADAGAS-
CAR.

S. with plain ears: color of the face, back, sides, tail, and outside of the limbs, of a dark glossy black: ears, end of the nose, cheeks, and all the under side of the limbs, yellowish white. The length of this species from the tip of the nose to the origin of the tail, is about eighteen inches: the tail is longer than the body, slender, and ends in a point.

Inhabits *Madagascar*: described by M. *de la Cépède*, in his supplement to M. *de Buffon*, vii. 256. tab. lxxiii.

S. with plain ears: smaller than the *European*: marked along the middle of the back with a ferruginous line from head to tail: the sides paler: belly of a pale ash-color, mottled with black: tail not so long, or so full of hair, as the common kind; of a ferruginous color, barred with black, and towards the end is a broader band of the same color. LEV. MUS.

341. HUDSON'S
BAY.

Inhabits the pine-forests about the *Bay*, and *Terra de Labrador*.

α. CAROLINA * SQUIRREL, with the head, back, and sides grey, white, and rust-colored intermixed: belly white, divided from the colors of the sides by a ferruginous line: lower part of the legs red: tail brown, mixed with black, and edged with white.

These are rather lesser than the *European* squirrels: vary in the colors: in most the grey predominates.

Quauhcolotlquapachtli. *Hernandez Nov. Hisp.* 8.
Le Coquallin. *De Buffon*, xliii. 109. tab. xliii.

342. VARIED.

S. with plain ears: upper part of the body varied with black, white and brown: the belly tawny†: twice the size of the common squirrel.

* Lesser Grey Squirrel of the old edition.

† Called by the *Indians*, *Cozticotequallin*, or Yellow Belly.

U 2

Inhabits

Inhabits *Mexico*: lives under ground, where it brings forth its young, and lays in a flock of winter food: lives on mayz: is never to be tamed.

These probably vary in size: I have seen one that seemed to be of this species, but not superior in size to the common squirrel: the colors were brown, orange, and cinereous: the belly orange.

343. FAIR.

Sciurus flavus. Sc. auriculis subrotundis, pedibus pentadactylis, corpore luteo. *Lin. ffl.* 86. *Amen. Acad.* 1. 561.

S. with the body and tail of a flaxen color: of a very small size, with plain round ears, and rounded tail.

Inhabits the woods near *Amadabad*, the capital of *Guzarat*, in great abundance, leaping from tree to tree *. *Linnaeus* says it is an inhabitant of *South America*.

344. BRASILIAN.

Sciurus Brasiliensis? *Marcgrave Brasl.*

330.

Sciurus coloris ex flavo et fusco mixti tæniis in lateribus albis. *Brisson quad.*

107.

Sciurus æstuans. Sc. griseus, subtus flavescens. *Lin. ffl.* 88.

S. with plain ears, and rounded tail: head, body, and sides, covered with soft dusky hairs, tipped with yellow: tail rounded: the hairs annulated with black and yellow: throat cinere-

• L'Ecureuil blond. *Della Valle*, p. 84.

OUS:

ous: inside of the legs, and the belly, yellow: the belly divided lengthways with a white line; which begins on the breast, is interrupted for a small space in the middle, and is then continued to the tail: length, from nose to tail, eight inches one quarter: tail ten.

Inhabits *Brazil* and *Guiana*. Mr. *Vandeck*, captain of a man of war in the *Portuguese* service, who procured them from their settlements in *S. America*, favored me with two.

Tlalmototli. *Hernandez Nov. Hisp.* 9.
Sciurus rarissimus ex Nov. Hispania. *Seb.*

Mus. i. 76. *tab.* xlvii. fig. 2, 3. *Briffon*
quad. 103.

345. MEXICAN.

S. of a mouse-color: the male marked on the back with seven white lines, which extend along the tail; the female, with only five: the tail of the male divided into four parts at the end: perhaps accidentally: its *scrotum* pendulous, like a goat's.

Inhabits *New Spain*. *Seba*, in *tab.* xlviii. *fig.* 5. has the figure of another, of an uniform color, distinguished also by its vast *scrotum*.

Mustela Africana. *Clus. Exot.* 112. *Raii*
syn. quad. 2162

Sciurus palmarum. Sc. subgriseus striis
tribus flavicantibus, caudaque albo ni-
groque lineata. *Lin. syst.* 86.

Sc. palmarum. Sc. coloris ex rufo et ni-
gro mixti, tenuis in dorso flavicantibus.
Briffon quad. 109.

Le Palmiste. *De Buffon*, x. 126. *tab.*
xxvi.

346. PALM.

S. with plain ears: an obscure pale yellow stripe on the middle of the back, another on each side, a third on each side of the

the belly; the two last at times very faint: rest of the hair on the sides, back, and head, black and red, very closely mixed; that on the thighs and legs more red: belly, pale yellow: hair on the tail does not lie flat, but encircles it; is coarse, and of a dirty yellow, barred with black. Authors describe this kind with only three stripes: this had five, so possibly they vary.

Governor *Loten* did me the favor of informing me that it lived much in the *Coco* trees, and was very fond of the *sury*, or palm-wine, which is procured from the tree; from which it obtained, among the *Indians*, the name of *Suricatsjé*, or the *little cat* of the *Sury* *.

According to *Clusius* and Mr. *Ray*, this species does not erect its tail like other squirrels, but has the faculty of expanding it sideways.

247. WHITE-
STRIPED.

Æ. BARBARY. *Sciurus getulus. Cail quad. 84. Brisson quad. 109.*
opusc. 77. Gesner quad. 847. Barbarian squirrel. *Edw. 198.*
Sc. getulus. Sc. fuscus striis quatuor albis
longitudinalibus. *Lin. syst. 87. Klein tab. xxvii.*
Le Barbarelique. De Buffon, x. 126.

S. with full black eyes and white orbits: head, body, feet, and tail, cinereous, inclining to red: lightest on the legs: sides, marked lengthways with two white stripes: belly white: tail bushy, marked regularly with shades of black, one beneath the other: size of the common squirrel.

* See the process of obtaining this liquor in *Rumphius's herbarium Amboinense*, vol. i. p. 5. The tree is engraved in *tab. i. ii.*

Both these squirrels inhabit *Barbary* and other hot countries : live in trees, especially *palms*, from which one takes its name.

THIS species resembles much the common squirrel, but is lighter colored, and has a yellow line extending along the sides, from leg to leg.

348. PLANTANE.

Common in *Java* and *Prince's island*; is called by the *Malayes*, *Ba-djing*; lives much on *Plantanes*; is very shy; retreats at the sight of mankind, and clatters over the dry leaves of the *Pitang* or *Plantanes* with vast noise. It also is common on the *tamarind* trees.

A. with membranes from fore leg to hind leg.

Sciurus Sagitta. Sc. hypochondriis prolixis volitans, cauda plano-pinnata lanceolata. *Lin. syst.* 88.

Sciurus petaurita. *Pallas Mysel. Zool.* 54. tab. vi.

Sciurus maximus volans, seu felis volans. Sc. castanei coloris, in parte corporis

superiore, in inferiore vero eximie flavescens; cute ab anticis cruribus ad postica membranæ in modum extensa volans. *Briffon quad.* 12.

349. SAILING.

Le Taguan ou grand Ecureuil volant. *De Buffon*, Suppl. iii. 150. tab. xxi. *Mus. Roy. Society* *.

S with a small rounded head : cloven upper lip : small blunt ears : two small warts at the outmost corner of each eye,

* Where there is the skin of one in fine preservation.

with

with hairs growing out of them: neck short: four toes on the fore feet; and instead of a thumb, a slender bone, two inches and a half long, lodged under the lateral membrane, serving to stretch it out: from thence to the hind legs extends the membrane, which is broad, and a continuation of the skin of the sides and belly; the membrane extends along the fore legs, and stretches out near the joint in a winged form: five toes on the hind feet, and on all the toes sharp, compressed, bent claws: tail covered with long hairs, disposed horizontally: color of the head, body, and tail, a bright bay; in some parts inclining to orange: breast and belly of a yellowish white: length, from nose to tail, eighteen inches; tail fifteen.

Inhabits *Java**, and others of the *Indian* islands: leaps from tree to tree as if it flew: will catch hold of the boughs† with the tail. Differs in size: that described by *Linnaeus* was the size of our squirrel: that killed by Sir *Edward Michelbourne*, in one of the *Indian* isles, was greater than a hare. *Nieuboff* describes this species under the name of the Flying Cat, and says the back is black: he has given two very good figures of it; one in his frontispiece, the other in the page he describes it in‡.

* *Hamilton's voy.* ii. 131.

† *Sir Edward Michelbourne's voy. in Purchas's Pilgrim.* i. 134.

‡ *Churchill's coll.* ii. 354.



Pileated Squirrel — N. 319.



Greater Flying Squirrel. *Ph. Tr.* lxii. 379.

350. SEVERN RIVER:

S. with back and sides of a deep cinereous color at the bottom; end ferruginous: under side of the body of a yellowish white; the hair every where long and full: tail covered with long hairs, disposed in a less flat way than those of the *European* kind; brown on the upper part, darker at the end, yellowish beneath the skin: the instrument of flying disposed from leg to leg; but does not border the fore-legs.

Size equal to the *European* squirrel.

Inhabits the southern parts of *Hudson's Bay*, about *Severn* river. *Mus. Roy. Society.*

SIZE.

PLACE.

Asapanick. Smith's Virginia, 27. *Jes-
seyn's voy.* 86. *De Laet*, 88.
Sciurus Americanus volans. Raii syn. quad.
215. *Lawson's Carolina*, 124. *Catesby*
Carolina, ii. 76, 77. *Edw.* 191. *Kalm*,

i. 321. *tab. i. De Pratz.* ii. 69.
Sciurus volans. Sc. hypochondriis prolaxis
volitans, cauda rotundata. Lin. syst. 88.
Sciurus volans. Brisson quad. 110. iii. No.
12. *LEV. MUS.*

351. FLYING.

S. with round naked ears: full black eyes: a lateral membrane from fore to hind legs: the fore legs for the most part clear of the membrane: tail with long hairs disposed horizontally, longest in the middle, and ending in a point: color above, a brownish ash: beneath white, tinged with yellow. Much less than the common squirrel.

VOL. II.

X

Inhabits

Inhabits *North America* and *New Spain* *: lives in hollow trees : sleeps in the day ; during the night very lively ; is gregarious, numbers being found in one tree : leaps from bough to bough sometimes at the distance of ten yards : this action improperly called *flying*, for the animal cannot go in any other direction than forward ; and even then cannot keep an even line, but sinks considerably before it can reach the place it aims at : sensible of this, the squirrel mounts the higher, in proportion to the distance it wishes to reach : when it would leap, it stretches out the fore legs, and extending the membranes, becomes specifically lighter than it would otherwise be ; and thus is enabled to spring further than other squirrels that have not this apparatus. When numbers leap at a time, they seem like leaves blown off by the wind. Their food the same as the other *American* squirrels : are easily tamed : bring three or four young at a time.

352. NORFOLK-
ISLE.

Stockdale's Bot. Bay, 151. White, 288.

S. with very short ears, almost hid in the fur : color very much resembling that of the *American* grey squirrel : a black line extends from the head along the middle of the back to the tail : the flying membrane black, edged with white : two thirds of the tail are of an elegant ash-color ; the rest black : size of the *American* grey squirrel.

Inhabits *Norfolk isle*.

* Where it is called *Quimichpatlan*. *Hernandez, Nov. Hisp.* 8.

In

In the isle of *Pulo Condore* is a flying squirrel striped with brown and white: possibly a new species.

Sciurus Virginianus volans. Seb. Mus. i. tab. xlv. Briffon quad. iii.
Mus volans. Lin. syst. 85.

353. HOODED.

S with the lateral membrane beginning at the chin and ears, and extending like the former from fore to hind leg: reddish above; cinereous, tinged with yellow, beneath: ears large and oval.

Inhabits *Virginia*, according to *Seba*; who is the only author who has described it. *Linnaeus's* synonyms, from *Ray* and *Edwards*, erroneous.

Mus Ponticus vel Scythicus. Gefner quad.
 743.

Sciurus Petaurista volans. Klein quad. 54.
 Flying squirrel. *Pb. Trans. abr. ix. 76.*
 tab. v.

Sciurus volans. Faun. suet. No. 38. Pal-
las, nov. sp. fasc. i. 355.

Sc. volans Sc. hypochondrii prolaxis

volitans, cauda rotundata. Lin. syst.
 88.

Sciurus Sibiricus volans. Briffon, 110.
 No. 13.

Le Poulatouche. De Buffon, x. 95. tab.
 xxii.

Quadrupes volatilis Russiæ. Com. acad.
Petrop. v. 218. LEV. Mus.

354. EUROPEAN
 FL. SQ.

S with naked ears, indented on the exterior side: full eyes: eyelids bordered with black: membranes extend to the very base of the fore feet, and form a large wing on the exterior side: tail full of hair, and round at the end: color of the

upper part of the body a fine grey, like that on a gull's back ; lower part of a pure white.

SIZE. From nose to tail four inches and a quarter ; of the tail to the tip of the hair, five.

PLACE. Inhabits *Finland* and *Lapland*, and the *Russian* dominions, from *Livonia* to the river *Kolyma* or *Kowyma*, in the N. E. part of *Siberia*, and is common in all the mountainous wooded tracts of that cold region : lives usually on birch-tree buds and fructifications, and on the cones of the pines and cedars : is not gregarious, and leads a solitary life, and wanders about even in winter : lives in hollow trees, and makes its nest in the moss of birch-trees : when at rest, it flings its tail over its back ; but in leaping, extends it.

NAMES. The *Germans* call it *Könige der Grauzerke**, or King of the Squirrels ; the *Russians*, *Polatucha*, and *Letaga* ; the *Poles*, *Wiejviorka* *Lataiaca*.

* *Klein*.

Two cutting teeth in each jaw.
Four toes before: five behind.
Naked ears.
Long tail, covered with hair.

XXXI.
DORMOUSE.

Moufe squirrel. *Joffelyn's voy.* 86.
Ground fquirrel. *Lawfon Carolina*, 124.
Catefby Carolina, ii. 75. *Edw.* 181.
Kalm, i. 327. *tab.* i.
Sciurus Lifteri. Raii fyn. quad. 216.
Sciurus minor virgatus. Nov. Com. Petrop.
v. 344.
Boern-doeskie. Le Brun, voy. Mofcov. ii.
342.

Sciurus striatus. Sc. flavus ftriis quinque
fulcis longitudinalibus. Lin. fyft. 87.
Klein quad. 53. *Pallas nov. fp. fafc.* i.
378.
Sciurus Carolinenfis. Briffon quat.
Le Suiffe. De Buffon, x. 126. *tab.* xxviii.
Charlevoix Nouv. France, v. 198. *LEV.*
Mus.

355. STRIPED.

D. with plain ears: ridge of the back marked with a black
freak: each fide with a pale yellow ftripe, bounded
above and below with a line of black: head, body, and tail, of
a reddifh brown; the tail the darkeft: breaft and belly white:
nofe and feet pale red: eyes full.

Inhabits the north of *Afia*, beginning about the river *Kama*,
and grows more and more frequent in the woody parts of *Sibe-*
ria; but found in the greateft abundance in the forefts of *North*
America: they never run up trees except purfued, and find no
other means of efcaping: they burrow, and form their habitations
under ground with two entrances, that they may get accefs to the
one, in cafe the other is ftopped up. Their retreats are formed with
great fkill, in form of a long gallery, with branches on each fide,
each of which terminates in an enlarged chamber, as a magazine

PLACE.

MANNERS.

MAGAZINES.

to store their winter provision in; in one they lodge the acorns, in another the *mayz*, in a third the hickery nuts, and in the last, their favorite food, the *chinquapin* chestnut. They very seldom stir out during winter, at least as long as their provisions last; but if that fails, they will dig into cellars where apples are kept, or barns where *mayz* is stored, and do a great deal of mischief; but at that time the cat destroys great numbers, and is as great an enemy to them as to mice.

During the *mayz* harvest, these animals are very busy in biting off the ears, and filling their mouths so full with the corn, that their cheeks are quite distended, having pouches in their jaws like the *hamster*. It is observable, that they give great preference to certain food; for if, after filling their mouths with rye, they happen to meet with wheat, they sling away the first, that they may indulge in the last. They are very wild, bite severely, and are scarcely ever tamed: the skins are of little use; but are sometimes brought over to line cloaks.

356. FAT.

Glis. *Gesner quad.* 550. *Raii syn. quad.* 229. *Sciurus* Glis. Sc. canus subtus albidus.
 Glis vulgaris. *Klein quad.* 56. *Lin. syst.* 87.
 Glis supra obscurè cinereus, infra ex albo Le Loir. *De Buffon*, viii. 158. *tab.* xxiv.
 cinerescens. *Briffon quad.* 113. *Mus* Glis. *Pallas nov. sp. fasc.* i. 88.

D. with thin naked ears: body covered with soft ash-colored hair: belly whitish: tail full of long hair: from nose to tail, near six inches; tail four and a half: thicker in the body than the squirrel.

Inhabits *France* and the south of *Europe*. Is found in the woods in the south-west parts of *Russia*, and was discovered by
 Doctor

Doctor *Pallas* in the rocky caverns about the rivers *Samara* and *Volga*. The late Doctor *Kramer* favored me with one from *Austria*. Lives in trees, and leaps from bough to bough: feeds on fruits and acorns: lodges in the hollows of trees: remains in a torpid state during winter, and grows very fat,

*Tota mihi dormitur hyems, et pinguior illo
Tempore sum, quo me nil nisi somnus alit*.*

Was esteemed a great delicacy by the *Romans*, who had their *Gliraria*, places constructed to keep and feed them in. I think that the *Italians* at present eat them.

Mus avellanarum major. *Gesner quad.* 735. *Mus quercinus.* *M. cauda elongata pilosa,* macula nigra sub oculos. *Lin. f. f.*
Greater Dormouse, or Sleeper. Raii syn. 84.
quad. 219.
Glis supra obscure cinereus, infra ex albo cinerefcens, macula ad oculos nigra. *Le Lerot. De Buffon, viii. 181. tab. xxv.*
Briffon quad. 114. *Mus nitedula. Pullas, nov. sp. fasc. i. 88.*

357. GARDEN.

D. with the eyes surrounded with a large spot of black, reaching to the base of the ears, and another behind the ears: head and whole body of a tawny color: the throat and whole under side of the body white, tinged with yellow: the tail long: the hairs at the beginning very short; at the end bushy: length, from nose to tail, not five inches: the tail four.

Inhabits *France* and the south of *Europe*: is found in magpies nests and hollow trees about the *Volga*, and other temperate

* *Martial Epig. Lib. xiii. Ep. 59.*

and

and southern parts of the *Russian* dominions. Neither this nor the former species extend beyond the *Uralian* mountains: infests gardens, and is very destructive to fruits of all kind: is particularly fond of peaches: lodges in holes in the walls: brings five or six young at a time: like the former, remains torpid during winter: has a strong smell, like a rat.

358. *DEGUS.**Sciurus Degus. Molina Chili, 284.*

D. of a dull white color, and with a blackish line cross the shoulders, reaching to the elbows: the tail ending in a tuft: ears rounded: larger than the common rat.

MANNERS.

Inhabits *Chili*, and lives under ground, near the hedges and bushes; and forms its retreat into various galleries communicating with each other: feeds on roots and fruits, and lays up a large provision of them for winter food. Is not torpid during that season like our dormouse.

359. *COMMON.*

Mus avellanarum minor, the Dormouse or Sleeper. *Raii syn. quad.* 220.

Rothe Wald Maufs. *Kramer Austria*, 317.

Glis supra rufus, infra albicans. *Briffon quad.*

Mus avellanarius. *M. cauda elongata pilosa*, corpore rufo, gula albicante, pol-

licibus posticis muticis. *Lin. syst.* 83. *Faun. suec.* No. 35. *Pallas nov. sp. fasc.*

i. 89.

Le Mulcardin. *De Buffon*, viii. 193. *tab. xxvi.*

Dormouse. *Edw.* 266. *Br. Zool.* i. 95. *LEV. MUS.*

D. with round naked ears: full black eyes: body of a tawny red: throat white: size of a mouse, but plumper: tail two inches

inches and a half long, and pretty hairy, especially towards the end.

Inhabits *Europe*: lives in thick hedges: makes its nest in the hollow of a low tree, or in a thick bush near the bottom, of grass, moss, or dead leaves: brings three or four young at a time: seldom appears far from its retreat: forms magazines of nuts: eats its food sitting up, like a squirrel: at approach of winter, retires and rolls itself up, lying torpid: sometimes in a warm day revives, takes a little food, and relapses into its former state.

D. with a flat head, obtuse nose, eyes full and black, upper lip bifid.

360. EARLESS.

Auricles very minute, scarcely apparent: long whiskers.

Head, back, sides, and front of the fore legs, pale ferruginous, mixed with black: from shoulder to hind parts, on each side, a white line: above each eye another: belly and feet of a dirty white.

Tail black in the middle; hoary on the sides.

Toes long and distinct: the knob on the fore feet large: claws very long.

Hind legs black behind, and naked.

Size of a common squirrel, but much broader and flatter.

800 miles above the *Cape of Good Hope*, about the mountain *Sneeberg*. Communicated by Sir *Joseph Banks*.

SIZE.

PLACE.

VOL. II.

Y

Never

MANNERS.

Never climbs trees: burrows, feeds on bulbous roots, and is particularly fond of potatoes: walks often on its hind feet; and often lies flat on its belly: very tame, and never offers to bite: frequently flirts up its tail: makes a warm nest, and forms in it a round hole, in which it lodges, and pulls to the orifice a quantity of materials, in order to close it: keeps sometimes in its retreat for three entire days.

361. GILT-TAIL.

Le Lerot: a queue doree. *Allamand Supplém.* iv. 164. tab. lxxvii.

SIZE.

PLACE.

D. with short broad ears, great whiskers, the face marked lengthways with a gold color line extending from the nose to the space between the ears. The rest of the head and whole body and beginning of the tail are a purplish chestnut color, the remaining half of the tail is black: the rest of a beautiful gold color. The tail is thick about the base.

Length from nose to tail is five inches; of the tail six and nine lines.


Inhabits *Surinam*. Lives on fruits and climbs up the trees.

GUERLINGUETS.

*M. de la Cépède** gives us the description of two species of animals, which he calls *Guerlinguets*. He denies that they are true squirrels: the ears are naked, and the tail grows taper, yet is covered with long hair, but by no means disposed like

* *Supplém. &c.* vii. 261. tab. lxxv. lxxvi.

that



Gilt tail Dormouse — . 1361.



that on the tail of the squirrel: they may come into this genus; at least let them remain here till we are better informed.

The larger is between seven and eight inches long, exclusive of the tail: the tail is of equal length: the hair on the body is very short, and at its extremity a bright bay. The tail is rayed with brown and tawny.

362. GREATER.

The lesser is little more than four inches long: the tail little more than three: the body, legs, and tail, are clouded with olive and ash-color: on the face, lower part of the belly, and sides of the legs are tawny.

363. LESSER.

XXXII.
JERBOA.

Two cutting teeth in each jaw.

Two very short fore legs: two very long hind legs, resembling those of cloven-footed water-fowl.

Very long tail, tufted at the end.

364. ÆGYPTIAN.

*Mus Juvæ. Theophr. apusc. 295. Ælian hist. an. lib. xv. c. 26.**Mus bipes. Plinii lib. x. c. 65. Textura's Travels, 21.**Gerbua. Edw. 219. Plaised's journal, 59.**Mus jaculus. M. cauda elongata floccosa,**palms subpentadactylis, femoribus longissimis, brachiis brevissimis. Lin. syst. 85. Hasselquist itin. 198.**Le Jerbo. De Buffon, xiii. 141.**Mus sagitta. Pallas nov. sp. fasc. i. 306. tab. xxi.*

J with thin, erect, and broad ears: full and dark eyes: long whiskers: fore legs an inch long; five toes on each; the inner, or thumb, scarce apparent; but that, as well as the rest, furnished with a sharp claw: hind legs two inches and a quarter long, thin, covered with short hair, and exactly resembling those of a bird; three toes on each, covered above and below with hair; the middle toe the longest; on each a pretty long sharp claw: length, from nose to tail, seven inches and one quarter: tail ten inches, terminated with a thick black tuft of hair; the tip white; the rest of the tail covered with very short coarse hair: the upper part of the body thin, or compressed sideways: the part about the rump and loins large: the head, back, sides, and thighs, covered with long hair, ash-colored at the bottom, pale tawny at the ends: breast and belly whitish: across the upper

per part of the thighs is an obscure dusky band: the hair long and soft.

Inhabits *Ægypt*, *Barbary*, *Palestine*, the deserts between *Basfora* and *Aleppo*, the sandy tracts between the *Don* and *Volga*, the hills south of the *Irtish*, from fort *Janiyschera* to the *seven palaces*, where the *Altaic* mountains begin: as singular in its motions as in its form: always stands on its hind feet; the fore feet performing the office of hands: runs fast; and when pursued, jumps five or six feet from the ground: burrows like rabbits: keeps close in the day: sleeps rolled up: lively during night: when taken, emits a plaintive feeble note: feeds on vegetables: has great strength in its fore feet. Two, which I saw living in *London*, burrowed almost through the brick wall of the room they were in; came out of their hole at night for food, and when caught, were much fatter and sleeker than when confined to their box.

This is the *Daman Israel*, or the *Lamb of the Israelites* of the *Arabs*, and is supposed to be the *Saphan**, the coney of *HOLY WRIT*: our rabbit being unknown in the *Holy Land*. Dr. *Shaw* met with this species on mount *Libanus*, and distinguishes it from the next species†. It is also the mouse of *Ijaiab*‡, *Achbar* in the original signifying a male *Jerboa*.

This and the following species, which is found to extend to the

PLACE.

* *Bochart* displays a vast deal of learning on the subject. *Vide Hierozoicon, lib. iii. c. 33. p. 1001.*

† *Travels*, 376.

‡ *Chap. lxvi. 17. Bochart, 1015.* This animal was a forbidden food with the *Israelites*.

colder

colder regions, on any approach of cold grow torpid, and remain so till they are revived by a change of weather. *Pallas* calls this class the *Species Lethargica*.

365. SIBERIAN.

Cuniculus pumilio saliens cauda longissima. Nov. Com. Petrop. v. 351. tab. ix. fig. 1.
Cuniculus pumilio saliens, cauda anomala longissima. Briffon quad. 103.

Dipus Jaculus. Gm. Lin. 157.

Flying hare. Strahlenberg's hist. Russ. 370.

Mus jaculus. Pallas nov. sp. fasc. 1, 275. tab. xx. Mus. LEV.

α. GREAT.

NOSE truncated; end edged with white: lower teeth slender; twice as long as the upper.

Ears large, pointed, tips with white, naked within: hairs on the back tawny, of a dark grey beneath, very soft: legs and whole under side of the body white: half the tail next to the body covered with short whitish hairs; from thence, with long black hairs; the end has a large white feathered tuft an inch long.

Five toes on the fore feet; the toe without a nail.

On the hind legs, an inch above the feet, are two long toes armed with nails: the back part of the legs naked.

Length eight inches and a half; of the tail ten.

PLACE.

This variety is nowhere very frequent, but is chiefly found from the *Caspian* sea to the river *Irtish*.

β. MIDDLE.

Of the size of a rat: of the color of the former, except that the rump on each side is crossed with a white line.

This middle species is found only in the eastern deserts of *Siberia* and *Tartary*, beyond lake *Baikal*; also in *Barbary** and *Syria*†, and even as far as *India*‡.

* *Shaw's Travels.*

† *Hayn's Tesoro Brit. ii. p. and tab. 124.*

‡ *Pallas.*
Differs



Differs from the *Great*, in wanting the white circle round the nose, in having a less tuft to the tail, and the end just tipped with white: agrees entirely in form; but is far inferior in size to even the *Middle*. Inhabits the same places with the *Great*.

γ. PYGMY.

These three agree in manners: burrow in hard ground, clay or indurated mud: not only in high and dry spots, but even in low and salt places. They dig their holes very speedily, not only with their fore feet but with their teeth, and sling the earth back, with their hind feet, so as to form a heap at the entrance. The burrows are many yards long, and run obliquely and winding, but not above half a yard deep below the surface. They end in a large space or nest, the receptacle of the purest herbs. They have usually but one entrance; yet by a wonderful sagacity they work from their nest another passage to within a very small space of the surface, which in case of necessity they can burst through, and so escape.

MANNERS.

It is singular, that an animal of a very chilly nature, should keep within its hole the whole day, and wander about only in the night.

They sleep rolled up, with their head between their thighs: and when kept in a stove, and taken suddenly out, they seem quite stupified, and for a time scarcely find the use of their limbs: perhaps this arises from an excess of heat; for when an attempt is made to take them out of their burrows, they are quickly alarmed on the noise of digging, and attempt their escape. At sun-set they come out of their holes, clear them of the filth, and keep abroad till the sun has drawn up the dews from the earth. On approach of any danger, they immediately take to flight, with leaps a fathom in height, and so swiftly that a
man

GREAT LEAPS.

man well mounted can hardly overtake them. They spring so nimbly, that it is impossible to see their feet touch the ground. They do not go strait forwards, but turn here and there, till they gain a burrow, whether it is their own, or that of another. In leaping, they carry their tails stretched out: in standing, or going or walking, they carry them in form of an S, the lower part touching the ground, so that it seems a director in their motions. When surprized, they will sometimes go on all fours, but soon recover their attitude of standing on their hind legs like a bird: even when undisturbed, they use the former attitude; then rise erect, listen, and hop about like a crow. In digging or eating they drop on their fore legs: but in the last action will often sit up and eat like a squirrel.

EASILY TAMED.

They are easily made tame: seek always a warm corner: foretell cold or bad weather by wrapping themselves close up in hay; and those which are at liberty stop up the mouths of their burrows.

FOOD.

In a wild state they are particularly fond of the roots of tulips: live much on oleraceous plants: the small stature of the *pygmy* kind is attributed to their feeding on saline plants. Those of the middle size, which live beyond the lake *Baikal*, live on the bulbs of the *Lilium Pomponium*, and they gnaw the twigs of the *Robinia Carugana*. When confined, they will not refuse raw meat, and the entrails of fowls.

They are the prey of all lesser rapacious beasts. The *Arabs*, who are forbidden all other kinds of mice, esteem these the greatest delicacies: as those people often are disappointed in digging after them, they have this proverb, “To buy a hole instead of a *Jerboa*.”

The *Mongols* have a notion that they suck the sheep: certain it is, they are during night very frequent among the flocks, which they disturb by their leaps.

The *Mongols* call this animal *Alagh-Daagha*. *Alagh* signifies variegated, *Daagha*, a foal. The *Calmuks* call it *Jalma*: the great sort they style *Morin Jalma*, or the *Horse Jerboa*; the lesser sort, *Chein Jalma*, or the *Sheep*.

NAMES.

They breed often in the summer; in the southern parts, in the beginning of *May*: beyond *Baikal*, not till *June*. They bring perhaps eight at a time, as they have so many teats. They sleep the whole winter without nutriment. About *Afracan*, they will sometimes appear in a warm day in *February*: but return to their holes on the return of cold.

Animals of this genus were certainly the *two-footed mice*, and the *Egyptian mice*, of the ancients, which were said to walk on their hind legs; and use the fore instead of hands. These, with the plant *Silphium*, were used to denote the country of *Cyrene*, where both were found, as appears from the figures on a beautiful gold coin preserved by Mr. *Haym**, and which I have caused to be copied above the animal, in the plate.

Dipus sagitta. *Gm. Lin.* 158. *Pallas nov. sp.* 87, 206. tab. xxi. *Edw.* tab. 219.

366, ARROW.

J. with ears shorter and broader than the preceding: nose longer and less obtuse: toes before, three behind: coat thicker

* *Tesoro Brit.* ii. 124.

and longer; a white band from the base of the tail to the junction of the thighs with the body: length from the tip of the nose to the rump, little more than five inches; of the tail six.

Inhabits *Barbary*, and all the north of *Africa*, *Aegypt*, *Arabia*, and *Syria*; and lives in the sandy deserts.

367. CAPE.

Grand Gerbo. *Allamand de Buffon*, xv. *Dipus Cafer*. *Gm. Lin.* 159. *Miller's plates*.
118. *Journal Historique*, 59. *tab. xxxi.*

J with a short head: broad between the ears: mouth placed far below the upper jaw: lower very short: two great teeth in each: ears one-third shorter than those of the common rabbit, thin and transparent: eyes large: whiskers great.

Fore legs short, five toes on each, with a great protuberance next to the inner toe: claws of the fore toes crooked, and two-thirds longer than the toes themselves: claws of the hind toes short.

Color above tawny; cinereous below, mixed with long hairs pointed with black: two-thirds of the tail tawny, the rest black.

Length from nose to tail one foot two inches; of the tail near fifteen inches; the ears near three.

Inhabits the great mountains far north of the *Cape of Good Hope*. It is called by the *Hottentots*, *Aerdmannetje*; and by the *Dutch*, *Springen Haas*, or the *Jumping Hare*.

It is very strong; will leap twenty or thirty feet at a time: its voice a grunting: when it eats, sits upright, with the legs extended horizontally, and with a bent back: uses its fore feet to bring the food to its mouth; burrows with them, which it does
so

so expeditiously as quickly to bury itself. In sleeping, it sits with its knees separate, puts its head between its hind legs, and with the fore legs holds its ears over its eyes.

Mus longipes. *M. cauda elongata vestita*, palmis tetradactylis, plantis pentadactylis, femoribus longissimis. *Lin. Syst.* 84.

Mus cauda longa vestita, pedibus posticis longitudine corporis, flavis. *Mus. Ad. Fr.* 9.

368. TORRID.

J with naked oval ears: long whiskers: four toes on the fore feet: the hind feet the length of the body, thick, strong and thinly haired: five toes on each foot: scarcely any neck: tail the length of the body, with very little hair on it: color of the upper part of the body yellow; the lower white: size of a common mouse.

Inhabits, according to *Linnaeus*, the torrid zone*: mentioned by no other writer.

* *Habitat in torridis regionibus.*

XXXIII. RAT.

Two cutting teeth in each jaw.

Four toes before; five behind.

Very slender taper tail; naked, or very slightly haired.

* Jerboid.

369. CANADA.

R. with the upper jaw projecting far beyond the lower: upper cutting teeth deeply divided by a longitudinal furrow: ears small, and hid in the fur, and placed far back: the three middle hind toes very long; those on each side very short: color of the fur on the upper part of the head and body, light rust; towards the bottom of the fur iron grey: belly whitish: length from nose to the tip of the tail scarcely two inches: tail very slender; three inches and a half long.

MANNERS.

This animal inhabits the woods of *Canada*. Its hind legs have more of the *Jerboas*, than any of the rest of this genus; are very long: it goes upright on these, like the Jerboa; and its pace is leaping like that animal: is exceeding nimble, and is with difficulty caught, except it can be forced into the open grounds: is torpid during winter: wraps itself up like the dormouse, and coils up its long tail; previously rolling itself into a round ball of clay, which it forms for its winter retreat.

I am indebted to Col. *Davies*, of the artillery, for the sight and the account of this curious animal.

R. with

R. with a blunt nose: mouth placed far beneath: upper lip bifid: ears large, naked, rounded; fore legs short, furnished with four toes, and a tubercle instead of a thumb: hind legs long and naked, like some of the *Jerboas*: thumb short: toes long, slender, and distinct; the exterior the shortest.

370. LABRADOR.

The whole length of the animal eight inches: of which the tail is four and three quarters.

Color deep brown above, white beneath, separated the whole length on each side by a yellow line.

Inhabits *Hudson's Bay*, and the *Labradore* coast. Sent by Mr. *Graham*, and deposited in the Museum of the *Royal Society*.

PLACE.

Since I wrote the above, I find that Doctor *Pallas* has described this species under the title of *Mus Longipes* *. It inhabits also the sandy desert of *Naryn*, or *Ryn Pesky*, between the *Volga* and the *Taik*, near the *Caspian Sea*, in lat. $46\frac{1}{2}$. In this tract scarcely any thing grows except the *Torlok*, or *Pterococcus Aphyllus*, and a few other poor plants on which it feeds. Two were then taken sporting in the mid-day sun; they were both males, and attempted to escape to different holes. The burrows had three entrances running obliquely, and were about a yard deep; lined or plaistered with mud. In the bottom was neither nest nor provision of grass.

The *Asiatic* animal differed in color from the *American*, being above of a light grey mixed with tawny, white below: these colors divided lengthways by a stripe of dusky red. The tail

* *Nov. sp. fasc. i. 314. tab. xviii. B. Mus mindianus? Itin. ii. 702.*

covered with longer and looser hair at the end than in the other parts: the soles of the feet clad with hair. This I could not well observe in the specimen from *Hudson's Bay*, as it was preserved in spirits. *Linnaeus* describes this species under the title of *Mus longipes*, *Syst. nat.* 84. Doctor *Pallas*, with great reason, supposes it to be the same with the *Jird* of Doctor *SHAW*, which our learned countryman describes with the *Jerboa*. It agrees in colors with the above; in its long tail being better clothed than that of a rat; and in its burrowing under ground. This is frequent in *Barbary*, and is reckoned there a good food*.

371. A. CIRCASSIAN.

To this I join, on the judgment of Doctor *Pallas*, another animal, which I described at N° 205 of the *Synopsis* of *Quadrupeds*, under the title of *Circassian Marmot*, or


M. with ears like those of mice: red sparkling eyes: sharp teeth: body long, and of an equal thickness: chestnut-colored hair, long, especially on the back: has sharp claws: tail long and bushy: fore feet shorter than the hind feet: size of the *Hamster*, N° 324.

Inhabits the neighborhood of the river *Terek*, which flows out of *Circassia* and falls into the *Caspian Sea*: runs fast up hill, very slowly down: burrows, and lives under ground †.

* *Shaw's travels*, 248.

† *Scheber's memorab. Asiatic. Russiæ in Muller's Sammlung Russl. viii. 124.*





1. *Middle Jerboa* ——— p. 166.

2. *Tamarisk Rat* ——— N^o 372.

Mus Tamaricinus. Pallas, nov. sp. i. 322. tab. xix. lit. ii. 7c2.

372. TAMARISK.

R. with an oblong head: great whiskers: nose blunt: nostrils covered with a flap: teeth yellow: eyes large and brown: ears large, naked, and oval: neck short: space round the nose and eyes, and beyond the ears, white: sides of the head and neck hoary: back and sides of a yellowish grey: tips of the hairs brown: breast and belly white: tail cinereous; above half annulated with brown: hind legs long: on the fore feet a warty tubercle instead of a thumb.

Length to the tail above six inches: tail not quite so long.

SIZE.

Inhabits the lower salt-marshes about *Saritschikofka*, on the *Lower Taik* or *Ural*, where they burrow beneath the knotty roots of the tamarisk bushes; each burrow has two entrances, and is very deep: they feed only at night: out of numbers which were taken in traps placed before their holes, not a female was taken. Their food is the succulent maritime tribe of plants, such as *Nittraria*, *Salifola*, and others, with which those deserts abound.

PLACE.

To this division of Rats I give the title of *Ferboid*, from the affinity it has to that genus in the length of the hind legs. To the other, *Murine*, as comprehending all the common species of Rats and Mice.

*• Murine.

** Murine.

373. BLACK.

Mus domesticus major, quem vulgò Rat-
tum vocant. *Gesner quad.* 731. *Ratt. syn.*
quad. 217.

Mus Rattus, *Mus Cisternarius*. *Klein quad.*

57.
Ratze. *Kramer Aufsr.* 316.

Mus cauda longissima obscurè cinerea.
Brisson quad. 118.

Mus Rattus. *M. cauda elongata* subnuda,
palmis tetradactylis cum unguiculopoli-
licari, plantis pentadactylis. *Lin. syst.*
Ratta. Faun. jucc. N° 33 *Br. Zool.* i. N°

27.

Le Rat. De Buffon, vii. 278. *tab.* xxxvi.
Pallas nov. sp. fosc. i. 93. *LEV. Mus.*

R. of a deep iron-grey color, nearly black: belly cinereous:
legs dusky, almost naked: a claw, in the place of a fifth
toe, on the fore feet: length, from nose to tail, seven inches; tail
near eight.

PLACE.

Inhabits most parts of *Europe*: of late, the numbers much less-
ened, and in many places extirpated, by the next species: very
destructive to corn, furniture, young poultry, rabbits, and pi-
geons: will gnaw the extremities of infants when asleep: breeds
often in a year: brings six or seven young at a time: makes its
nest, in a hole near a chimney, of wool, bits of cloth, or straw:
will destroy and devour one another: its greatest enemy is the
weasel. First introduced by the *Europeans* into *South America**;
about the year 1544, in the time of the Viceroy *Blasco Nunnez*.
Is now the pest of all that continent.

The word *Rattus* is modern. The *Romans* probably compre-
hended all kinds under the word *Mus*. The *Welsh* call this *Llygoden*
Frengig, or the *French* *Mouffe*, which evinces it not to be a native

* *Garcilasso de la Vega*, 384. *Ovalle. Churchill's coll.* iii. 43.

of our island. There is a very minute variety of this kind about the *Volga*, in the deserts of the lower part (for they have not reached the upper) which scarcely weighs seven drachms.

I cannot trace the original place of the black rat: none are found in *Siberia* or *Kamtschatka*.

Rats (I know not of what species) are found in the *Papuas* islands, off *New Guinea**; but according to the account given by Doctor *Forster*†, the common black rats swarm in *Otaheite*, and other of the *Society* islands, and are also met with in the other groupes of islands, in *New Zealand*, and in *New Holland*. They feed in *Otaheite* on the fruits of the country, and are so bold as even to attack the inhabitants when they are asleep. The natives hold them in the utmost detestation, as unclean animals, and will even avoid killing them, lest they should be polluted by the touch. They will not even eat the bread-fruit these animals should happen to run over.

SOUTH-SEA
ISLANDS.

Le Coypu. *Molina Chili*. 268. *Mus Coypus*. *Gm. Lin.* 125.

374. COYPU.

R. with round ears: nose elongated, covered with whiskers: legs short: tail thick, and of a moderate length, well covered with hairs: two very sharp cutting teeth in each jaw: on the fore feet are five toes, all separated; on the hind feet five, palmated: has the appearance of the otter in hair and size.

This animal lives equally well in the water as on the land; and

MANNERS.

* Captain *Forrest*.

† *Observations*, &c. 185, 187.

VOL. II.

A a

frequents

frequents also houses: is easily tamed, and very content in the domestic state: attaches itself to those who treat it kindly: has a piercing cry on being abused: the female brings forth five or six young, which always follow her.

375. BROWN.

Muscauda longissima, supra dilute fulvus,
infra albicans. Le Rat de Bois. *Briffon*
quad. 120.
Le Surmulot. *De Buffon*, viii. 206. *tab.*

xxvii.
Norway Rat. *Br. Zool.* i. N° 26.
Mus Decumanus. *Pallas nov. sp. fasc.* i.
91. *LEV. Mus.*

R. with the head, back, and sides, of a light brown color, mixed with tawny and ash-color: breast and belly dirty white: feet naked, and of a dirty flesh-color: fore feet furnished with four toes, and a claw instead of the fifth: length, from nose to tail, nine inches; tail the same: weight eleven ounces: is stronger made than the last.

IN EUROPE.

Inhabits most parts of *Europe*: but was a stranger to that continent 'till the present century: came into *Great Britain* about fifty years ago: not known in the neighborhood of *Paris* half that time. This rat is common in *India*, both on the land and in ships. May we not go to *Indostan* for their origin? They swarm in *Petersburg*: have reached *Prussia*, but not the opposite side of the *Baltic*; at least *Linnaeus* takes no notice of them.

ASIA.

Are numerous in *Persia*, where they burrow in the fields*. In *Hyrkania* they occupy the deserted holes of the porcupine.

* Doctor *Pallas*, among his other epistolary communications,

Some

Some years ago an immense migration arrived from the west at the town of *Taïk*; and in the year 1727 an equal number appeared about *Astracan*, filled the whole bed of the *Volga*, and infested the houses to that degree, that nothing could be preserved from them. They have not yet reached *Siberia*. These probably were the *Mures Caspii* of *Ælian*, which he says were little less than *Ichneumons*; and made periodical visits in infinite multitudes to the countries bordering on the *Caspian Sea*: swimming boldly over the rivers, holding by one another's tail *.

Burrow, like the water-rat, on the sides of ponds and ditches: swim well and dive readily: live on grain and fruits, and will destroy poultry and game: encrease fast; bring from fourteen to eighteen young at a time: are very bold and fierce; will turn when closely pursued, and fasten on the stick or hand of those who offer to strike them: have destroyed the common black rat in most places. Inhabit fields part of the year, but migrate in great numbers into houses, and do infinite mischief.

Le Rat Perchal, *De Buffon, Supplem.* vii. 276. tab. lxi.

376. PERCHAL*

R. with ears rounded on the top: nose long and turning up: body longer than that of common rats: hair on the upper parts deep brown: hind legs larger than the fore: tail naked and scaly: length from nose to tail above a foot; tail between eight and nine inches.

Common in *India*, and infests the houses in *Pondicherry*,

PLACE.

* *Æliani hist. ar.* xvii. c. 17.

as our rats do those of *Europe*: are very numerous: the inhabitants use them for food.

377. BANDICOTE.

At p. 440 of the former edition I imagined that the *Brown rat* was the same as the *Bandicote* of the *East Indies*. My good and intelligent friend Doctor *Patrick Ruffel*, who has made a long residence on the eastern coast of *Indo-Java*, convinces me of my mistake. His remarks are such that do not at present enable me to give so full a description of this species as I could wish. It is generally agreed that the *Bandicote* is at least five times the weight of the *Brown rat*: that, comparative with that kind, it has a shorter and thicker tail: that its general form is much thicker, and the back arched; so that, at first sight, it looks like a little pig: it is less active and alert than the brown-rat: is infinitely mischievous in gardens: burrows under the houses, and will even undermine them so as to cause them to fall: never go on board ships. The *Palinquin-boys* eat this kind, but will reject the common rat. A more satisfactory account of the *Bandicote* may be expected in the course of a year.

378. AMERICAN.

LEVERIAN MUSEUM.

Mus Caraco? *Pallas nov. sp. fasc. i. 335. tab. xxiii.*

R. with the upper jaw much longer than the lower: head long: nose narrow and pointed: ears large and naked: whiskers fine, but long: tail naked, and like that of the black rat, but not so long.

Color

Color a deep brown; on the belly inclines to ash-color: hair ruder than in the preceding species.

In size larger than the black, lesser than the brown rat.

Inhabits *North America*; but I am uncertain whether it is entirely wild, or whether it has yet found its way into houses and out-houses. Mr. *Bartram** mentions the rat (but does not determine the species) which lives among the stones and caverns in the *Blue Mountains*, far from mankind: comes out at night, and makes a terrible noise; but in very severe weather keeps silent within its holes.

The *Mus CARACO* of Doctor *Pallas* is so nearly allied to this species, that I do not at this time venture to separate them: the whiskers of the former seem rather shorter, and the tail, in proportion to its length, thicker; but the thinness of that part might, in the specimen in the *LEVERIAN MUSEUM*, arise from its being dried; neither could I examine it thoroughly, as it was within a glass case. The *Caraco* has not as yet appeared to the west of the *Jenesai*, but swarms about and beyond lake *Baikal*. It has much agreement with the last kind, being, as the *Mongals* report, conversant among lakes and waters, and is called by them *Chavacko*, and *Jike-Cholgonach* or the *Great Mouse*. It burrows in the banks of rivers: is supposed to extend to *China*, and to be very noxious there.

β. CARACO.

* In *Kalm's trav.* ii. 48.

379. SCHERMAN.

Le Scherman. *De Buffon, Supplem.* vii. 278. tab. lxx.

R. with a short head and thick nose: small eyes: ears so very small as to be scarcely visible: color of the hair dusky, mixed with grey and tawny: edges of the mouth bordered with white: body six inches long; tail above two.

Common about *Strasbourg*, in the gardens and places near the water: make great havoc among the plants and the cultivated grounds: swim and dive very well, and are often taken by the fishermen in their weels: burrow under ground, and are frequently caught in the traps used by the people who are employed in taking the *Hamster* rat.

380. WATER.

Le Rat d'Eau. *Belon, Aquat.* 30. tab. xxxi.
Mus aquatilis. Agricola An. Subter. 488.
Gesner quad. 732. *Raii Syn. quad.* 217.
Klein quad. 57.
 Wasser-maus *Kramer Austr.* 316.
Mus Amphibius. M. cauda elongata pilosa, plantis palmatis. Lin. syst. 82. *Faun. succ.* N° 32. *Pallas Nov. sp. fasc.* i. 20.
M. cauda longa pilis supra ex nigro et flavescente mixtis, infra cinereis vestitus. Brisson quad. 124.
 Le Rat d'Eau. *De Buffon*, vii. 348. tab. xliii.
 Water Rat. *Br. Zool.* i. N° 27. *Lev. Mus.*

R. with a thick blunt nose: ears hid in the fur: eyes small: teeth yellow: on each foot five toes; inner toe of the fore foot very small; the first joint very flexible: head and body covered with long hairs, black mixed with a few ferruginous hairs: belly of an iron grey: tail covered with short black hairs; the tip whitish: weight nine ounces: length, from nose to tail, seven inches;

inches; tail only five: shape of the head and body more compact than the former species*.

Inhabits *Europe*, the north of *Asia*, and *Norib America*†; burrows in the banks of rivers, ponds, and wet ditches: feeds on small fish, and the fry of greater; on frogs, insects, and roots: is itself the prey of pike: swims and dives admirably, though it is not web-footed, as Mr. Ray supposed, and *Linnaeus* copied after him: brings six young at a time. This animal and the Otter are eaten in *France* on *maigre* days.

Le Guanque. *Molina*, 281. Mus Cyanus. *Gmelin*, 132.

381. SKY-COLORED.

R. with rounded ears: fur of a blue color: size and appearance of my field rat.

Inhabits *Chili*: burrows a gallery ten feet long, with seven correspondent chambers on each side of a foot in depth: these are the magazines for winter provision, which are of roots, most nicely laid in order one upon the other: at the approach of the rainy season retire to the burrows: breed twice in the year, and bring forth six

* It has some resemblance to the Beaver, which induced *Linnaeus*, in the first edition of his *Fauna Suecica*, to style it *Casor cauda lineari teriti*.

† *Larsson* *bijl. Carolina*, 122. He also mentions another, which he calls the *Marsk Rat*, being more hairy than the common rat; but apparently is the same with this. Those of *Canada* vary to tawny and white. Vide *De Buffon*, xiv. 401. xv. 146.

at a time: the first brood is left to provide for itself; the second retires under ground with the parents: are very timid, and very cleanly in their retreats: the peasants hunt for the hoards, and by robbing them leave the family to perish.

382. MOUSE.

- Mus domesticus communis seu minor.* *Lin. syst.* 83. *Pallas Nov. sp. fasc.* i. 95.
Gefner quad. 714. *Raii syn. quad.* 218. *Mus. Faun. suec.* N° 34.
Mus minor, musculus vulgaris. Kleinquad. *Mus cauda longissima, obscure cinereus,*
Maus. Kramer Austr. 316. *ventre subalbescente. Brisson quad.* 119.
Mus musculus. M. cauda elongata, palmis *La Souris. De Buffon, vii. 309. tab. lix.*
tetradactylis, plantis pentadactylis. *Br. Zool. i. N° 30. LEV. MUS.*

AN animal that needs no description: when found white, is very beautiful, the full bright eye appearing to great advantage amidst the snowy fur.

Inhabits all parts of the world, except the *Arctic*: follows mankind.

383. FIELD.

- Mus agrestis minor. Gefner quad.* 733. *Mus sylvaticus. M. cauda longa, palmis*
Mus domesticus medius. Raii syn. quad. *tetradactylis, plantis pentadactylis,*
218. corpore griseo pilis nigris abdomine al-
Maus mit weissen bauch. Kramer Austr. *bo. Lix. syst.* 84. *Pallas Nov. sp. fasc.* i.
317. 94. Faun. Suec. N° 36.
Mus cauda longa supra e fusco flavescens, infra ex albo cinerefcens. Brisson quad. *Le Mulot. De Buffon, vii. 325. tab. xli.*
123. Long-tailed Field-mouse. Br. Zool. i.
N° 28. LEV. MUS.

R. with full and black eyes: head, back, and sides of a yellowish brown, mixed with some dusky hairs: breast of an ochre-color: belly white: length, from the tip of the nose to the tail,

four inches and a half: tail four inches, slightly covered with hair.

Inhabits *Europe*: found only in fields and gardens: feeds on nuts, acorns, and corn: forms great magazines of winter provision: hogs, tempted by the smell, do much damage in the fields by rooting up the hoards: makes a nest for its young very near the surface, and often in a thick tuft of grass: brings from seven to ten at a time: called; in some parts of *England*, *Bean Mouse*, from the havock it makes among the beans when just sown.

Is common in *Russia*, and about the *Uralian* chain, but not beyond.

«. **AMERICAN R.** with very long whiskers, some white, others black: ears large, naked, and open: from the head to the tail, along the middle of the back, a broad dark stripe, ferruginous and dusky: the cheeks, space beneath the ears, and sides, quite to the tail, orange-colored: under side, from nose to tail, of a snowy whiteness: feet white: hind legs longer than those of the *European* kind: tail dusky above, whitish beneath. *New York*.

The less long-tailed Field-Mouse. *Br. Zool. ii. App. 498. LEV. MUS.*

384. HARVEST.

R. with eyes less prominent than those of the former: ears prominent: of a full ferruginous color above, white beneath: a strait line along the sides divides the colors: tail a little hairy: length, from nose to tail, two inches and a half: tail two inches: weight one-sixth of an ounce.

VOL. II.

B b

Inhabits

Inhabits *Hampshire*; where it appears in greatest numbers during harvest: never enters houses; but is carried into the ricks of corn in the sheaves; and often hundreds are killed on breaking up the ricks: during winter, shelters itself under ground: burrows very deep, and forms a warm bed of dead grass: makes its nest for its young above ground, between the straws of standing corn; it is of a round shape, and composed of blades of corn: brings about eight young at a time.

385. ORIENTAL.

Mus orientalis. Seb. Mus. ii. 22. tab. xxi.

fig. 2.

M. cauda mediocri subnuda, palmis tetradactylis, plantis pentadactylis, corporis striis punctatis. Lin. syst. 84.

M. cauda longa, striis corporis longitu-

dinalibus & punctis albis. Mus. Ad. Fred. 10.

Mus cauda longa, rufus, lineis in dorso albicantibus, margaritarum æmulis. Brisson quad. 124.

R. with round naked ears: of a grey color: the back and sides elegantly marked with twelve rows of small pearl-colored spots, extending from the head to the rump: tail the length of the body: in size, half that of a common mouse.

Inhabits *India*. In the same country, and in *Guinea*, is another very small species, which smells of musk. The *Portuguese* living in *India* call it *Cheroso*, and say its bite is venomous. *Boullaye la Gouz. 256. Barbot's Guinea, 214.*

Mus Barbarus. *M. caudamediocri corpore fusco, striis decem pallidis, palmis tri-*

daetylis, plantis pentadaetylis. Lin. syst. tom. i. pars ii. addenda.

386. BARBARY.

LESS than the common mouse: of a brown color: marked on the back with ten slender streaks: three toes with claws on the fore feet, with the rudiments of a thumb: tail of the length of the body.

Inhabits *Barbary*.

Mus Mexicanus maculatus. Seb. Mus. 74. tab. xlv. fig. 5.

387. MEXICAN.

R. of a whitish color, mixed with red: head whitish: each side of the belly marked with a great reddish spot.

According to *Seba* inhabits *Mexico*.

Mus agrestis Americanus albus. Seb. Mus. i. 76. tab. xlvii. fig. 4.

388. VIRGINIAN.

R. with pointed ears and nose; the last black: whiskers long: fur very short: limbs very weak and slender: tail at the base thick, growing gradually so from the rump, so that the junction cannot be distinguished; decreases gradually, and becomes very long and slender; ends in a point, and is in all parts beset with long hair.

Color of this animal universally white.

According to *Seba*, found in *Virginia*. The thickness at the base of the tail is its specific difference.

PLACE.

B b 2

Mus

389. WANDER-
ING.

Mus Vagus. Pallas Nov. sp. fasc. i. 327. tab. xxii. fig. 2.

R. with an oblong head: blunt nose, with a red tip: cutting teeth yellow; the upper truncated: eyes placed midway between the nose and the ears: ears large, oval, naked; the tip dusky and downy: limbs slender: instead of a thumb, on the fore feet, is a conic wart: tail longer than the body, and very slender.

Color above a pale ash, mixed and undulated with black: along the back to the tail is a black line: ends of the limbs whitish.

Length, from nose to tail, between two and three inches; the tail near three.

Inhabits the whole *Tartarian* desert; and at certain times wanders about in great flocks, and migrating from place to place during night. Observed as high as lat. 57, about the *Irtish*, and between the *Oby* and *Jenesai*, in birch woods: is of a very chilly nature; soon becomes torpid, and sleeps rolled up in the cold night, even of the month of *June*. Lives in fissures of rocks, under stones, and in hollow fallen trees: has carnivorous inclinations; for on being placed in a box with a mouse of another species, it killed and devoured part, notwithstanding it had seeds to feed on. Is called by the *Tartars*, *Djickis-fitkan*, or *gregarius Mousé*.

Mus

Mus Betulinus. Pallas Nov. sp. fasc. i. 322. tab. xxii. fig. 1.

390. BEECH.

R. with a sharp nose, with the end red: ears smaller than those of the former, brown, bristly at the end: limbs very slender: toes long, slender, and very separable: tail very long and slender, much exceeding the length of the body; brown above, white below.

Color of the head and body a cinereous rust, with a few dusky hairs interspersed: breast and belly, pale ash: along the top of the back is a dusky line.

Less than the former.

Inhabits the birch woods about the plains of *Ischim* and *Baraba*, and between the *Oby* and *Jenessei*: lives in the hollows of decayed trees: very tender, and soon grows torpid in cold weather: runs up trees, and fastens to the boughs with its tail; and, by assistance of its slender fingers, adheres to any smooth surface: emits a weak note.

SIZE.
PLACE.

Mus Agrarius. Pallas nov. sp. fasc. i. 341. tab. xxiv. A. Itin. i. 454.
Mus Rubeus. Schwenkfeldt Anim. Silf. 114.

391. RUSTIC.

R. with a sharp nose: oblong head: small ears lined with fur: color of the body and head ferruginous, with a dusky line along the back: belly and limbs whitish: above each hind foot is a dusky circle.

A little less than the field mouse. The tail only half the length of the body.

4.

Inhabits.

PLACE.

Inhabits the temperate tract of *Russia*, and *Siberia*, as far as the *Irtish*: in the former, chiefly about villages and corn-fields; in the latter, in woods. In *Russia* is often migratory, and often very noxious to the grain: it is called there *Sbitnik*, or the *Corn Mouſe*, for it abounds in the sheafs and ricks. At times they wander in vast multitudes, and destroy the whole expectations of the farmer. This plague did in particular, in the years 1763 or 1764, make great ravages in the rich country about *Cafan* and *Aiſk*; but came in ſuch numbers as to fill the very houſes, and became through hunger ſo bold as to ſteal even the bread from the table before the very faces of the gueſts. At approach of winter they all diſappeared.

They make their retreats a little below the ſurface, which in thoſe places appears elevated: each has a long gallery, with a chamber at the end, in which they place their winter food, which conſiſts of various ſorts of feeds.

392. SORICINE.

Mus Soricinus. Schreber, tab. clxxxiii. Gm. Lin. 130.

R. with an elegant ſlender head: ears rounded and covered with hair: tail long and ſlender: hair on the head and upper part of the body cinereous, mixed with yellow: belly white: length two inches.

PLACE.

Inhabits the neighborhood of *Strasburg*: diſcovered by Profeſſor *Herman*.

Mus

Mus pumilio. *Gmel. Lir. Syst.* 130. *Sparman's voy.* ii. 349. *tab.* vii.

303. LINEATED.

R. with black forehead and hind part of the head: from the last extend along the back to tail four black lines: color of the rest of the animal a cinereous brown: tail of a light color, very small: not superior in size to the following.

Inhabits the forest of *Siticania* on the *Slangen* river, at a vast distance to the eastward of the *Cape of Good Hope*.

PLACE.

Mus minutus. *Pallas Nov. sp. fasc.* i. 345. *tab.* xxiv. *B. Ilin.* i. 454.

394. LITTLE.

R. with a sharpish nose: dusky, with a whiteness at the corner of the mouth: ears small, half hid in the fur: body more slender than that of the common mouse: tail shorter and more slender.

Color, a deep tawny above, white below: feet grey.

The rest of the genus; little more than two inches long from nose to tail; weight not half a dram.

SIZE.

Inhabits the temperate parts of *Russia* and *Siberia*, in corn-fields and barns; is also plentiful in the birch-woods. More males among them than females. Seem to wander without any certain places for their nests.

PLACE.

** With

325. ROCK.

** With tails of middle length.

Mus Saxatilis. Pallas Nov. sp. fasc. i. 255. tab. xxiii. B.

R. with an oblong head; nose rather pointed: ears rising above the fur; oval, downy, at the edges brown: whiskers short: limbs strong: tail not half so long as the body, with a few hairs scattered over it.

Color above, brown slightly mixed with grey: sides incline more to the last color: belly of a light cinereous: snout dusky, surrounded with a very slender ring of white.

SIZE.

Length four inches: tail one and a half.

PLACE.

Inhabits the country beyond lake *Baikal*, and the *Mongolian* desert: makes its burrows in a wonderful manner, considering the weakness of its feet, between the crannies of the rocks which had been forced open by violence of frost, or the insinuation of roots of plants: it chuses its habitation amidst the rudest rocks, and lives chiefly on the seeds of *Astragali*. The burrows consist, firstly, of a large winding oblique passage, through which the animal flings out the earth: secondly, of one or more holes pointing downwards, which likewise wind among the rocks; and at their bottom is the nest, formed of soft herbs.

Viverra

Viverra fasciata. Gmelin Lin. i. 92. Chat sauvage, &c. Sonnerat voy. ii. 143. tab. lxxix.

396. INDIAN.

R. with short pointed ears: sharp nose: two cutting teeth in each jaw, and fourteen grinders in each: five toes to each foot: claws strong and crooked: color grey, tinged on the lower part of the head and neck with red: belly white: back and sides marked with four black lines, commencing near the hind part of the head, and ending at the rump: on each thigh is a bifurcated black stroke, the forks pointing backwards.

Length two feet; tail nine inches. Inhabits *India*. No further account is given by M. *Sonnerat* of this and the following obscure species. I place them in this genus, as they have no canine teeth, and only two incisores in each jaw.

Le Zenik des Hottentots. Sonnerat voy. Viverra Zenik. Gmelin Lin. i. 84; ii. 145. tab. xcii.

397. ZENIK.

R. with short ears: very long sharp nose: two cutting teeth; sixteen grinding teeth: four toes on each foot: claws on the fore feet very long, and almost strait: color of a reddish grey, striped transversely with ten black lines falling from the back almost to the belly.

Size of a water rat: tail not so long as the body: of a gilded red on three parts of its length; the rest black.

Inhabits the land of the *Hottentots*.

VOL. I.

C c

Mus

398. **ÆCONOMIC.** *Mus Œconomus. Pallas Nov. sp. f. sc. i. Tegoulitchek. Deser. Kamtschatka, Engl. 234. tab. xiv. A. Itin. iii. 692. ed. 104.*

R. with small eyes: ears naked, and usually hid in the fur: limbs strong: teeth very tawny: color black and yellow, intimately mixed; dusky on the back; from throat to tail hoary; beneath the hair a dark down; ends of the feet dusky.

SIZE. Length four inches and a quarter; of the tail, more than an inch: in form of body like the meadow mouse, but is rather longer, and the belly bigger. The females are far superior to the males in size, as on the former rests the chief labor of providing the food.

PLACE. Inhabits in vast numbers all *Siberia*, especially the eastern parts, and *Kamtschatka*; and even found within the *Arctic* circle.

MANNERS. They are called by Doctor *Pallas*, *Mures Œconomi* or *Œconomic Mice*, from their curious way of living. They inhabit damp soils, and shun the sandy; form burrows beneath the upper crust of the turfy ground; and have in them many chambers, and several entrances. Never more than two animals are found in these extensive nests, and those fondly attached to each other; sometimes only one inhabits these dwellings, except towards autumn, when the whole family make it their residence. In the first they form magazines for winter food, consisting of various sorts of plants, which they collect in summer with great pains; and in sunny days draw them out of their nests, in order to give them a more effectual drying. During summer they never touch their hoards, but live on berries, and other vegetable productions.



1. Long-eared Rat — . V^o 303. 2. Vietnamese Rat — . V^o 398.
3. Sulphur — . V^o 422

Twenty, and even thirty pounds of fresh roots, have been found in one hoard. Besides man, these mice find a cruel animal in the wild boars, which ransack the magazines, and devour the little defenceless owners.

They in certain years make great migrations out of *Kamtchatka*; they collect in the spring, and go off in incredible multitudes. Like the *Lemmus*, they go on in a direct course, and nothing stops their progress, neither rivers nor arms of sea: in their passage they often fall a prey to the ravenous fishes and birds; but on land are safe, as the *Kamtchatkans* pay a superstitious regard for them; and when they find them lying, weak or half dead with fatigue, on the banks, after passing a river, will give them all possible assistance. They set out on their migration westward. From the river *Pengin* they go southward, and about the middle of *July* reach *Ochotska* and *Judoma*, a tract of amazing extent. They return again in *October*. The *Kamtchatkans* are greatly alarmed at their migrations, as they presage rainy seasons, and an unsuccessful chase; but on their return, expresses are sent to all parts with the good news.

Many fables are related of them, such as that they cover their provisions with poisonous herbs before their migrations, in order to destroy other rats which may attempt to plunder their magazines; and if by chance they should be pillaged, they will strangle themselves through vexation, by squeezing their necks between the forks of shrubs; for this reason the natives never take away all their store, but leave part for their subsistence, or leave in its place some caviare, or any thing that will serve for their support. It is certain that the roots of certain poisonous plants are

often found in their nests half eaten: but this is no wonder, as it is well known that divers animals will feed on noxious vegetables which would prove the certain bane of others.

399. WOOLLY.

La Chinchilla, *Molina Chili*. 283. *Mus laniger*. *Gm. Lin.* 134.

R. with very small ears: short nose: tail of a middling length: whole body covered with long wool of exquisite fineness, grey, and long enough to be spun. The length of this species is six inches.

These animals live in society under ground, and feed on the bulbous roots of the country. It breeds twice a year, and brings five or six at a time: it is a very gentle tame animal: very fond of being caressed, and will lie down without fear by mankind: it is often domesticated. The ancient *Peruvians* manufactured many small articles from the wool, which they sold at a great price.

400. RED.

Mus Rutilus. *Pallas Nov. sp. fasc.* i. 146. *tab.* xiv. B.

R. with the nose and face very bristly: ears, like those of the former, naked, except the tip, on which is a rusty down: tail full of hair: color, from the middle of the forehead, along the back, to the rump, an uniform pleasant tawny red: the sides light grey and yellow: under side of the body whitish: feet white: tail dusky above, light below.

SIZE.

Length not four inches; tail above one.

Inhabits

Inhabits *Siberia*, from the *Oby* eastward to *Kamtschatka*, in woods and mountains; and also within the *Arctic* circle. Creeps sometimes into houses and granaries; lives abroad under logs of wood, or trunks of trees: they wander out the whole winter, and are very lively even amidst the snows: eat any thing which comes in their way; even flesh.

A variety is found about *Casan*, a little lesser than the *Siberian* kind, and the tail longer and more slender: the red on the back is not so much diffused as in the other. The same kind has also been discovered in the botanical garden at *Gottengen*.

PLACE.

Mus Alliarius. Pallas nov. sp. fasc. i. 252. tab. xiv. C.

401. GARLIC.

R. with great open naked ears, very apparently out of the fur: tail clothed with hair: color on the back cinereous, mixed with longer hairs tipped with dusky grey: sides of a whitish ash: breast, belly, and feet white: tail marked along the top with a dusky line, the rest white.

Length a little above four inches; tail one and a half.

Inhabits the country about the *Jenesei* and *Lena*: is frequent in the subterraneous magazines of bulbous roots, especially the *Allium angulatum*, or angular garlic, formed by the *Siberian* peasants.

SIZE.

PLACE.

R. with the nose a little extended; four toes on the fore feet, with a tubercle instead of a thumb: five toes on the hind feet: round ears covered with fur: tail of a middling length, and hairy:

402. SORICINE.

hairy: color of the upper part of the body yellowish grey: belly white.

PLACE. Inhabits the neighborhood of *Straßbourg*. Discovered by Professor *Herman*.

*** With short tails.

403. LEMMUS. Lemmar vel Lemmus. *Olaus magnus de gent. Septentr.* 358.
 Leem vel Lemmer. *Gesner quad.* 731.
 Mus Norvegicus vulgò Leming. *Worm Mus.* 321, 325. *Scheffer Lapland*, 136.
Pontop. Norw. 5, ii. 30. *Sirom. Sundmor.* 154. *Raii syn. quad.* 227.
 Sable-mice. *Ph. Tr. abridg.* ii. 875.
 Cuniculus caudatus, auritus, ex flavo, rufo et nigro variegatus. *Briffon quad.* 100.
 Mus Lemmus. M. cauda abbreviata, pedibus pentadactylis, corpore fulvo nigro vario. *Lin. syst.* 80. *Pallas nov. sp.* f. sc. i. 186. *tab.* xii. A. & B.
 Fial-Mus, Sabell-Mus, Lappis.
 Lummick. *Faun. Suec.* N° 29.
 Le Leming. *De Buffon*, xiii. 314.

R. with two very long cutting teeth in each jaw: head pointed: long whiskers; six of the hairs on each side longer and stronger than the rest: eyes small and black: mouth small: upper lip divided: ears small, blunt, and reclining backwards: fore legs very short: four slender toes on the fore feet, covered with hairs; and in the place of the thumb a sharp claw, like a cock's spur: five toes behind: the skin very thin: the color of the head and body black and tawny, disposed in irregular blotches: belly white, tinged with yellow.

SIZE. Length, from nose to tail, about five inches: in large specimens a little more: the tail about half an inch. Those of *Russian Lapland* are much less than those of the *Norwegian* or *Swedish*.

PLACE. Inhabits *Norway* and *Lapland*, the country about the river *Oby*,

2



1



M. Agate sculp.

1. *Lemmus* ——— . 1. 403.

2. *Vandy* of the same.



Oby, and the north extremity of the *Uralian* chain. They appear in numberless troops, at very uncertain periods, in *Norway* and *Lapland*: are the pest and wonder of the country: they march like the army of locusts, so emphatically described by the prophet *Joel*: destroy every root of grass before them, and spread universal desolation: they infect the very ground, and cattle are said to perish which taste of the grass which they have touched: they march by myriads, in regular lines: nothing stops their progress, neither fire, torrents, lake, or morass. They bend their course strait forward, with most amazing obstinacy; they swim over the lakes; the greatest rock gives them but a slight check, they go round it, and then resume their march directly on, without the least deviation: if they meet a peasant, they persist in their course, and jump as high as his knees in defence of their progress: are so fierce as to lay hold of a stick, and suffer themselves to be swung about before they quit their hold: if struck, they turn about and bite, and will make a noise like a dog.

They feed on grass, on the rein-deer liverwort, and the catkins of the dwarf birch. The first they get under the snow, beneath which they wander during winter; and make their lodgements, and have a spiracle to the surface for the sake of air. In these retreats they are eagerly pursued by the *Arctic* foxes.

They make very shallow burrows under the turf; but do not form any magazines for winter provision: by this improvidence it seems that they are compelled to make these numerous migrations, in certain years, urged by hunger to quit their usual residences.

They breed often in the year, and bring five or six young at a time:

time: sometimes they bring forth on their migration; some they carry in their mouths, and others on their backs.

They are not poisonous, as is vulgarly reported; for they are often eaten by the *Lapländers*, who compare their flesh to that of squirrels.

Are the prey of foxes, lynxes, and ermines, who follow them in great numbers: at length they perish, either through want of food, or by destroying one another, or in some great water, or in the sea. They are the dread of the country: in former times spiritual weapons were exerted against them; the priest exorcised, and had a long form of prayer to avert the evil*: happily it does not occur frequently; once or twice in twenty years: it seems like a vast colony of emigrants, from a nation over-stocked; a discharge of animals from the great *Northern* hive, that once poured out its myriads of human creatures upon Southern *Europe*. Where the head-quarters of these quadrupeds are, is not very certainly known; *Linnaeus* says, the *Norwegian* and *Lapland Alps*; *Pontoppidan* seems to think, that *Kolens* rock, which divides *Nordland* from *Sweden*, is their native place: but wherever they come from, none return: their course is predestinated, and they pursue their fate.

* *Worm. Mus.* 333. where the whole form is preserved. It was once seriously believed that these animals were generated in the clouds, and fell in showers upon the ground: *Per tempestates et repentinos imbres e celo decidunt, incompertum unde, an ex æmulatoribus insulis, et huc vento delatae, an ex nubibus sæculentis natae deferantur.* *Olaus Magnus* de Gent. Septentr. 358.

Mus torquatus. Pallas Nov. sp. fasc. i. 205:

404. RINGED.

R. with a blunt nose: ears hid in the fur: legs strong and short: soles covered with hair: claws very strong, hooked at the end: the hair on the whole body very fine.

Color of the upper part of the body ferruginous, mixed with grey and yellow; sometimes pale grey, clouded with undulated lines of dusky rust-color: from the ears, down each side of the cheeks, is a bed of the same color, and behind that a stripe of white, so that the neck appears encircled with a collar; behind these again is another bed of the former color.

Length to the tail little more than three inches; of the tail one; at its end is a hard tuft of bristles.

SIZE.

Inhabits the northern parts about the river *Oby*. Makes its burrows, with many passages, beneath the turfy soil. The nests are filled with rein-deer and snowy liverworts. They are said to migrate at the same seasons with the *Lemmus*.

PLACE.

Mus Hudsonius. Pallas Nov. sp. fasc. i. 203.

405. HUDSON'S.

R. with slender brown whiskers: very fine long soft hair: cinereous, tinged with tawny, on the back, with a dusky stripe running along its middle: along each side a pale tawny line: belly pale cinereous: limbs very short: fore feet very strong: the two middle claws of the male very strong, thick, and compressed; divided at the end: those of the supposed females (of the

VOL. II.

D d

lesser

leffer skins) small: tail very short, terminated by some stiff bristles.

SIZE. Length about five inches. Described by Doctor *Pallas*, from some skins sent to him from *Labrador*, one of which he favored me with.

406. HARE-
TAILED.

Mus Lagurus. Pallas Nov. sp. fasc. i. 210. tab. xiii. A. lin. ii. App. 704.

R. with a long head, and blunt nose: rough lips, and swelling out: ears short, round, flat, just appearing out of the fur: limbs short and slender: tail the shortest of all the genus, scarcely appearing out of the hairs: fur very soft and full, cinereous on the upper part, mixed with dusky: along the back is a dark line: belly and feet of a pale ash-color.

Length between three and four inches.

SIZE,
PLACE,
MANNERS.

Inhabits the country above the *Taik*, *Irish*, and *Jenesei*. They love dry soils, but firm; in which they make burrows with two entrances; one oblique, leading to the nest, the other perpendicular, but both end at it, or unite beyond; the nest is formed of grass. Usually the male has a different habitation, but sometimes they live together. When more males than one get together, they fight, and the conqueror devours the vanquished; the mate of the deceased instantly submits to the embraces of the former, even though pregnant. They are very falacious, and bring their young frequently in the air: they bring six at a time: emit often a musky smell when in heat: the males fight sitting up, and bite very hard, and make a noise by striking their teeth together. They sleep very much, and like the Marmots, rolled up;

up; and, like them, are slow in their motions: are very fond of the dwarf iris, but feed on all sorts of seeds: they have also carnivorous appetites, for they will devour one another, and even others of different species, of the same size with themselves; for which reason few other kinds live near them. They migrate in great troops; therefore are called by the *Tartars*, *Dybilkis-Zizchan*, the *Rambling Mouse*.

Mus socialis. *Pallar Nov. sp. fasc. i. 218. tab. xiii. B. Itin. ii. App. 703.*

407. SOCIAL.

R. with a thick head and blunt nose: whiskers white: ears oval, naked: limbs short and strong: tail slender: nose dusky: upper part of the body a light grey; palest on the sides: sides, shoulders, and belly, white.

Length above three inches; tail an inch.

Inhabits the *Caspian* desert, between the *Volga* and the *Tarik*, and the country of *Hircania*. They live in sandy, low, and herby places, in large societies; and in many places the whole ground is covered with the little hills formed by the earth they cast out of their burrows: the burrows are about a span in depth, with eight or more passages. They are always found to live in pairs, or with a family. They live much on tulip-roots. They rarely appear in autumn, but swarm in the spring. They are said either to migrate or change their places in autumn, or to conceal themselves among the bushes; and in the winter to shelter in hay-ricks. They breed later than other kinds. Are the prey of weasels, sitchets, crows, and vipers.

SIZE.

PLACE.

MANNERS.

408. BAIKAL.

Mus Gregalis. Pallas nov. sp. 238. Gmel. Lin. syst. 133.

R. with large thin ears appearing above the fur; whiskers black; hair rough and hard; color above a pale grey: the back darkened with dusky hairs, which gradually decline into the lighter color: body below of a dirty white: the legs stronger, the tail thicker, than in the SOCIAL species: about the size of that kind.

Inhabits *Siberia*, but not like the country beyond the *Oby*: most plentiful about the *Baikal* lake and *Trans-Baikal* region; especially those places which abound most with the *Lilium pomponium* and *allium tenuissimum*; and *Siberia* and *Hircania*. They collect the roots of these and of the *Trifolium Lupinistrum*, for winter food. They form their lodge beneath the turf, and have many minute entrances: the earth that they sling out is carefully heaped above their lodge, in form of a hillock, to divert the rain. In this retreat the male, female, and the progeny of one year, reside. This species is never observed to migrate.

Mus agrestis capite grandi brachiurus.
Raii syn. quad. 218.

Mus terrestris. *M. cauda mediocri sub-*
pilosa, palmis subtetradaetylis, plantis
pentadaetylis, auriculis vellere brevio-
ribus. *Lin. syst.* 82.

Molle. *Faun. suec.* N° 31*.

Mus cauda brevi, pilis e nigricante &
fordidè luteo mixtis in dorso, & satu-

ratè cinereis in ventre vestitis. *Briffon*
quad. 125.

Le Campagnol. *De Buffon*, vii. 369. *tab.*
xlvii.

The short-tailed Field-mouse. *Br. Zool.* i.
N° 31.

Erdzeiß. *Kramer Austr.* 316.

Mus arvalis. *Pallas Nov. sp. fasc.* i. 78.
LEV. MUS.

409. MEADOW.

R. with a large head: blunt nose: ears short, and hid in the
fur: eyes prominent: tail short: color of the head and
upper part of the body ferruginous, mixed with black: belly
deep ash-color: feet dusky.

Length, from nose to tail, six inches; tail only one and a half,
thinly covered with hair, terminated by a small tuft.

SIZE.

Inhabits *Europe*, *Siberia* and *Hircania*; also in great abundance in
Nezefoundland, where it does much mischief in the gardens: in *Eng-*
land, seldom infests gardens: makes its nest in moist meadows:
brings eight young at a time: has a strong affection for them: re-
sides under ground: lives on nuts, acorns, and corn.

PLACE.

* The species, N° 30. *Faun. suec.* described by the style of *Mus cauda abbreviata*,
corpore nigro fusco, abdomine cinerescente, seems the same with this.

410. GREGARIOUS.

Mus gregarius. M. cauda corpore triplo brevior subpilosa, corpore griseo sub-

tus pedibusque albis. *Lin. f. f. 84.*

R. with a small mouth and blunt nose: ears naked, and appearing above the fur: hair on the upper part of the body black at the roots and tips, ferruginous in the middle: throat, belly, and feet whitish: tail thrice as short as the body, covered with thin white hairs; the end black and ash-color: is a little larger than the common mouse.

Inhabits *Germany* and *Sweden*: eats sitting up: burrows, and lives under ground.

*** Short-tailed.

With pouches in each jaw.

411. HAMSTER.

Hamster, *Cricetus Agricola An. Subtr.*

486. *Gesner quad.* 738. *Raii Syn. quad.*

221. *Mayer An. i. tab.* lxxxi. lxxxii.

Skizetzel, Chomik, *Rzaczinski Po on.*

232.

Porcellus frumentarius. *Sclavenkfelds The-*

riol. 1b. 118.

Krichtsch, Hamster. *Kramer Austr.* 317.

Pallas Nov. sp. f. f. i. 83. *Zimmerman.*

341. 511.

Mus cricetus. M. cauda mediocri, auri-

culis rotundatis, corpore subtus nigro, lateribus rufescentibus maculis tribus albis. *Lin. f. f. 82.*


Glis ex cinereo rufas in dorso, in ventre niger, maculis tribus ad latera albis. *Briffon quad.* 117.

Le Hamster. *De Buffon.* xiii. 117. *tab.* xiv. xvi. *Suppl.* iii. 183.


German Marmot. *Syn. quad.* N° 2co. *LEV. Mus.*

R. with large rounded ears: full black eyes: color on the head and back, reddish brown: cheeks red: beneath each ear a white spot, and another behind; a fourth near the hind legs:

2



1



1. Hamster — . V. 411.

2. Black Variety of the same.



legs: breast, upper part of the fore legs, and the belly, black: tail short, almost naked: four toes, and a fifth claw, on the fore feet, five behind: about nine inches long; tail three.

The males are always bigger than the females; some weigh from twelve to sixteen ounces: the females seldom exceed four or six. They vary sometimes in color. About *Casan* is found frequently a family entirely black.

SIZE.

Inhabits *Austria*, *Silesia*, and many parts of *Germany*, *Poland*, and *Ukraine*; in all the southern and temperate parts of *Russia* and *Siberia*; and even about the river *Jenesi*, but not farther to the east. They are also found in the *Tartarian* deserts, in sandy soil, disliking moist places. They are very fond of such spots which abound with liquorice, whose seeds they feed on. They swarm so in *Gotha*, that in one year 11,564, in another 54,429, and in a third 80,139 of their skins were delivered at the *Hotel de Ville* of the capital*, these animals being proscribed on account of their vast devastations among the corn.

PLACE.

They are very destructive to grain; eating great quantities, and carrying still more to its hoard: within its cheeks are two pouches, receptacles for its booty, which it fills till the cheeks seem ready to burst: the *Germans* therefore say of a very greedy fellow, *Er frisst wie ein Hamster*.

MANNERS.

They live under ground; first form an entrance, burrowing down obliquely: at the end of that passage the male sinks one perpendicular hole; the female several: at the end of these are formed various vaults, either as lodges for themselves and young, or store-houses for their food. each young has its different apart-

* *De Buffon, Suppl. iii. 185.* quoted from *Mr. Sulzer*.

ment;

ment; each sort of grain its different vault; the first they line with straw or grafs: these vaults are of different depths, according to the age of the animal; a young *Hamster* makes them scarcely a foot deep; an old one sinks them to the depth of four or five; and the whole diameter of the habitation, with all its communications, is sometimes eight or ten feet.

The male and female have always separate burrows; for excepting their short season of courtship, they have no intercourse. The whole race is so malevolent as to constantly reject all society with one another. They will fight, kill, and devour their own species, as well as other lesser animals; so may be said to be carnivorous as well as granivorous. If it happens that two males meet in search of a female, a battle ensues; the female makes a short attachment to the conqueror, after which the connexion ceases. She brings forth two or three times in a year, and brings from sixteen to eighteen at a birth. Their growth is very quick; and at about the age of three weeks, the old one forces them out of the burrows to take care of themselves: she shews little affection for them; for if any one digs into the hole, she attempts to save herself by burrowing deeper into the earth, and totally neglects the safety of her brood: on the contrary, if she is attacked in the season of courtship, she defends the male with the utmost fury.

They lie torpid from the first colds to the end of the winter; and during that time are seemingly quite insensible, and have the appearance of being dead; their limbs stiff, and body cold as ice: not even spirits of wine, or oil of vitriol, poured in to them, can produce the least mark of sensibility. It is only in places beyond the reach of the air in which it grows torpid; for the severest

cold on the surface does not affect it, as has been proved by experiment.

In its annual revival, it begins first to lose the stiffness of its limbs; then breathes deeply, and by long intervals: on moving its limbs, it opens its mouth, and makes a rattle in the throat; after some days it opens its eyes, and tries to stand; but makes its efforts like a person much concerned in liquor; at length, when it has attained its usual attitude, it rests for a long time in tranquillity, to recollect itself, and recover from its fatigue.

They begin to lay in their provisions in *August*; and will carry grains of corn, corn in the ear, and peas and beans in the pods, which they clean in their holes, and carry the husks carefully out: the pouches above mentioned are so capacious as to hold a quarter of a pint *English*. As soon as they have finished their work, they stop up the mouth of their passage carefully. As they lie torpid during the whole severe season, these hoards are designed for their support on their first retreat, and in the spring and beginning of the summer, before they can supply themselves in the fields. In winter, the peasants go what they call a *Hamster-nesting*; and when they discover the retreat, dig down till they discover the hoard, and are commonly well paid; for, besides the skins of the animals, which are valuable furs, they find commonly two bushels of good grain in the magazine. These animals are very fierce; will jump at a horse that happens to tread near them, and hang by its nose, so that it is difficult to disengage them: they make a noise like the barking of a dog. In some seasons are so numerous as to occasion a dearth of corn. Pole-cats are their greatest enemies; for they pursue them into their holes, and destroy numbers.

It is remarkable, that the hair sticks so close to the skin, as not to be plucked off without the utmost difficulty.

412. VORMELA.

In my former edition I supposed the *Vormela* of *Agricola* * to have been a variety of this kind. He says it is less; the whole body marked with yellow and tawny spots; the tail cinereous, and white tipped with black; but as he adds that it is a palm and a half long, I must refer it to another species, or perhaps genus; for it is not unlikely but that it is the same with the *Sarmatian Weefel*, N° 239.

413. YAIK.

Mus accedula. *Pallas Nov. sp. fasc. i. 257. tab. xviii. A.*
Mus migratorius. *Pallas Itin. ii. App. 703.*

R. with a thick snout: blunt nose: very fleshy lips: upper lip deeply divided: upper fore teeth small, yellow, convex outwards, truncated; the lower slender, pointed: eyes large: ears great, oblongly oval, high above the fur, naked: tail very short, cylindrical: color about the face white: upper part of the body of a cinereous yellow, mixed with brown; below of a hoary whiteness.

SIZE.

Length near four inches.

PLACE AND
MANNERS.

Inhabits the deserts about the *Yaik*: runs about during night, when it quits its burrow. It is said by the *Cossacks* to migrate in great numbers out of the deserts, and to be followed by multitudes of foxes, presaging a good hunting-season: but Doctor *Pallas* doubts whether this species, or any of the pouched kinds,

* *De anim. fabter.* 486.

go far from their homes, as those receptacles for provision are calculated only for short excursions.

Mus Phæus. *Pallas Nov. sp. fasc. i. 261. tab. xv. A.*

414. ZARIZYN.

R. with the forehead much elevated: edges of the eyelids black: ears naked, oval, standing far out of the fur: tail very short, slightly furred: color above, a hoary ash-color, with long dusky hairs, running from the neck, along the middle of the back, to the tail: the sides whitish: the circumference of the mouth, under side of the body, and the extremities of the limbs, of a snowy whiteness.

Length about three inches and a half.

Inhabits the deserts of *African*, about *Zarizyn*; and is taken in traps frequently in winter, in places near to stables and out-houses. It is also common among the *Hyrcanian* mountains, about the *Persian* villages, where it commits great ravages among the rice. It does not grow torpid during winter, as is proved by the stomachs of such which are taken in that season, being found full of food.

SIZE.

PLACE.

Mus arenarius. *Pallas Nov. sp. fasc. i. 266. tab. xvi. A. Ilin. ii. App. 704.*

415. SAND.

R. with a longish head and snout, and sharp nose: the pouches very large: ears great, oval, brownish: body short: nails white: color of the upper part of the body hoary: sides, belly, limbs, and tail, of a pure white.

E c 2

Length

SIZE.

Length near four inches; tail above one.

PLACE.

Inhabits the sandy plains of the *Baraba*, not far from the river *Irtish*. The males inhabit a very deep burrow, with a single entrance, at the bottom of which is the nest, made of the *Elymus arenarius*, and other plants: other burrows, perhaps of the females, had three entrances: in another, discovered in *May*, were five young in three nests; two were preserved alive; were untameable, very fierce, and would fling themselves on their back, and defend themselves by biting: they went out only in the night, and hid themselves during day in their fodder.

416. SONGAR.

Mus songarus. *Pallas Nov. sp. fasc. i. 269. tab. xvi. B. Itin. ii. App. 703.*

R. with a thick head and blunt nose: ears oval, very thin, appear above the fur, are very slightly clothed with hoary down: tail very short, blunt, thick, and hairy: color above, a cinereous grey, marked along the back, from head to tail, with a black line: sides of the head and body marked with great white spots in certain parts, running into one another, in others bounded with brown: belly and legs white.

SIZE.

Length three inches.


PLACE.

Inhabits, with the former, the *Baraba*, usually in the dry sandy saline places: dwells during summer in the shallow new-begun burrows; those of the females have a very deep oblique passage at the end of it: the nest formed of herbs; in one of which were seven young; from this nest ran another deep hole, perhaps the winter retreat. The young were much grown, yet blind. Doctor *Pallas* preserved them long: they grow soon familiar, contrary

2



1



1. *Sciurus* —. 1844.

2. *Sciurus* —. 1846.



to the nature of other mice; would feed from his hand, lap milk, and when placed on a table, shew no desire of running away; but were slower in all their motions than the other species. They washed their faces with their paws, and eat sitting up: wandered about in the day and morning: slept all night rolled up: seldom made any cry, and when they did, it was like that of a bat.

Mus furunculus. *Pallas Nov. sp. fasc. i. 273.*
Mus Barabensis. *Itin. ii. App. 704.*

417. BARABA.

R. with a sharp nose: large broad naked ears, dusky edged with white: tail longer than that of the preceding: color of the upper part of the body cinereous yellow, growing paler towards the sides: below of a dirty white: from the hind part of the neck extends a black line, reaching not quite to the tail: tail white, marked above with a dusky line.

Length about three inches and a quarter: tail near one inch.

Inhabits the sandy plain of *Baraba*, towards the *Ob*; and between the *Onon* and *Argun*, and about the lake *Dalai* in the *Chinese* empire. Nothing is known of their manners: the specimens from whom the descriptions were formed, were taken running about the fields.

The last division of mice is of those which lead a subterraneous life, like the Mole, which I take the liberty of naming

SIZE.
 PLACE.

*** Mole-

*** Mole-Rat.

418. BLIND.

*Mus Typhlus. Pallas Nov. Sp. fasc. i.**Lemni. Rzaczynsk. Austr. Polon. 325. De**Buffon, xv. 142.**Slepezh. Gmelin Itin. i. 131. tab. xxii.**Spalax microphthalmus. Gueldenst. Nov.**Com. Petrop. xiv. 411. tab. viii. ix.**Mus oculis minutissimis, auriculis cauda-**que nullis. Lepechen. ibid. 509. tab. xv.**Podolian Marmot. Syn. quad. N° 204.*

R. with a great head broader than the body: not the least aperture for the eyes; yet beneath the skin are the rudiments of those organs, not bigger than the seed of a poppy: no external ears: the end of the nose covered with a thick skin: nostrils very remote, and placed below: the mouth gaping, and the teeth exposed: upper fore teeth short, lower very long, and none of them hid by the lip; ends quite even: body cylindrical: limbs very short: five toes on each foot, all separated, except by a thin membrane near the base: claws short: hair universally short, thick, and very soft; dusky at the bottom, at the ends of a cinereous grey: the space about the nose, and above the mouth, white.

SIZE.

Length between seven and eight inches: weight of a male above eight ounces.

PLACE.

Inhabits only the southern parts of *Russia*, from *Poland* to the *Volga*, but is not found any where to the east of that river; but is very common from the *Sybian* to the *Sarpa*: is frequent along the *Don*, even to its origin, and about the town of *Rask*, excepting the sandy parts, for it delights in moist and turfy soils.

It

It lives in great numbers in the same places with the EARLESS MARMOTS.

It forms burrows beneath the turf for a very considerable extent, with several lateral passages made in quest of roots, on which it feeds. At the interval of some yards, there are openings to the surface to discharge the earth, which forms in those places hillocks of two yards in circumference, and of a great height. It works its way with its great teeth, and casts the earth under its belly with the fore feet, and again behind it, with its hind feet: it works with great agility; and on any apprehension of an enemy, it forms instantly a perpendicular burrow. The bite of this animal is very severe. It cannot see its assailant, but lifts up its head in a menacing attitude. When irritated, it snorts, and gnashes its teeth, but emits no cry. It often quits its hole, especially in the morning, and during the amorous season basks with the female in the sun. It does not appear that it lies torpid during winter, nor whether it lays in provision for that season. It is particularly fond of the bulbous *Chærophyllum*.

The *Russians* call it *Slepez*, or the blind: the *Cossacks*, for the same reason, style it *Sfochor Nomon*. In *Ukraine*, the vulgar believe that the touch of a hand, which has suffocated this animal, has the same virtue in curing the king's-evil, as was once believed to be inherent in the abdicated family of *Great Britain*.

419. DAUURIAN.

Mus Aspalax. *Pallas Nov. sp. fasc. i. 165. tab. x. Itin. iii. 692.*
 Mus Myofpalax. *Laxman.*

R. with a thick flat head: short snout: blunt nose, fit for digging: upper fore teeth naked; lower covered with a moveable lip: no external ears: eyes very small, yet visible, lodged deeply in their sockets, which are so minute as scarcely to admit a grain of millet: body short, and depressed: limbs very strong, especially the fore legs: fore feet large, and adapted for digging; naked, and furnished with five toes, and very long and strong claws, slightly bent, on the three middle: hind feet naked to the heel; on each are five toes with small claws: tail short: hair soft, and loose: color at bottom dusky, outwardly of a dirty cinereous grey: in some is a white line on the hind part of the head.

SIZE.

Differs in size. Those of the *Altaic* chain are near nine inches from nose to tail: those about lake *Baikal* not six: the tail of the former is near two inches long.

PLACE.

Inhabits, first, the *Altaic* mountains; and again beyond lake *Baikal*, and from thence for some space southward; but none are found to the north. In the former it lives on the bulbs of the *Erythronium*; in the latter on those of the *Lilium Pomponium*.

It burrows like the former, a little below the surface, and spreads over an extent of a hundred fathoms; and the direction it takes is known by the number of hillocks.

Its voice is weak and plaintive. It digs with both nose and fore feet; but less than the preceding with the teeth: by comminuting the earth, and flinging it up in hillocks, it prepares the ground

ground for the reception of various kinds of rare seeds; which grow usually in greater plenty about such places than any others.

The *Tanguts*, about lake *Baikal*, call this species *Monon Zokor*, or blind; yet it is not quite deprived of sight. The *Russians* style it *Semiunaja Medwedka*, or *Earth Bear*.

R. with a large head: nose black; end flattened and corrugated: eyes minute, much hid in the fur: no ears: upper teeth one-third of an inch long, fulcated lengthways; lower, one inch and a quarter, exposed to view: legs short: on the fore legs are four toes and a thumb, detached and free: inmost toe the longest, the others gradually shorten: on the thumb is a short claw; the other claws are very long, and slightly bent: the soles are naked, and distinguished by two great tubercles: hind feet very long, large, and naked, which the animal rests on even to the heel: they have five toes with short claws.

420. AFRICAN.

Tail compressed, and covered above and below with short hairs: on the sides beset with very long bristles disposed horizontally.

Color a cinereous brown, palest on the lower parts.

Length to the tail thirteen inches: tail two.

SIZE.

PLACE:

Inhabits the sandy country near the *Cape of Good Hope*, where it is called *Sand Moll*. It burrows, and flings up hillocks, like the former; and renders the ground so hollow, as to be very inconvenient to travellers; for it breaks every six or seven minutes under the horses feet, and lets them in up to the shoulders. This animal feeds on the roots of *Ixia*, *Gladioli*, *Antholyzæ*, and *Irides*;

VOL. II.

F f

grows

grows to the size of a rabbit, and is by some esteemed a good dish *. This, from its superior size, I suppose to be the *Sand Moll* of Mr. *Missou*.

421. CAPE.

Mus Capensis. Pallas Nov. sp. fasc. i. 172.
tab. vii.

xlv.

Long-toothed Marmot. *Brown's Zool. tab.*

La Taupe du Cap. *Journal hist. fig. 64.*




R. with a blunt nose: minute round nostrils: eyes small, but larger than those of the preceding: no ears: upper fore teeth contiguous, truncated; lower, an inch long, not contiguous, bend upwards, excavated on the upper surface: end of the nose naked and black, the rest white: chin, and lower sides of the cheeks, of the same color: space round the ears and eyes white: on the hind part of the head is a white spot; rest of the head, cheeks, back, and sides, of a rusty brown, and cinereous: belly cinereous: five slender toes on each foot, furnished with small claws: tail very short, beset with bristles.

SIZE.


Length, from nose to tail, about seven inches.

Is very common about the *Cape*, and very destructive to gardens; flings up hillocks, and eats roots of various kinds.

* *Masson's trav. Ph. Transf. lxxvi. 304. De la Caille, 292.*



2. Blind Mole-Rat . . . 1. 418.
 3. Laurian ————— . 1. 419.
 1. Cape ————— . 1. 421.



Mus Talpinus. *Pallas Nov. Sp. fasc. i. 176. tab. xi. B. Nov. Com. Petrop. xiv. 563. tab. xxi. fig. 3.*

422. TALPINE.

R. with a large short head: thick snout: nose truncated: upper teeth extending out of the mouth, long and flat: lower longer, rounded at the ends: eyes small, hid in the fur: no ears: the aperture bounded behind by a small rim: body short: fore feet strong: on those, and on the hind feet, five toes furnished with small claws: tail very short, scarcely appearing beyond the fur: color of the head, nose, back, and sides, dusky: cheeks greyish: chin white: belly and limbs whitish.

Length near four inches.

Inhabits all the open grounds and commons of the temperate parts of *Russia* and western *Siberia*, but scarcely any beyond the *Irtish*, and none as far as the *Oby*.

SIZE.

PLACE.

Loves a black turfy soil, and is frequent in meadows near villages: seldom in sandy or mudded tracts: always abound where there is plenty of *Phlomis tuberosa*, and *Lathyrus esculentus*. Its place is known by the little hillocks it flings up along the course of its burrow, which is of great extent; for this reason the *Russians* call it *Semleroiika*, or *Earth-digger*. In these burrows it lurks all the day, but in evening and morning renews its labors; nor does it quit its hole unless to fling out the earth, or in the season of love to seek a mate, or to change the place of its habitation. It does not bear the full light of day; therefore its few excursions are usually in the evenings.

MANNERS.

It does not grow torpid in winter; but makes its nest beneath some shrub or hay-rick, and deep in the ground, and keep themselves

felves warm by lining it with soft grafs: and often make a lodge, which they fill with tuberous roots. During the cold feafon their fur grows univerfally thicker and longer.

It is very eafily taken: but foon grows fick in confinement, unlefs a quantity of earth is put into the place. They emit a puling note, but that rarely: they often gnafh, and, as it were, whet their teeth againft each other.

They are in heat the end of *March*, or beginning of *April*; at that time the females have a ftrong musky fmell. They bring three or four at a time.

They fometimes vary in color, and are found quite black.

Two cutting teeth in each jaw, pointing forward.

Long slender nose: small ears.

Five toes on each foot.

XXXIV.
SHREW.

- Mus aquaticus. Clusii exot.* 373. *Worm.* Dæfman, *Faun. fusc.* No. 28. *De Buffon,*
Muf. 334. x. 1.
Muscovy or Musk rat. Raii syn. quad. 217. *Castor cauda verticaliter plana, digitis*
Nov. Com. Petrop. iv. 383. *omnibus membranis inter se connexis.*
Castor moschatus. C. cauda longa com- *Briffon quad.* 92
presso-lanceolata, pedibus palmatis. Long-nosed Beaver. *Syn. quad.* N^o 192.
Lin. fyfl. 79.

423. MUSKY:

S. with a long slender nose, like that of a shrew-mouse: no external ears: very small eyes; tail compressed sideways: color of the head and back dusky; the belly whitish ash-color: length, from nose to tail, seven inches; tail eight.

Inhabits the river *Volga* and lakes adjacent, from *Novogorod to Saratof*; never found in *Russia*, and its existence in *Lapland* doubted*. Never goes upon dry land, but wanders from lake to lake, only in fortuitous floods: is often seen swimming or walking under the water: comes up for air to the surface, or in clear weather sporting on the surface: loves stagnating waters with high banks, in which it makes burrows twenty feet long: feeds on leeches, and the *larvæ* of water insects: a few fragments of roots have also been found in the stomach. Is not torpid during winter, being often in that season taken in nets†. Is very slow in its pace: makes holes in the cliffs, with the entrance far beneath the lowest fall of the

* *Dr. Pallat, MSS.*

† The same.

water;

water; works upwards, but never to the surface, only high enough to lie beyond the highest flow of the river: feeds on fish: is devoured by the Pikes and *Siluri*, and gives those fish so strong a flavor of musk, as to render them not eatable: has the same scent as the former, especially about the tail: out of which is expressed a sort of musk, very much resembling the genuine kind*. The skins are put into chests among cloaths, to drive away moths†, and to preserve the wearers from pestilence and fevers.

At *Orenburg*, the skins and tails sell for fifteen or twenty *copecks* per hundred. They are so common near *Nisney Novogorod*, that the peasants bring five hundred apiece to market, where they are sold for one ruble per hundred. The German name for these animals is *Biesem-ratze*; the Russian, *Wychozhol*.

424. PERFUMING.

Mus Pilorides? *Pallas Nov. sp. fasc.* i. 91.

Mus albus Ceylonicus? *Brisson*, 122. LEV. MUS.


S. with a long slender nose: upper jaw extending far beyond the lower: upper fore teeth short: lower long, slender, incurvated: whiskers long and white: eyes small: ears transparent, broad, and round: hair short and close, on head and body, of a fine pale cærulean: the belly lighter: feet naked and pink-colored.

Length from nose to tail near eight inches; tail three and a

* *Scheber* in *Müller's Samlung Russ.* vii. 41. 42.

† *Ratckoff Orenb. Tepegr.* i. 286.

half;



Peromyscus leucopus. V. 24.



half: quite naked, round, thick at the base, tapering to a point; and of the same color with the feet.

Inhabits *Java*, and others of the *East Indian* islands; eats rice; has so strong a scent of musk as to perfume every thing it runs over. I have it from the most undoubted authority, that it will render the wine in a well-corked bottle not drinkable, by merely passing over it. Cats will not touch them.

Tucan. *Hernandez Nov. Hsp.* 7.

Le Tucan. *De Buffon*, xv. 159.

425. MEXICAN.

S. with a sharp nose: small round ears: without sight: two long fore teeth above and below: thick, fat, and fleshy body: short legs, so that the belly almost touches the ground: long crooked claws: tawny hair: short tail: length, from nose to tail, nine inches.

Inhabits *Mexico*: burrows, and makes such a number of cavities, that travellers can scarcely tread with safety: if it gets out of its hole, does not know how to return, but begins to dig another: grows very fat, and is eatable: feeds on roots, kidney-beans, and other seeds. *M. de Buffon* thinks it is a Mole; but by the ears, it should be classed here.

Mus araneus figura muris. *Marcgrave* La musaraigne de Brasil. *De Buffon*, xv. 160.

426. BRASILIAN.

S. with a sharp nose and teeth: pendulous scrotum: of a dusky color, marked along the back with three broad black strokes: length, from nose to tail, five inches; tail two.

Inhabits *Brasil*: does not fear the cat: neither does that animal hunt after it.

427. MURINE.

S. murinus. *S. cauda mediocri*, corpore fusco, pedibus caudaque cinereis, *Lin.*
syft. 74.

S. with a long nose, hollowed beneath: very long hairs about the nostrils: ears rounded, and rather naked: of an ashy-color: body of the size of a common mouse: tail a little shorter than the body, and not so hairy.

Inhabits *Java*.

428. FÆTID.

Μυγανν. *Ælian hist. An. lib. vi. c. 22.*

Μυγανν. *Dioscorid. lib. ii. c. 42.*

Mus araneus. *Agricola An. Subter. 485.*

Gesner quad. 747.

Mus araneus, mus cæcus. *Gesner icon.*

116.

Mus araneus, Shrew, Shrew-mouse, or hardy Shrew, *Rail syn. quad. 233.*

Mus araneus rostro productiore Spitzmaus. *Klein quad. 57. Kramer Austr.*

317.

Sorex araneus. S. cauda mediocri, corpore subtus albido. *Lin. syft.* 74.

Nabbus. Faun. suec. No. 24.

Mus araneus supra ex fusco rufus, infra albicans. *Briffon quad. 126.*

La Musaraigne. De Buffon, viii. 57. tab. x.

Shrew-mouse. Er. Zool. i. 112.

S. with short rounded ears: eyes small, and almost hid in the fur: nose long and slender, upper part the longest: head and upper part of the body of a brownish red: belly of a dirty white: length, from nose to tail, two inches and a half; tail one and a half.

PLACE,

Inhabits *Europe*, *Siberia*, and even the *Arctic* flats, and *Kamtshatka*; it is also found about the *Caspian* sea; lives in old walls, heaps

heaps of stones, or holes in the earth: is frequently near hay-ricks, dunghills, and necessary-houses: lives on corn, insects, and any filth: is often observed rooting in ordure, like a hog: from its food, or the places it frequents, has a disagreeable smell: cats will kill, but not eat it: brings four or five young at a time. The antients believed it was injurious to cattle, an error now detected. There seems to be an annual mortality of these animals in *August*, numbers being then found dead in the paths.

Mus araneus dorso nigro, ventreque albo.

Merret Pinax, 167.

Sorex fodiens. Pallas.*

La Musaraigne d'Eau. De Buffon, viii.

64. tab. xi.

Water Shrew-mouse. Br. Zool. illustr.

tab. cii. LEV. MUS.

429. WATER.

S. with a long slender nose: very minute ears; and within each a tuft of white hairs: very small eyes, hid in the fur: color of the head and upper part of the body black: throat, breast, and belly, of a light ash-color: the feet white: beneath the tail a triangular dusky spot: much larger than the last: length, from nose to tail, three inches three quarters; tail two inches.

Inhabits *Europe* and *Siberia*, as far at least as the river *Jenesei*; long since known in *England*, but lost till *May 1768*, when it was discovered in the fens near *Revesby Abby, Lincolnshire*: burrows in the banks near the water; and is said to swim under wa-

PLACE.

* Doctor *Pallas* favored me with several prints of this animal in 1765, but never published them: he discovered it near *Berlin*: it is called there *Græber*, or The Digger.

VOL. II.

G g

ter:

ter*: is called by the Fen-men the *Blind Mouse*: chirrups like a grasshopper, and its note often mistaken for one.

430. ELEPHANT.

S. with a very long, slender and little nose: the whole animal of a deep brown color.

PLACE.

Inhabits the neighborhood of the Cape of *Good Hope*: called the *Elephant*, from its *proboscis*-like snout: engraven from a drawing by Mr. *Paterfon*.* This animal has been very ill represented by *Petiver* in his *Gazoph.* Dec. iii. tab. xxiii. fig. 9. under the title of *Mus araneus maximus Capensis*.

Sorex marinus. Gm. Lin. 114.

431. MARINE.

S. with elongated snout, channel'd below: ears rounded, and naked: fur of a dusky color; whiskers grey: tail a little shorter than the body: size of the common mouse.


Inhabits *Java*.

432. SURINAM.

S. with the upper part of the body bay; the lower pale ash, mixed with yellow: tail one half shorter than the body.

Inhabits *Surinam*.

* *L. Baldner*, iii. 137.



Elephant Shrew. —. N.º 30



Sorex pusillus, *Erxleben*, 122. *Gm. Lin.* 114.

433. PERSIAN.

S. with the body hoary above, cinereous beneath: tail (*subdisticha*) short, and whitish: length of the body three inches seven lines; tail one inch one line.

Inhabits the north of *Persia*: burrows and lives below ground.

Sorex minutus. *S. rostro longissimo*. *Lin. Syst.* 73.

434. MINUTE.

S. with a head near as big as the body: very slender nose: broad short naked ears: whiskers reaching to the eyes: eyes small, and capable of being drawn in: hair very fine and shining; grey above, white beneath: no tail.

Inhabits *Siberia*, about the *Oby* and near the *Kama*: lives in a nest made of *lichens*, in some moist place beneath the roots of trees: lives on feeds: digs: runs swiftly: has the voice of a bat.

Sorex exilis. *Gm. Lin.* 115.

435. PYGMY.

S. with a very long slender nose: in shape and color like the **FOETID**, but paler: the tail very slender near the roots, then suddenly grows remarkably thick and round; and again grows gradually less to the end.

LINNÆUS imagines that the last is the left of quadrupeds. **DOCTOR PALLAS**, who communicated this species, thinks this has

a better clame to that title, as its weight is only equal to, or very little above half a drachm.

Is very common between, and about the rivers *Jeneſei* and *Oby*.

436. WHITE-
TOOTHED.

S. of a dusky cinereous color: belly white: cutting teeth white: tail slender and hairy.

437. SQUARE-
TAILED.

S. of a dusky cinereous color: belly paler: cutting teeth brownish: tail inclines to a squared form.

This species has no bad smell.

438. CARINATED.

S. of a dusky cinereous whitish on the belly, with brownish fore teeth: a white spot beyond each eye: tail slender and taper, carinated or ridged below.

439. UNICOLOR.

S. of an uniform dusky cinereous color: base of the tail narrow, or compressed.

PLACE.

The above four species inhabit the neighborhood of *Strasbourg*, and were discovered by Professor *Herman*.

Long nose: upper jaw much longer than the lower.

No ears.

Fore feet very broad, with scarcely any apparent legs before:
hind feet small.

XXXV.
MOLE.

Talpa. *Agricola An. Subter.* 490. *Gesner*
quad. 931. *Klein quad.* 6

Talpa, the Mole. Mold-warp, or Want.

Rain. syn. quad. 236.

Kret. *Rzeczynski Polon.* 236.

Scheer, Scheer-maus. Maul-wurf. *Kra-*
mer Austr. 314.

Talpa Europæus. T. caudata, pedibus

pentadactylis. *Lin. syst.* 73.

Mullvad, Surk. *Faun. suec.* No. 23. *Br.*
Zool. i. 108.

Talpa caudata, nigricans pedibus anticis
et posticis pentadactylis. *Briffon quad.*
203.

La Taupe. *De Buffon*, viii. 81. *tab.* xii.
Lev. Mus.

440. EUROPEAN.

M. with very minute eyes, hid in the fur: long snout: six cutting teeth in the upper, eight in the lower jaw, and two canine in each: no external ears, only an orifice: fore part of the body thick and muscular; hind part taper: fore feet placed obliquely, broad, and like hands: five toes, each terminated by strong claws: hind feet very small, with five toes to each: tail short: skin very tough, so as scarcely to be cut through: hair short, close set, softer than the finest velvet: usually black, sometimes spotted* with white; sometimes quite white: length five inches three quarters; tail one.

Inhabits *Europe*, and the temperate or southern parts of *Russia* and *Siberia*, as far as the River *Lena*. The *Siberian* is much larger than the *European Mole*.

PLACE.

* Spotted Mole, *Edw.* 268.

It


MANNERS.

It lives under ground: burrows with vast rapidity with its fore feet; flings the earth back with its hind feet: has the sense of smelling exquisite, which directs it to its food—worms, insects, and roots: does vast damage in gardens, by flinging up the soil and loosening the roots of plants: is most active before rain, and in winter before a thaw, worms being then in motion: breeds in the spring: brings four or five young at a time: makes its nest of moss, a little beneath the surface of the ground, under the greatest hillock: raises no hillocks in dry weather, being then obliged to penetrate deep after its prey: makes a great scream when taken. *Palma Christi* and white *hellebore*, made into a paste, and laid in their holes, destroys them. None in *Ireland*.

β. **YELLOW M.** in form resembling the *European*, but larger, being six inches two-tenths long; the tail one inch: hair soft, silky, and glossy, of a yellowish brown color at the ends; dark grey at the roots: brightest about the head; darkest about the rump: belly of a deep cinereous brown: feet and tail white.

Inhabits *N. America*. Described from a skin in which the jaws were taken out.





Cape . Mole . L. 3. 3. 11

- Talpa Sibiricus verficolor, *Aspalax* dictus. anticistridactylis, posticistetradactylis.
Seb. Mus. i. 51. tab. xxxii. fig. 4, 5. *Brisson quat.* c6.
Klein quad. 65. La Taupe dorée. *De Buffon*, xv. 145.
 Talpa Asiatica. T. ecaudata, palmis tri- Variable Mole. *Brown's Zool.* 118. tab.
 daetylilis. *Lin. Syst.* 73. 44.
 Talpa ecaudata, ex viridi aurea, pedibus

441. SIBERIAN.

M. with the nose short and blunt: space between the tip, and corner of the mouth covered with pale brown hair: from the corner of the mouth, a broad whitish bar points upwards along the sides of the head: color of the hair on the upper part of the body varied with glossy green and copper-color: below is of a cinereous brown: in the upper jaw are two sharp cutting teeth; in the lower the same, with a sharp canine tooth contiguous to them on each side.

TEETH.

On the fore feet three toes with vast claws; that on the outmost toe exceedingly large: on the hind feet five small toes and weak claws: no tail: rump round.

Length four inches.

SIZE.

Inhabits the *Cape of Good Hope*, not *Siberia*, as *Seba* supposes: Whether this is the *Bles Moll* of the *Dutch*, which lives in the harder grounds about the *Cape**, I cannot determine.

PLACE.

* *Maffon's Trav. Ph. Transf.* lxxvi. 305.

442. RADIATED.

Sorex cristatus. S. naribus carunculatis, cauda brevior. Lin. Syst. 73.
 LEV. MUS.

M. with small but broad fore legs; five long white claws on each: nose long; the edges beset with radiated tendrils: hair on the body dusky, very short, fine, and compact; on the nose longer: the hind legs scaly: five toes on each foot: length, from nose to tail, three inches three quarters: tail slender, round, and taper; one inch three-tenths long.

PLACE.

Inhabits *N. America*. Forms subterraneous passages, in different directions, in uncultivated fields; raises walks about two inches high and a palm broad: the holes often give way and let in the walkers. Feeds on roots: has great strength in its legs.

443. LONG-TAIL-
ED.

M. with a radiated nose: the fore feet pretty broad, hind feet very scaly, with a few short hairs on them: the claws on the fore feet like those of the common Mole; on the hind very long and slender: hair on the nose and body soft, long, and of a rusty brown color: tail covered with short hair; the length two inches; that of nose and body four inches six-tenths.



PLACE.

Inhabits *N. America*. LEV. MUS.

444. BROWN.

Sorex aquaticus. S. plantis palmatis, palmis caudaque brevior albis.
Lin. Syst. 74. LEV. MUS.

M. with a slender nose: upper jaw much longer than the lower; two cutting teeth in the upper, four in the lower, the two middle of which are very small: no canine teeth: fore feet



1. *Radiated Mole* —. M. 242.

2. *Long-tailed Mole* —. M. 243.



feet very broad: nails long: hind feet small; five claws on each: hair very soft and glossy, brown at the ends, deep grey at the bottom: tail and feet white: length, from nose to tail, five inches and a half: tail very slender, not an inch long.

Inhabits *N. America*: called there the Brown Mole: sent from *New York* by Mr. *A. Blackburne*, with β . Yellow Mole, and No. 442 and 443. The black and shining purple *Virginian* Mole, described by *Seba** as the same with the common kind, was not among those that gentleman favoured us with. *Linnaeus* places this, and our radiated Mole, in his class of *Sorex*, or *Shrew*, on account of the difference of the teeth; but as these animals possess the stronger characters of the *Mole*, such as form of nose and body, shape of feet, and even the manners, we think them better adapted to this genus than to the preceding.

PLACE.

Talpa rubra Americana. Seb. Mus. i. 51. tab. xxxii. fig. 2.

445. RED.

M. of a cinereous red color: three toes on the fore feet, four on the hind: form of the body and tail like the *European* kind.

According to *Seba*, it inhabits *America*; but he does not inform us whether it is *North* or *South*.

* *L. 51. tab. xxxii. fig. 4.*

XXXVI.
HEDGE-HOG.

Five toes on each foot.
Body covered with strong short spines.

416. COMMON.

Erinaceus, *Agricola An. Subter.* 481.
Echinus terrestris, *Gesner quad.* 368.
Echinus fr. Erinaceus terrestris, Urchin,
or Hedge-hog. *Rail syn. quad.* 231.
Jez. *Kzeczinski Poloz.* 233.
Acanthion vulgaris nostras, *Kleinquad.* 66.
Igel. *Kramer Duffr.* 314.
Erinaceus Europeus. E. auriculis rotun-

datis naribus cristatis. *Lin. syst.* 75.
Igelkott. *Faun. succ.* N^o 22. *Br. Zool.* i.
106.
Erinaceus auriculis erectis. *Briffon quad.*
128. *Sib. Mus.* i. 78. *tab.* xlix.
L'Herisson. *De Buffon*, viii. 28. *tab.* vi.
Hærbe, vel Ganfud. *Førskæl*, iii. *LEV.*
Mus.

H. with a long nose: nostrils bordered on each side with a loose flap: ears rounded, short, broad, and naked: eyes small: legs short, naked, and dusky: inner toe the shortest: claws weak: upper part of the face, the sides, and rump, covered with strong coarse hair of a yellowish and cinereous color; the back, with strong sharp spines of a whitish color, with a bar of black through their middle: tail an inch long: length, from nose to tail, ten inches.

PLACE.

Inhabits *Europe* and *Madagascar**; is common in many parts of *Russia*, but scarcely or ever found in *Siberia*: is in motion during night; keeps retired in the day: feeds on roots, fruits, worms, and insects: erroneously charged with sucking cows and hurting their udders: resides in small thickets, in hedges, and at the bottom of ditches covered with bushes: lies still and curled up in

* *Pl. com. rep. Madagascar*, 52, et *reth y d'Arabie* V. *Sua.*

moſs, graſs, or leaves, and during winter rolls itſelf up and ſleeps out that dreary ſeaſon: a mild and helpleſs animal; on approach of an enemy, rolls itſelf into the form of a ball, and is then invulnerable.

Erinaceus Auritus. Pallas & Gmelin, in *Nov. com. Petrop.* xiv. 519. 573. tab. xvi. and xxi. fig. 4.

447. SIDERIAN.

H. with the upper jaw long and ſlender: with very large open oval ears, naked, brown round the edges, with ſoft whitish hairs within: tail ſhorter than that of the common hedge-hog: upper part of the body covered with ſlender brown ſpines, encompassed at the baſe, and near the ends, with a ring of white: the limbs and belly cloathed with a moſt elegant ſoft white fur.

Generally much inferior in ſize to the common kind; but beyond *Baikal* is found much larger than that ſpecies.

Is very common in all the ſouthern deſerts, from the *Don* to the *Oby*.

PLACE.

Grows very fat: ſleeps all the winter, lodged in a hole a few inches deep: lives on inſects, even the moſt cauſtic, and will eat (as experiment has been made) above a hundred *Cantharides* without any injury: rolls itſelf up, and has all the manners of the common kind.

418. ASIATIC. Le petit Tandre. *Sounerat, voy. ii. 146.* Le Tendrac, et Le Tanrec. *De Buffon, tab. xviii. xii. 438.*

H. with a long slender nose: short rounded ears: short legs: the body marked longitudinally with five broad lines of black, and the same of white; which are continued over the shoulders and thighs: the white marks consist of short spines; the black marks are furnished with long loose hairs, which fall quite to the ground: head and face quite black: no tail: length seven inches. *M. de Buffon* has given the figure of a young one.

The other, or the *Tanrec*, is rather larger: covered with spines only on the top and hind part of the head, the top and sides of the neck, and the shoulders; the longest were on the upper part of the neck, and stood erect: the rest of the body was covered with yellowish bristles, among which were intermixed some that were black, and much longer than the others. Each of these animals, which are varieties or young of the same species, had five toes on each foot.

PLACE.

Inhabit the isles of *India*, and that of *Madagascar*: are, when of their full growth, of the size of* rabbits: grunt like hogs: grow very fat: multiply greatly: frequent† shallow pieces of fresh or salt water: they burrow on land: lie torpid during six months, during which time their old hair falls off. Their flesh is eaten by the *Indians*, but is very flabby and insipid.

* *Dutch voy. East Indies, 203.* Those in the cabinet of the *French King* were much smaller; probably young.

† *Cauche voy. Madagascar, 53. Flacourt hist. Madagascar, 152.*



1. Common Hedge-Hog — . N. 46.

2 O Asiatic ——— . N. 48.



American Hedge-hog. *Bancroft Guiana*, *Liv. fig. 75. Erisson quat. 131.*
 144. *Erinaceus Americanus albus. Seb. Musf.*
Erinaceus inauris. E. auriculis nullis. i. 78. tab. fig. 3.


449. GUIANA.

H. without external ears, having only two orifices for hearing: has a short thick head: back and sides covered with short spines of an ash-color, tinged with yellow: face, belly, legs, and tail, covered with soft whitish hair: above the eyes, of a chestnut color; the hind part and sides of the head of a deeper color: length, from nose to tail, eight inches: tail short: claws long and crooked.

Inhabits *Guiana*.

PLACE.

DIV.



DIV. II. SECT. IV.

DIGITATED QUADRUPEDS:

Without Cutting Teeth.

DIV. II. SECT. IV. Digitated Quadrupeds.

XXXVII.
S L O T H.

Without cutting teeth in either jaw.
 With canine teeth and grinders.
 Fore legs much longer than the hind.
 Long claws.

450. THREE-
TOED.Ardopithecus. *Gesler quad.* 869. *Icon.*
quad. 65.Ignavus five per $\alpha\iota\lambda\phi\gamma\gamma\omega\iota$ Agilis. *Chyf.*
exot. 110. 372.Ai, five Ignavus. *Marsgrave Brasl.* 221.Sloth. *Raii syn. quad.* 245. *Edw.* 310.Ignavus Americanus, nulum sletu iniscens.
Klein quad. 43.Tardigradus pedibus anticis & posticis
tridactylis. *Briffon quad.* 21.Ai, five Tardigradus gracilis Americanus. *Seb. Mus.* xxxiii. *fig.* 2. *Schreber,*
ii. 7. *tab.* lxiv.Ouaikaré, Pareffieux. *Barrere France*
Æquin. 154.Bradypus tridactylus. B. pedibus tridac-
tylis cauda brevi. *Lin. syst.* 50.L'Ai. *De Buffon*, xiii. 44. *tab.* v. vi.
Br. Mus. LEV. MUS.

S with a blunt black nose, a little lengthened: very small ex-
 ternal ears: eyes small, black, and heavy; from the corner
 of each a dusky line: color of the face and throat a dirty white:
 hair on the limbs and body long and very uneven, of a cinereous
 brown color, with a black line along the middle of the back:
 each side, about the shoulders, is dashed with rust-color; the rest
 of the back and limbs spotted irregularly with black. The
 young, such as I suspect that to be in the *British Museum*, have
 few or no spots. Tail short, a meer stump: legs thick, long, and
 awkwardly



Three-toed Sloth — S. 45.



awkwardly placed: face naked: three toes, and three very long claws on each foot.

It grows, as *Nieuboff* remarks, to the bulk of a middle-sized fox*.

Inhabits most parts of the eastern side of *South America*: the most sluggish and most slow of all animals; seems to move with the utmost pain; makes a great progress if it can go a quarter of a league in a day†: ascends trees, in which it generally lives, with much difficulty: its food is fruit, or the leaves of trees; if it cannot find fruit on the ground, looks out for a tree well loaded, and with great pains climbs up: to save the trouble of descending, flings off the fruit, and forming itself into a ball, drops from the branches; continues at the foot till it has devoured all; nor ever stirs, till compelled by hunger‡: its motion is attended with a most moving and plaintive cry, which at once produces pity and disgust, and is its only defence; for every beast of prey is so affected by the noise, as to quit it with horror||: its mouth is never without foam: its note, according to *Kircher*, is an ascending and descending *hexachord*§, which it utters only by night: its look is so piteous as to move compassion; it is also accompanied with tears, which dissuade every body from injuring so wretched a being: its abstinence from food is remarkably powerful; one that had fastened itself by its feet to a pole, and was so suspended cross two beams, remained forty days without meat,

SIZE.

PLACE.

MANNERS.

* *Nieuboff's trav. Churchill's collect.* ii. 18.

† *Gamilla Orenoque*, ii. 13.

‡ *Ulloa's voy.* i. 103.

|| *Ibid.*

§ *Kircher's Musurgia*, as quoted by Mr. STILLINGFLEET, in his miscellaneous tracts, p. 100.

drink, or sleep*: the strength in its feet is so great, that there is no possibility of freeing any thing from its claws, which it happens to seize on. A dog was let loose at the above-mentioned animal, when it was taken from the pole; after some time the *Sloth* layed hold of the dog with its feet, and held him four days, till he perished with hunger†.

451. TWO-TOED. *Tardigradus Ceilonicus* fœmina. *Seb.* *Tardigradus* pedibus anticis didactylis, posticis tridactylis. *Briffon quad.* 22. *Mus.* i. tab. xxxiv.
Bradypus didactylus. *Br.* manibus didactylis cauda nulla. *Lin. syst.* 51. *L'Unau. De Buffon,* xiii. 34. tab. i. *Br. Mus.*
Schreber, ii. 10. tab. lxx.

S with a round head: short projecting nose: ears like the human, lying flat to the head: two long strong claws on the fore feet, three on the hind: hair on the body long and rough; on some parts curled and woolly: in some, of a pale red above, cinereous below; in others, of a yellowish white below, cinereous brown above. No tail. Length of that in the *British Museum* eleven inches: I believe a young one.


PLACE.

Inhabits *South America* and the isle of *Ceylon*. The last is strenuously denied by *M. de Buffon*, who has fixed the residence of this genus to *America* only: but, besides the authority of *Seba*, who expressly says his specimen was brought from *Ceylon*, a gentleman, long resident in *India*, and much distinguished in the literary world, has informed me he has seen this animal brought from the *Paliacat* mountains that lie in sight of *Madras*; which satisfies me that it is common to both continents. Farther enquiry is desired into the identity of this species.

* *Kircher.*

† *Ibid.*

There



Ursiform Noth — . V. 452.

There is reason to think that it is met with also in *Guinea*, or at least some species of this genus; for *Barbot* and *Bosman* describe an animal by the name of *Potto*, to which they give the attributes of the former, and describe as being grey when young, red, and covered with a sort of hair as thick set as flocks of wool. Both these writers were sensible men, and, though not naturalists, were too observant of the animals of *Guinea* to mistake one whose characters are so strongly marked as those of the *Sloth* *.

Bradypus uriformis. Naturalists Miscellany. tab. 58.

452. URSIFORM.

S. with a long and strong nose, truncated at the end: the forehead rises suddenly above it: that and the nose whitish, and almost naked: eyes very small; above is a black line: ears short, and lost in the hair: the hair on the top of the head points forward, that in the neck is parted in the middle; on head and neck, back and sides, is extremely long, shaggy and black; in most parts twelve inches long, and on the upper part of the body shines in the sun with a most brilliant purple gloss; on the breast and belly short; across the first is a line of white: the tail is only five inches long, and is quite hid in the hair: the limbs are very strong and bear-like: on each foot are five toes: on those of the fore feet the claws are three inches long, pointing forward, and slightly incurvated; pointing forward and admirably adapted for digging or burrowing: the claws of the hind feet are very short: the bottoms

HAIR.

* *Bosman*, 237. *Barbot*, 212.

of the feet are black and naked. This animal wants the *incisores*, or cutting teeth, above and below. In each jaw are two canine teeth, remote from the grinders: the roof of the mouth is marked with transverse fulci: the tongue is smooth, and not so long as the mouth.

LIPS. The nostrils are transverse, and appear like a narrow slit: the lips are very loose, and capable of being protruded to a great length, and drawn in again; they serve the use of a hand, and by their means it conveys apples or any sort of food, into its mouth: its principal food was vegetables, and also milk: it was very fond of honey, sugar, and other sweets; but did not willingly eat any animal food.

MANNERS. In its manners it was gentle, and very good natured; it suffered me to put my hand far down its mouth to examine the inside, and to tumble it up and down, to examine the different parts; nor did it ever offer to bite: it did no more than emit a short abrupt roar when I had provoked it highly.

I class it, from the teeth, among the *Bradypi*, or *Sloths*, not from its inactivity, or any of its natural properties: it was neither slow nor languid, but was moderately lively: it appeared to have a habit of turning itself round and round, every now and then, as if for amusement, in the manner of a dog about to lie down to sleep: it is said to have a strong propensity to burrowing; and that it was first dug out of its retreat by those who discovered it.

PLACE. It inhabits *Bengal*, and lives in certain sand hills not remote from *Patna*. It was about the size of a black *American* bear, not half grown. When I saw this animal in 1790 it was between four and five years old, so probably had attained its full growth.

I saw it in company with the ingenious Doctor *Shaw*, of the
British

British Museum. My figure is copied from his *Naturalists Miscellany*: but it was before engraved by Mr. *Catton* in his book of *Quadrupeds*. Mr. *Bewick* has also given a very good figure of it at p. 266 of his beautiful *History of Quadrupeds with wooden plates*.

XXXVIII.
ARMADILLO.

Without either cutting teeth or canine teeth.

Head, and upper part of the body, guarded by a crustaceous covering; the middle with pliant bands, formed of various segments, reaching from the back to the edges of the belly.

453. THREE-
BANDED.

Tatu apara. *Margrave Brasl.* 232. *Raii syn. quad.* 234.

Armadillo seu Tatu genus alterum. *Chuf.*

Exot. 109. *Klein quad.* 48.

Tatu seu Armadillo orientalis. *Seb. Mus.* i. tab. xxxviii. fig. 2, 3.

Dafypus tricinctus. D. cingulis tribus,

pedibus pentadactylis. *Lin. syst.* 53.

Cataphraetus scutis duobus cingulis tribus. *Briffon quad.* 24.

L'Apar, ou le Tatou a trois bandes. *De Buffon*, x. 206. *Schreber*, ii. 28. tab. lxxii. A. lxxvi. fig. 1. 2.

A with short but broad rounded ears: the crust on the head, back, and rump, divided into elegant pentangular tuberculated segments: three bands in the middle: five toes on each foot: short tail.

PLACE AND
MANNERS.

The whole genus inhabits *South America*: the manners of all much the same: burrows under ground; the smaller species in moist places, the larger in dry, and at a distance from the sea: keeps in its hole in the day, rambles out at night: when overtaken, rolls itself into the form of a ball, which it does by means of the pliant bands on its middle, and thus becomes invulnerable: when surprized, runs to its hole, and thinks itself secure if it can hide its head and some part of its body. The *Indians* take it by the tail, when the animal fixes its claws in the earth so strongly that there is no moving it till the *Indian* tickles

it

it with a stick: is hunted with little dogs, who give notice to their master of its haunts by barking, who digs it out; to take it out incautiously is very dangerous, on account of the snakes that commonly lurk in the burrows. Feeds on potatoes, melons, and roots, and does great damage to plantations: drinks much: grows very fat, and is reckoned very delicious eating when young; but when old, has a musky disagreeable taste: is very numerous; breeds every month, and brings four at a time: is very inoffensive*.

Tatou. *Belon obs.* 211. *Portraits*, 106.
Tatu & Tatu paba *Brazil*: Armadillo
Hispanis, Lusitanis Encuberto. *Marcgrave Brazil*. 131.
Cataphraëtus scutis duobus, cingulis sex.
Briffon quad. 25.

Dasyus sex cinctus. D. cingulis senis,
pedibus pentadactylis. *Lia. fyst.* 54.
L'Encourbert, ou Le Tatou a six bandes.
De Buffon, x. 209. *tab.* xlii. *Supplem.*
iii. 285. *tab.* lvii. *Schreber*, ii. 31. *tab.*
lxi. B. LEV. MUS.

454, SIX-BANDED.

A. with the crust of the head, shoulders, and rump, formed of angular pieces: the bands on the back six; between which, also on the neck and belly, are a few scattered hairs; tail not the length of the body, very thick at the base, tapering to a point: five toes on each foot.

Inhabits *Brazil* and *Guiana*.

PLACE.

* The authorities for the natural history: *Marcgrave*, 231. *Dampier*, ii. 61. *Gumilla Orenoque*, iii. 223 to 226. *Niuehoff*, 19. *Bancroft's Guiana*, 145. *Rechebert Antilles*, i. 286.

Ayotochtli?

455. EIGHT-
BANDED.

Ayotochtli? *Hernandez Mex.* 314.
Tatuete *Brasiliterjibus*, Verdadeiro *Lufitanis*. *Maregrave Brasil.* 231. *Chuf. exot.* 330.
Cataphractus scutis duobus cingulis octo. *Briffon quad.* 26.
Erinaceus loricatus cingulis septenis pal-

mis tetradaetylis, plantis pentadaetylis. Ameen. Acad. i. 560.
Dafypus septem cinctus. Lin. fyft. 54.
Le Tatuete. ou Tatou a huit bandes. *De Buffon*, x. 212. *Schreber*, ii. 34. 36. *tab.* lxxii. lxxvi. *fig.* 3. 4.

A. with upright ears, two inches long: small black eyes: eight bands on the fides: four toes on the fore feet, five on the hind: length, from nose to tail, about ten inches; tail nine.

PLACE.

Inhabits *Brasil*. Reckoned more delicious eating than the others.

456. NINE-
BANDED.


Armadillo. *Form. Mus.* 335.
Tatu porcinus, Schildverkel. *Klein quad.* 48.
Pig-headed Armadillo. *Grew's rarities*, 18. *Raii syn. quad.* 233.
Tatu five Armadillo Americanus. *Seb. Mus. tab.* xxix. *fig.* 1.
Dafypus novem cinctus. D. cingulis novem, palmis tetradaetylis, plantis pentadaetylis. Lin. fyft. 54. *Phil. transf.*

liv. 57. *tab.* vii.
Cataphractus scutis duobus, cingulis novem. *Briffon quad.* 27.
Le Cachichame, ou Tatou a neuf bandes. *De Buffon*, x. 215. *tab.* xxxviii. *Supplem.* iii. 287. *tab.* lviii. *Schreber*, i. 37. *tab.* lxxiv. lxxvi. *fig.* 7. 10.
American Armadillo. *Phil. Transf.* liv. 57. *tab.* vii. *Lev. Mus.*

A. with long ears: crust on the shoulders and rump marked with hexangular figures; the crust on the head marked in the same manner: nine bands on the fides, distinguished by transverse cuneiform marks: breast and belly covered with long hairs: four toes on the fore feet, five on the hind: tail long and taper: length of the whole animal three feet: the tail a little longer than the body.

In





Twelve banded Armadillo — . 1857.

In the LEVERIAN MUSEUM is a specimen of the same form, number of bands, and proportions, with this; but the crusts on the head, and other parts, are covered with large scales not angular.

Inhabits *South America*. One was brought a few years ago to *England*, from the *Mosquito* shore, and lived here some time: it was fed with raw beef, and milk, but refused our grains and fruit*.

PLACE.

Tatu five Armadillo Africanus. *Seb.* Cataphractus scutis duobus, cingulis duodecim. *Briffon quad.* 27. *Schreber*, ii. 40. *tab.* lxxv. lxxvi. *fig.* 11. 12.
Muf. i. *tab.* xxx. *fig.* 3, 4.
 Le Kabaffou, ou Tatou a douze bandes.
De Buffon, x. 218. *tab.* xl.

457. TWELVE-BANDED.

A. with broad upright ears: the crust on the shoulders marked with oblong pieces; that of the rump with hexangular: twelve bands on the sides: five toes, with very large claws, on the fore feet; five lesser on the hind: tail shorter than the body: some hairs scattered over the body.

M. de Buffon† mentions another of twelve bands, with a tail covered with rhomboid figures, which he is doubtful whether to refer to this species. It is the largest I ever heard of, being from nose to tail two feet ten inches long; the tail about one foot eight: by the figure (for I never saw the animal) it varies greatly from the other.

* This corroborates what *Marcgrave* says of one of these animals, *Cuniculus, aves mortuas aliisque devorans*; which is very extraordinary in quadrupeds which want both cutting and canine teeth.

† *P.* 256. *tab.* xli.

VOL. II.

K k

Weevil-headed

458. EIGHTEEN-
BANDED.Weeble-headed Armadillo. *Grev's rarities*, 19.Tatu Multelinus. *Raii syn. quad.* 235.
Dasypus uncinatus. D. tegmine tripartito, pedibus pentadactylis. *Lin. syst.* 53.Cataphractus scuto unico, cingulis octodecim. *Briffon quad.* 23.Le Cingaron, ou Tatou a dixhuit bandes. *De Buffon*, x. 220. *tab. xlii. Schre-ler*, ii. 42.

A. with a very slender head: small erect ears: the crust on the shoulders and rump consisting of square pieces: eighteen bands on the sides: five toes on each foot: length, from nose to tail, about fifteen inches; tail five and a half.

PLACE.

Inhabits *South America*.

DIV. II. SECT. V.

DIGITATED QUADRUPEDS:

Without Teeth.

DIV. II. SECT. V. Digitated Quadrupeds.

XXXIX. MANIS.

Back, sides, and upper part of the tail, covered with large strong scales.

Small mouth: long tongue: no teeth.

459. LONG-TAIL-
ED.

Lacertus peregrinus squamosus. *Clus. exot.* 374. *Rati syn. quad.* 274. *Pholidotus pedibus anticis et posticis tetradactylis, squamis mucronatis, cauda longissima.* *Brisson quad.* 19.
Scaly Lizard. Grew's rarities. 46. *Le Phatagin.* *De Buffon, x.* 180. *tab. xxxiv. Ass. Mus. LEV. MUS. BR. MUS.*
Manis tetradactyla. M. pedibus tetradactylis. *Lin. syst.* 53. *Schreber, ii.* 23. *tab. lxx.*


M. with a slender nose; that and the head smooth: body, legs, and tail, guarded by large sharp-pointed striated scales: the throat and belly covered with hair: short legs: four claws on each foot, one of which is very small: tail a little taper, but ends blunt. The color of the whole animal, chocolate.

PLACE.

Inhabits the islands of *India*. These animals approach so nearly the genus of *Lizards*, as to be the links in the chain of beings which connect the proper quadrupeds with the reptile class.

They grow to a great length: that which was preserved in the *Museum* of the *Royal Society*, was a yard and a half long*: from the tip of the nose to the tail, was only fourteen inches; the tail itself a yard and half a quarter.

• *Grew.**Lacertus*



(Long tailed Marmoset. V. 189.)



- Lacertus squamosus. Bontius Java, 60. Pet. Gaz. tab. xx. fig. 11.*
Armadillas squamatus major. Ceilanicus, seu Diabolus Trinovanticus dictus. Seb. Mus. i. tab. liti. lv. Klein quad. 47. Schröber, ii. 22. tab. lxix.
Pholidotus pedibus anticis et posticis pentadactylis, squamis subrotundis. Brisson quad. 18.
Manis pentadactyla. Lin. syst. 52.
Le Pangolin. De Buffon, x. 180. tab. xxxiv. Abb. Mus. LEV. MUS. BR. MUS.

460. SHORT-
TAILED.

M. with back, sides, and legs, covered with blunt scales, with bristles between each: five toes on each foot: tail not longer than the body: ears not unlike the human: chin, belly, and inside of the legs, hairy: tail broad; much shorter in proportion to the body than that of the preceding, and obtuse at the end: the color of the whole animal a pale yellow.

Inhabits the islands of *India*, and that of *Formosa*. The *Indians* call it *Pangoelling*; and the *Chinese*, *Chin Chion Seick* *. Feeds on lizards and insects: turns up the ground with its nose: walks with its claws bent under its feet: grows very fat: is esteemed very delicate eating: makes no noise, only a snorting.

PLAC.

It is also found in *Bengal*, where it is called in the *Sanskrit* language, *Vajracite*, or the *Thunderbolt reptile*, from the excessive hardness of its scales: in its stomach is found a number of small stones, probably taken in to help the digestion. In the second volume of the *Asiatic Researches*, p. 376, published under the direction of the able and learned Sir WILLIAM JONES, is a very good account of this animal; under the direction of that gentleman, a

* *Dallman in As. Stock's. 1749, 255.*

second inundation of knowledge is pouring upon the western world from its primeval seat, the *East*.

Perhaps is a native of *Guinea*: the *Quogelo* of the *Negroes*; which *Des Marchais** says grows to the length of eight feet, of which the tail is four: lives in woods and marshy places: feeds on ants, which it takes by laying its long tongue cross their paths, that member being covered with a sticky *saliva*, so the insects that attempt to pass over it cannot extricate themselves: walks very slowly: would be the prey of every ravenous beast, had it not the power of rolling itself up, and opposing to its adversary a formidable row of erected scales. In vain does the leopard attack it with its vast claws, for at last it is obliged to leave it in safety†. The *Negroes* kill these animals for the sake of the flesh, which they reckon excellent.

461. BROAD
TAILED.

A new *Manis*. Phil. Transf. vol. lx. p. 36. tab. 11.

M. with five toes on the fore feet, and four on the hind: scales of the shape of a muscle: belly quite smooth: the exterior scales end in a sharp point somewhat incurvated: tail very broad, decreasing to a point: whole length of the animal a *German* ell and five eighths: the tail half an ell and a span broad in the broadest part.

* *Voyage du des Marchais*, i. 200. *Barbot*, 114.

† Is said to destroy the *Elephant*, by twisting itself round the trunk, and compressing that tender organ with its hard scales.

This

This species was found in the wall of a merchant's house at *Tan-quebar*: when pursued it would roll itself up so that nothing but the back and tail could be seen: it was with great difficulty killed, although it was often struck with rice-stampers, or poles armed with iron: a blow on the belly deprived it of life. The scales of this genus are so hard as to strike fire.

Body

XL.
ANT-EATER.

Body covered with hair.
Small mouth: long cylindric tongue.
No teeth.

462. GREAT.

Tamandua-guacu. *Maregrave Brasl.*

225.

Tamandua-guacu five major. *Piso Brasl.*

320.

Pismire-eater, *Nituboff*, 19.

Tamandua major cauda panniculata.

Barrere France Equin. 162.Mange-fourmis. *Des Marchais*, iii. 307.Great Ant-Ear. *Rati syn. quad.* 241.

Myrmecophaga rostro longissimo, pe-

dibus anticis tetradactylis, posticis pentadactylis, cauda longissimis pilis vestita. *Briffon quad.* 15.Myrmecophaga jubata. M. palmis tetradactylis, plantis pentadactylis. *Lin. ffl.* 52. *Klein quad.* 45. tab. v.Le Tamanoir. *De Buffon*, x. 141. tab.xxix. *Suppl.* iii. 278. tab. lv. *Schreber*,ii. 14. tab. lxxvii. *Br. Mus.*

A. E. with a long slender nose: small black eyes: short round ears: slender tongue, two feet and a half long, which lies double in the mouth: legs slender: four toes on the fore feet, five on the hind: the two middle claws on the fore feet very large, strong, and hooked: the hair on the upper part of the body is half a foot long, black mixed with grey: from the neck, cross the shoulders, to the sides, is a black line bounded above with white: the fore legs are whitish, marked above the feet with a black spot: the tail is clothed with very coarse black hairs a foot long: length, from nose to tail, about three feet ten inches; the tail two and a half: weight about a hundred pounds.

PLACE AND
MANNERS.

Inhabits *Brasl* and *Guiana*: runs slowly: swims over the great rivers; at which time it flings its tail over its back: lives on ants; as soon as it discovers their nests, overturns them, or digs them

them up with its feet; then thrusts its long tongue into their retreats, and penetrating all the passages of the nest, withdraws it into its mouth loaded with prey: is fearful of rain, and protects itself against wet by covering its body with its long tail. This (as well as every species of this genus) brings but one young at a time, at which season it is dangerous to approach the place: it does not arrive at its full growth under four years. The flesh has a strong disagreeable taste, but is eaten by the *Indians*. Notwithstanding this animal wants teeth, it is fierce and dangerous; nothing that gets within its fore feet can disengage itself. The very Panthers of *America** are often unequal in the combat; for if the Ant-eater once has opportunity of embracing them, it fixes its talons in their sides, and both fall together, and both perish; for such is the obstinacy and stupidity of this animal, that it will not extricate itself even from a dead adversary †: sleeps in the day; preys by night.

The following history of this animal is given in *Dillon's Travels* through *Spain*, p. 76, in his account of the Royal Cabinet of Natural History at *Madrid*. “ The Great Ant-bear from *Buenos Ayres*, the *Myrmecophaga jubata* of *Linneus*, called by the *Spaniards* *Osa Palmera*, was alive at *Madrid* in 1776, and is now stuffed and preserved in this cabinet. The people who brought it from *Buenos Ayres* say, it differs from what *they* call the Ant-eater, which only feeds on emmets, and other insects; whereas this would eat flesh, when cut in small pieces, to the amount of four or five pounds. From the snout to the extremity of the tail, this animal is two yards in length; and his height is about two feet: the head very

* *Gumilla Orenoque*, iii. 232.

† *Piso Brasil*. 320.

narrow; the nose long and slender. The tongue is so singular, that it looks more like a worm, and extends above sixteen inches. His body is covered with long hair, of a dark brown, with white stripes on the shoulders; and when he sleeps, he covers his body with his tail."

The specimen of the Great Ant-eater in the LEVERIAN *Museum*, is superior in size to any we have before heard of.

	Feet. Inches.	
Its whole length is	—	— 7 4
Tail	—	— 2 9
From tip of the nose to the ears	—	— 1 0
Length of the hairs of the mane	—	— 1 0
————— of the tail	—	— 1 2
Height to the top of the shoulders	—	— 2 0

Both of the above are extremely rare, and in an uncommon fine state of preservation.

463 MIDDLE.

Tamandua-i. *Marcgrave Brasl.* 225.

Raii syn. quad. 242.

Tamandua minor. *Piso Brasl.* 320. *Barre France Equin.* 162.

Tamandua-guacu. *Nieubuff,* 19.

Myrmecophaga rostro longissimo, pedibus anticis tetradactylis, posticis pen-

tadactylis, cauda ferè nuda. *Briffon quad.* 16.

Myrmecophaga tetradactyla. *Lin. syst.* 52. *Zoepl. Gronov.* No. 2.

Le Tamandua. *De Buffon,* x. 144. *Schreber,* ii. 16. tab. lxviii.

A. E. with a long slender nose, bending a little down: small black mouth and eyes: small upright ears: bottoms of the fore feet round; four claws on each, like those of the former; five

five on the hind feet: hair shining and hard, of a pale yellow color: along the middle of the back, and on the hind legs, dusky: each side of the neck is a black line, that crosses the shoulders and meets at the lower end of the back: the tail is covered with longer hair than the back, is taper, and bald at the end: length, from nose to tail, one foot seven inches; the tail ten inches.

Inhabits the same country with the last: its manners much the same: when it drinks, part spurts out of the nostrils: climbs trees, and lays hold of the branches with its tail.

PLACZ.

Le Tamandua. *De Buffon*, Supplem. iii. 281. tab. lvi.

464. STRIPED.

A. E. with a taper nose, the upper mandible extending very far beyond the lower: eyes exceedingly small: ears round and short: tail covered equally with long hairs: five toes on the fore feet.

Body and tail tawny; the first marked downwards with broad stripes of black; the last annulated: legs and nose striped in the same manner: belly of a dirty white.

Length from nose to tail thirteen inches *French*; of the tail seven and a half.

M. de Buffon speaks of one, which he supposes to be the same with this; but the difference in size and colors forbid us to subscribe to his opinion. The account was transmitted to him by *M. de la Borde*, physician at *Cayenne*. The hair, says he, is whitish, and about two inches long: it has very strong talons; eats only

L 1 2

in

in the day-time; keeps in the great woods: the flesh is good: it is much more rare than the *great Ant-cater*.

Weights sixty pounds.

PLACE.

Both these inhabit *Guiana*.

465. LEST.

Tamandua minor flavescens; Ouatirouazou. *Errere France Equin* 163.

Tamandua five Coati Americana alba.

Seb. Mus. i. tab. xxxvii

Myrmecophaga rostro brevi, pedibus anticis didactylis, posticis tetractylis.

Briffon quad. 17.

Myrmecophaga didactyla. M. palmis didactylis, plantis tetractylis, cauda villosa *Lin. Syst. 51. Zooph. Gronov.*

No. 1.

Little Ant-eater. *Edw. 220.*

Le Fourmillier. *De Buffon, x. 144 tab. xxx. Schreber, ii. 17. tab. lxvi.*

A. E. with a conic nose, bending a little down: ears small, and hid in the fur: two hooked claws on the fore feet, the exterior much the largest; four on the hind feet: head, body, limbs, and upper part and sides of the tail, covered with long soft silky hair, or rather wool, of a yellowish brown color: from nose to tail seven inches and a half; tail eight and a half, the last four inches of which, on the under side, naked: the tail is thick at the base, and tapers to a point.

Inhabits *Guiana*: climbs trees, in quest of a species of ants which build their nests among the branches: has the same prehensile power with its tail as the former.

PLACE.

There is a fourth species found at the *Cape of Good Hope*, and in *Ceylon*; but being described from a mere foetus*, we shall avoid giving a transcript of Dr. *Pallas's* account of it, but wait for further information. We shall only say, that it has four toes

* *Pallas Miscel. Zool. 64.*



Lest. Ant-eater—1865.



on the fore feet, and pendulous ears, which distinguishes it from other kinds. *Kolben* * describes their manners particularly, and says they have long heads and tongues, and are toothless; and that they sometimes weigh 100lb.†: that if they fasten their claws in the ground, the strongest man cannot pull them away: that they thrust out their clammy tongue into the ants nest, and draw it into their mouth covered with insects. That the *African* species agrees with the *American* in every external particular, is confirmed; but that the last is furnished with grinding teeth, like the *Armadillo*, in the lower end of the jaws, is a discovery proved from the remarks of Doctor *Camper*, a celebrated zoologist in *Holland*. Mr. *Stracklan*, in his account of *Ceylon* ‡, gives the same account of the manners of what the natives call the *Talgai*, or Ant Bear. It is not therefore to be doubted, but that these animals are common to the old and new continents.

Fourmillier d'Afrique. *Allamand Suppl.* V. 26. tab. xi.

466. CAPE.

A. E. with a long nose, truncated at the end like that of a hog; and the nostrils resembling those of that animal: ears six inches long, thin as parchment, and covered with very fine hairs: tongue very long and slender: the hairs on the head and upper part of the body and tail very short, and so closely adhering to the skin as if they were glued to it, their color a

DESCRIP.

* *Hist. Cape*, 118; where they are called *Earth Hogs*.

† As quoted by Dr. *Peters*; I suppose from the *Dutch* edition.

‡ *Phil. Trans. abridg.* v. 180.

dirty

dirty grey; those on the sides and belly long and of a reddish hue; those on the legs still longer, black and straight: the tail thick near the base, and tapering to a point: on the fore feet are four toes; on the hind five; all armed with strong claws: those behind equal even the length of the toes: all are blunted at the end and calculated for burrowing.

SIZE. The length is three feet five to the origin of the tail, the tail one foot nine.


MANNERS. This species inhabits the neighborhood of the *Cape of Good Hope*. It lives under ground; feeds on ants like the other species; but when it has found an ants nest it looks carefully around to see whether it can feed in safety, then puts out its long tongue to catch its prey. Is an object of chase among the *Hottentots*, and is reckoned good food.

467. ACULEATED.

Porcupine Ant-eater. *Naturalist's Miscellany*, pl. 109.

A. E. Length about a foot: coated on the upper parts with spines resembling those of a porcupine, being white tipped with black; the two colors separated by a ring of tawny or dull orange: spines on the back and sides somewhat recumbent, over the tail perpendicularly erect: snout long, naked, black and tubular, opening very small: tongue lumbriciform; forehead, cheeks, and whole under parts of the body, coated with dark brown stiff hairs: legs very short, toes short, broad rounded: claws on the fore-feet, five very strong, somewhat obtuse; on the hind-feet four, of which the two first are much longer, and sharper than the others: thumb unarmed: tail very short. Inhabits *New South Wales*: preys on ants, and is found

PLACE.



. Ant-eater ——— . 1807.




found about ant-hills. A most extraordinary quadruped, connecting in some measure the two very distant genera of Porcupine, and Ant-Eater. This singular animal is more fully described by Dr. *Shaw* in the Naturalist's Miscellany, and from the figure in that work the representation here given is faithfully copied. Dr. *Shaw* is of opinion that the genera of Manis, and Myrmecophaga, ought to be either united, or else that this animal should form a distinct genus.



D I V. III.

PINNATED QUADRUPEDS:

Having fin-like feet: fore legs buried deep in the skin: hind legs pointing quite backwards.



O Rubber Seal—S. 476.

DIV. III. Pinnated Quadrupeds.

XLI. WALRUS.

With two great tusks in the upper jaw, pointing downwards.
 Four grinders on both sides, above and below.
 No cutting teeth.
 Five palmated toes on each foot.

463. ARCTIC.

Rosmarus. *Gesner Pisc.* 211. *Klein quad.* 92.

Walrus, Mors, Rosmarus. *Werm. Mus.* 289. *Raii syn. quad.* 191.

Sea-horse, or Morke. *Morten's Spitzberg*, 107, 182. *Egede Greenland*, 82.

Sea-cow. *Grantz Greenl.* i. 125. *Schreber*, ii. 88.

Odobenus. La Vache marine. *Briffon quad.* 30.

Trichecus Rosmarus. T. dentibus laniariis superioribus exsertis. *Lin. syst.* 49.

Le Morse. *De Buffon*, xiii. 358. *tab. liv.* *Br. Mus. Alb. Mus.* Lev. Mus.

W with a round head : small mouth : very thick lips, covered above and below with pellucid bristles as thick as a straw : small fiery eyes : two small orifices instead of ears : short neck : body thick in the middle, tapering towards the tail : skin thick, wrinkled, with short brownish hairs thinly dispersed : legs short : five toes on each, all connected by webs, and small nails on each : the hind feet very broad : each leg loosely articulated ; the hind legs generally extended on a line with the body : tail very short : penis long.

SIZE.

Length, from nose to tail, sometimes eighteen feet, and ten or twelve



(Arctic Walrus) — A. 1848.



twelve round in the thickest part: the teeth have been sometimes found of the weight* of 20 lb. each.

Inhabit the coast of *Spitzbergen*, *Nova Zembla*, *Hudson's Bay*, and the Gulph of *St. Laurence*, and the *Icy Sea*, as far as *Cape Tschuktschi*, and the islands off it; but does not extend southward as far as the mouth of the *Anadyr*, nor are any seen in the islands between *Kamtschatka* and *America*. Are gregarious: in some places appear in herds of hundreds: are shy animals, and avoid places which are much haunted by mankind†: are very fierce; if wounded in the water, they attempt to sink the boat, either by rising under it, or by striking their great teeth into the sides; roar very loud, and will follow the boat till it gets out of sight. Numbers of them are often seen sleeping on an island of ice; if awakened, fling themselves with great impetuosity into the sea; at which time it is dangerous to approach the ice, least they should tumble into the boat and overset it: do not go upon the land till the coast is clear of ice. At particular times, they land in amazing numbers: the moment the first gets on shore, so as to lie dry, it will not stir till another comes and forces it forward by beating it with its great teeth; this is served in the same manner by the next, and so in succession till the whole is landed, continuing tumbling over one another, and forcing the foremost, for the sake of quiet, to remove further up.

PLACE.

MANNERS.

* Teeth of this size are only found on the coast of the *Icy Sea*, where the animals are seldom molested, and have time to attain their full growth. *Hist. Kamtschatka*, 120.

† In 1608, the crew of an *English* vessel killed on *Cherry Isle* above 900 *Walrus* in seven hours time; for they lay in heaps, like hogs huddled one upon another. *Marten's Spitzberg*. 181, 182.

CHACE.

The method of killing them on the *Magdalene* isles, in the gulph of *St. Laurence*, as I am informed, is thus:—The hunters watch their landing, and as soon as they find a sufficient number for what they call a *cut*, go on shore, each armed with a spear sharp on one side like a knife, with which they cut their throats: great care must be taken not to stand in the way of those which attempt to get again to sea, which they do with great agility by tumbling headlong; for they would crush any body to death by their vast weight. They are killed for the sake of their oil, one *Walrus* producing about half a tun. The knowledge of this chace is of great antiquity; *Oëlher*, the *Norwegian*, about the year 890, made a report of it to King *Alfred*, having, as he says, made the voyage beyond *Norway*, for the more commoditie of fishing of horse-whales, which have in their teeth bones of great price and excellencie, whereof he brought some at his returne unto the King*. In fact, it was in the northern world, in early times, the substitute to ivory, being very white and very hard. Their skins, *Oëlher* says, were good to cut into cables. I do not know whether we make any use of the skin; but *M. de Buffon* says, he has seen braces for coaches made of it, which were both strong and elastic.

They bring one, or at most two, young† at a time: feed on sea herbs and fish; also on shells, which they dig out of the sand with their teeth: are said also to make use of their teeth to ascend rocks or pieces of ice, fastening them to the cracks, and drawing their bodies up by that means. Besides mankind, they seem to have no other enemy than the white Bear, with whom they have

* *Hakluyt's coll. Voy.* i. 5.† *Barentz voy.* 4.

terrible

terrible combats; but generally come off victorious, by means of their great teeth.

Le Dugon. *De Buffon*, xiii. 374. *tab. lvi. Schreber*, ii. 93.

469. INDIAN.

W. with two short canine teeth, or tusks, placed in the upper jaw pretty close to each other: in the upper jaw four grinders on each side, placed at a distance from the tusks; in the lower, three on each side.

Inhabits the *Cape of Good Hope* and the *Philippine* isles. The head described above being supposed to belong to an animal resembling a *Walrus*, found in the seas of *Africa* and *India*, as appears from some citations from travellers, too unsatisfactory to merit repetition. It is said by one, that it goes upon land to feed on the green moss; and that it is called in the *Philippines*, the *Dugung* *.

PLACE.

* *De Buffon*, xiii. 377. *the note*.

Cutting

XLII. SEAL.

Cutting teeth, and two canine teeth in each jaw.

Five palmated toes on each foot.

Body thick at the shoulders, tapering towards the tail.

470. COMMON.

Φωκη. *Arist. hist. An. lib. vi. c. 12. Op-
pian Halieut. v. 376.*

Vitulus Oceani. Rondeletii, 453. 458.

*Le Veau Marin, ou Loup de Mer. Belon
Poissons, 25.*

*Phoca. Gesner Pisc. 830. Worm. Mus. 289.
Klein quad. 93. Brisson quad. 162.*

*Seal, Seole, or Sea Calf; Phoca five
Vitulus Marinus. Raii syn. quad. 189.*

Phil. trans. abridg. vol. xlvii. 120. tab.

vi. fig. 3.

Kassigiak. Crantz hist. Greenl. i. 123.

*Phoca vitulina. Ph. capite laevi inauri-
culato. Lin. syst. 56.*

Sial. Faun. succ. N° 4.

*Le Phoque. De Buffon, xiii. 333. tab.
xlv. Schreber, cxxxiv.*

*Seal. Br. Zool. i. 71, Br. Zool. illustr.
xlviii. LEV. Mus.*

S with large black eyes: large whiskers: oblong nostrils: flat head and nose: tongue forked at the end: two canine teeth in each jaw: six cutting teeth in the upper jaw; four in the lower: no external ears: body covered with thick short hair: short tail: toes furnished with strong sharp claws: usual length from five to six feet: color very various, dusky, brinded, or spotted with white or yellow.

PLACE.

Inhabit most quarters of the globe, but in greatest multitudes towards the North and the South; swarm near the *Arctic* circle, and the lower parts of *South America**, in both oceans; near the

* *Dampier* says, that they are seen by thousands on the isle of *Juan Fernandez*; that the young bleat like lambs; that none are found in the *South Sea*, north of the equator, till lat. 21; that he never saw any in the *West Indies*, except in the *Bay of Campeachy*; nor yet in the *East Indies*. i. 88, 89.

southern

southern end of *Terra del Fuego*; and even among the floating ice as low as *south lat.* 60. 21 *. Found in the *Caspian* † Sea, in the lake *Aral*, and lakes ‡ *Baikal* and *Oron*, which are fresh waters. They are lesser than those which frequent salt waters; but so fat that they seem almost shapeless. In lake *Baikal* some are covered with silvery hairs; others are yellowish, and have a large dark-colored mark on the hind part of the back, covering almost a third of the body.

They are found in the *Caspian* sea, in most amazing multitudes: they vary infinitely in their colors: some are wholly white; others wholly black; others of a yellowish white; others mouse colored; and others again spotted like a leopard: they creep out of the sea on the shores, and are killed as fast as they come; and are followed by a vast succession of others, who undergo the same fate. It is singular that the seals of the *Caspian* are very tenacious of life; it is well known that the smallest blow on the nose kills those of *Europe*. At approach of winter they go up the *Jaik*, and are killed in great numbers on the ice: they are sought for the skins and the oil: numbers are destroyed by the wolves and jackals; for which reason the seal-hunters watch most carefully the haunts of the seals in order to drive away their enemies. The seasons for hunting the seals are spring and autumn ||.

Seals bring two young at a time, which for some short space are white and woolly; bring forth in *autumn*, and suckle their young in caverns, or in rocks, till they are six or seven weeks old,

* *Cook's voy.* i. 34.

† *Bell's travels*, i. 49.

‡ The same, 280.

|| *Decouvertes, &c. faites par les Russes.* ii. 36. 4to. ed.

when they take to sea: cannot continue long under water; are therefore very frequently obliged to rise to take breath, and often float on the waves. In summer, sleep on rocks, or on sand-banks: if surprized, precipitate into the sea; or if at any distance, scramble along, and fling up the sand and gravel with great force with their hind feet, making a piteous moaning: if overtaken, will make a vigorous defence with their feet and teeth: a slight blow on the nose kills them, otherwise they will bear numbers of wounds. I imagine that the *Caspian* seal-hunters are not acquainted with the method.

Swim with vast strength and swiftness; frolic greatly in their element, and will sport without fear about ships and boats; which may have given rise to the fable of *Sea-nymphs* and *Sirens*. Their docility is very great, and their nature gentle: there is an instance of one which was so far tamed as to answer to the call of its keeper, crawl out of its tub at command, stretch at full length, and return into the water when directed; and extend its neck to kiss its master as often and as long as required*.

They never go any great distance from land: feed on all sorts of fish: are themselves good food, and often eaten by voyagers: killed for the sake of the oil made from their fat; a young seal will yield eight gallons: their skins very useful in making waist-coats, covers for trunks, and other conveniences: those of the lake *Baikal* are sold to the *Chinese*, who dye, and sell them to the *Mongals*† to face their fur-coats: are the wealth of the *Greenlanders*, supplying them with every necessary of life.

* Dr. *Parsons* in *Ph. transf.* xlvii. 113.

† *Müller's Ruff. Samlung.* iii. 559.





Seal, head ———, l. 474.

Br. Zool. i. p. 122.

Le Phoque à ventre blanc. *De Buffon*, Supplém. vi. 310. tab. xlv.

471. PIED.

S. with the nose taper and elongated: fore feet furnished with five toes, inclosed in a membrane, but very distinct; the claws long and strait: the hind feet very broad; five distinct toes, with the claws just extending to the margin of the membrane, which expands into the form of a crescent.

This I saw at *Chester*; it was taken near that city in *May* 1766. On the first capture its skin was naked, like that of a porpoise; and only the head, and a small spot beneath each leg, was hairy. Before it died the hair began to grow on other parts: the fore part of the head was black, hind part of the head and the throat white; beneath each fore leg a spot of the same color; hind feet of a dirty white; the rest of the animal of an intense black. I believe they vary in the disposition of the colors: that given by *M. de Buffon* had only the belly white. These species, according to that great writer, frequent the coast of the *Adriatic*: the length of that described by *M. de Buffon* was seven feet and a half; that which I saw was very much less, and probably a young one.

PLACE.

Vitulus Maris Mediterranei. Rondel.
Phoca Monachus, capite inauriculato,
dentibus incis: utriusque maxillæ qua-

tuor, palmis indivisis plantis exungui-
culatis. *Herman.*

472. MEDITER-
RANEAN.

S. with a small head: neck longer than that of the common
seal: orifices of the ears not larger than a pea: hair short
VOL. II. N n and

and rude: color dusky, spotted with ash-color: above the navel, of the specimen described by Mr. *Herman*, was a tawny spot: the toes on the fore feet furnished with nails: the hind feet pinniform, and without nails.

When the animal is placed on its back, the skin of the neck folds like a monk's hood.

SIZE. Length of the specimen described by Mr. *Herman* was eight feet seven inches: the greatest circumference above five feet.

PLACE. Inhabits the *Mediterranean Sea*, and as yet not discovered in the ocean. The common, or oceanic species, is probably an inhabitant of the same sea, for the species described by *Aristotle**, is of that kind; he minutely describes the feet, and attributes to the hind, as well as the fore feet, five toes, every one furnished with nails: that species therefore is the *Phoca* of the ancients, not the kind just under consideration.

473. LONG-
NECKED.

Long-necked Seal. *Grew's Museum*, 95.

S. with a slender body: length from the nose to the fore legs as great as from the fore legs to the tail: no claws on the fore feet, which resemble fins.

This was preserved in the *Museum* of the *Royal Society*.

Doctor *Parsons* has given a figure of it in the xlviiiith vol. of *Ph. Tr.* tab. vi. but we are left uninformed of its place.

* *Hist. an. lib.* ii. c. i.

ALLIED to this is another SEAL in the same *Museum*, sent of late years from the *Falkland isles*: its length is four feet; hair short, cinereous tipped with dirty white.

Nose short, beset with strong black bristles: short, narrow, pointed auricles.

Upper cutting teeth fulcated transversely; the lower in an opposite direction: on each side of the canine teeth, a lesser, or secondary one: grinders conoid, with a small process on one side near the base.

No claws on the fore feet; but beneath the skin evident marks of the bones of five toes: the skin extends far beyond their ends. On the toes of the hind legs are four long and strait claws; but the skin stretches far beyond, which gives them a very pinniform look.

This species probably inhabits also the seas about *Juan Fernandez*; for *Don Ulloa** informs us of one kind, which is not above a yard long. The small Seals inhabit from the *Falkland Islands*, round *Cape Horn*, even as far as *New Zealand*; and are seen further from shore than any other kind. They are very sportive, dipping up and down like porpoises, and go on in a progressive course like those fish. When they sleep, one fin generally appears above the water. They perhaps extend as far as the *Society Islands*, at least the natives have a name for the Seal, which they call *Himi*.

474. FALKLAND
ISLE.

PLACE.

* *Ulloa* says, the first species of Seal found near that isle, is not above a yard long. ii. 226.

475. TORTOISE-
HEADED.Tortoise-headed Seal. *Ph. Transf.* xlvii. 120. *tab.* vi.

S. with a head like that of a tortoise : neck slenderer than head or body : feet like those of the common Seal.

We are indebted to Doctor *Parsons* for the account of this species, who says it is found on the shores of many parts of *Europe*.

476. RUBBON.

S. with very short fine glossy bristly hair, of an uniform color, almost black ; marked along the sides, and towards the head and tail, with a stripe of a pale yellow color, exactly resembling a rubbon laid on it by art ; words cannot sufficiently convey the idea, the form is therefore engraven on the title of Division III. *Pinnated Quadrupeds*, from a drawing communicated to me by Doctor *Pallas*, who received it from one of the remotest *Kuril* islands.

Its size is unknown, for Doctor *Pallas* received only the middle part, which had been cut out of a very large skin, so that no description can be given of head, feet or tail : a. shews the part supposed to be next to the head ; b. that to the tail.

OBSCURE SPECIES.

Other obscure species in those seas, which are mentioned in *Steller's MSS.* are, I. A middle-sized Seal, elegantly speckled in all parts : II. One with brown spots, scarcer than the rest : III. A black species with a peculiar conformation of the hind legs.

Phoca Leporina. Lepechin. aſt. acad. Petrop. pars i. 264. tab. viii. ix.

477. LEPORINE.

S. with fur, ſoft as that of a hare, upright and interwoven; of a dirty white color: whiſkers long and thick, ſo that the animal appears bearded: head long: upper lip thick: four cutting teeth above; the ſame below: nails on fore and hind feet.

Uſual length ſix feet and a half; greateſt circumference five feet two.

SIZE.

Inhabits the White ſea during ſummer; aſcends and deſcends the rivers in queſt of prey; found alſo off *Iceland*, and from *Spitzbergen* to the *Tchutkinofs*.

PLACE.

Sea Calf. *Phil. Tranſ. ix. 74. tab. v.*
Le grand Phoque. *De Buffon, xiii. 345.*

Uſuk? *Crantz Greenl. i. 125. Schreber*
Cab. i. 43. LEV. MUS.

478. GREAT.

S. reſembling the common, but grow to the length of twelve*. feet: that deſcribed in the *Phil. Tranſ.* was ſeven feet and a half long, yet ſo young as to have ſcarce any teeth; the common Seal is at full growth when it has attained the length of ſix.

Inhabits the coaſt of *Scotland*, and the ſouth of *Greenland*. The ſkin is thick, and is uſed by the *Greenlanders* to cut thongs out of for their Seal fiſhery. Perhaps is the ſame with the great *Kamtſchatkan* Seal, called by the *Ruſſians*, *Lachtach*, weighing 800lb.†, whoſe cubs are black.

PLACE.

* A gentleman of my acquaintance ſhot one of that ſize in the north of *Scotland*.

† *Muller's Voy. Kamtſchatka, Co.*

Neitſek.

479. ROUGH.

Neitick. *Grantz Greenland*. i. 124. *Schreber*, clxxxvi.

S. with rough bristly hair, intermixed like that of a hog; of a pale brown color.

PLACE.

Inhabits *Greenland*: the natives make garments of its skin, turning the hairy side inmost. Perhaps what our *Newfoundland* Seal-hunters call *Square Phipper*; whose coat, they say, is like that of a water-dog, and weighs sometimes 500 lb.

480. PORCINE.

Phoca porcina. *Molina Chili*. 260.

S. agreeing in general form with the *Ursine*, N° 485, but the nose is longer, and resembles a hog's snout; it has also the vestiges of ears: the feet have five distinct toes, covered with a common membrane.


PLACE.

Inhabits the coast of *Chili*, but is a rare species.

481. EARED.

S. with conoid head: nose rather pointed: ears an inch long, very narrow and pointed: whiskers very long and white: fore feet pinniform; neither toes nor nails apparent, terminated membranously: in the hind feet the toes apparent, and each furnished with its nail; the membrane extends beyond, and then divides into five narrow divisions, correspondent to each toe: the tail a little





Mazell Sculp.

Harp Seal. . 1. 1833.

little more than an inch long: the whole body is covered with longish hair of a whitish or creme-color: the length from nose to tail is rather more than two feet.

Inhabits the freights of *Magellan*. This species is finely preserved in Mr. *Parkinson's* Museum, on the southward side of *Black-friars Bridge*. That gentleman has very properly placed seizing on it the *Condor* vulture, the vast co-inhabitant of the *Magellanic* regions. Every one knows that Mr. *Parkinson* is now possessed of the late Sir *Ashton Lever's* Museum; I have therefore still retained the words *LEV. MUS.* to the description of every animal contained in that matchless collection.

PLACE.

Clap-myss. *Egede Greenl.* 84.Neiterfoak. *Crantz Greenl.* i. 124.

482. HOODED.

S. with a strong folded skin on the forehead, which it can fling over its eyes and nose, to defend them against stones and sand in stormy weather: its hair white, with a thick coat of thick black wool under, which makes it appear of a fine grey.

Inhabits only the south of *Greenland*, and *Newfoundland*: in the last is called the *Hooded Seal*: the hunters say they cannot kill it till they remove the integument on the head.

PLACE.

Black-sided Seal. *Egede Greenl.* plate iii.
Attarfoak. *Crantz Greenl.* i. 124. *Schre-
ber, Cab.* i. 39.

Phoca oceanica. *Krylatca* Ruff. *Lefechin-
ast. acad. Petrop.* pars i. 259. tab. vi.
vii.

483. HARP.

S. with a pointed head and thick body, of a whitish grey color, marked on the sides with two black crescents, the horns pointing

pointing upwards towards each other; does not attain this mark till the fifth year; till that period, changes its color annually, and is distinguished by the *Greenlanders* by different names each year.

PLACE.

Inhabits *Greenland* and *Newfoundland*, *Iceland*, the *White Sea*, and *Frozen Ocean*, and passes through the *Asiatic* strait, as low as *Kamschatka*: is the most valuable kind; the skin the thickest and best, and its produce of oil the greatest: grows to the length of nine feet. Our Fishers call this the *Harp*, or *Heart Seal*, and style the marks on the sides the saddle. There is a blackish variety, which they say is a young *Harp*, called *Bedlemer*.

484. LITTLE.

Le petit Phoque. *De Buffon*, iii. 341. tab. liii. *Schreber*, cxxxv:
LEV. MUS.

SIZE.

S. with the four middle cutting teeth of the upper jaw bifurcated; the two middle of the lower jaw slightly trifurcated: a rudiment of an ear: the webs of the feet extending far beyond the toes and nails: hair soft, smooth, and longer than in the common Seal: color dusky on the head and back; beneath brownish: length two feet four inches.

Our Seal-hunters affirm, that they often observe, on the coast of *Newfoundland*, a small species, not exceeding two feet, or two feet and a half, in length. *M. de Buffon* says the specimen in the cabinet of the *French* king came from *India*; but from the authority of *Dampier*, and of modern voyagers to the *East Indies*,





Musque Seal — . 185.

who have assured me they never saw any Seals * there, I suspect he was imposed on.

Captain *Abraham Dixon* assured me that he saw off the coast of *North America*, in his voyages of 1785 to 1788, multitudes of small Seals, not exceeding a foot in length: they were perpetually dipping and rising again, but were so active that he never could procure a specimen.

Ursus Marinus. Steller. Nov. Com. Pe- *Phoca Ursina. Ph. capite auriculato. Lin.*
trop. ii. 331. tab. xv. *syn. 55.*
Sea Cat. Hist. Kamtschatka, 123. Mul- *L'Ours Marin. Brisson quad. 166. Schre-*
ler's Exped. 59. *ber, cxxxii.*

485. URSINE.

THERE are three marine animals, which keep a particular situation, and seem divided between the N. E. of *Asia*, and N. W. of *America*, in the narrow seas between those vast continents. These are what are called the *Sea Lion* and *Sea Bear*, and the *Manati*. They inhabit, from *June* to *September*, the isles that are scattered in the seas between *Kamtschatka* and *America*, in order to copulate, and bring forth their young in full security. They never land upon *Kamtschatka*. The accurate and indefatigable naturalist *Steller* was the first who gave an exact description of them; he and his companions, in the *Russian* expedition of 1742, were in all probability the first *Europeans* who gave

PLACE.

* A gentleman, the most curious, and greatest navigator of the *Indian* seas now living, informed me, that he not only never met with any Seals in those seas, but even none nearer than the isles of *Galapagos*, a little north of the line, on the coast of *America*.

VOL. II.

O o

them

them any disturbance in those their retreats. In *September*, these animals quit their stations, vastly emaciated; some return to the *Asiatic*, others to the *American* shores; but, like the Sea Otters, are confined in those seas between lat. 50 and 56.

They are not, as far as I can discover, found from those places, any where nearer than *New Zealand**, where they are very common, and again about *Staten Land*†, the frozen island of *New Georgia*‡, and the *Falkland* islands§. I suspect that they are also found in the island of *Juan Fernandez*; for, among the Seals so imperfectly described by Don *Ulloa*§, his second kind seems to be of this species. I may add, that *Alexander Selkirk* speaks of Seals which come on shore in that island in *November* to whelp†, which nearly corresponds with the time our late circumnavigators saw them in *New Year's islands*, where they found them and their young in *December*. Lastly, I may mention the isles of *Gallopagos*, where Captain *Woodes Rogers* says he was attacked by a fierce Seal, as big as a bear, and with difficulty escaped with his life**.

The *Ursine* Seal, a name we substitute for the sea-bear, leads, during the three months in summer, a most indolent life: it arrives at the islands vastly fat; but during that time they are scarce ever in motion: confine themselves for whole weeks to one spot, sleep a great part of the time, eat nothing, and, except the employment the females have in suckling their young, are totally inactive. They live in families; each male has from eight to fifty females, whom he guards with the jealousy of an eastern monarch, and though they lie by thousands on the shores, each family

* *Forster's obs.* 189. † *Cook's voy.* ii. 203. ‡ *Cook's voy.* ii. 213.
Forster's voy. ii. 529. § *Pernetti, Engl. ed.* 187. tab. xvi. § *Voy.* ii. 226.
 † In *Woodes Rogers's voy.* 136. ** The same, 265.

keeps itself separate from the rest, and sometimes, with the young and unmarried ones, amount to a hundred and twenty. The old animals, which are destitute of females, or deserted by them, live apart, and are excessively splenetic, peevish, and quarrelsome: are excessively fierce, and so attached to their old haunts, that they would die sooner than quit them. They are monstrously fat, and have a most hircine smell. If another approaches their station, they are roused from their indolence, and instantly snap at it, and a battle ensues; in the conflict, they perhaps intrude on the seat of another: this gives new cause of offence, so in the end the discord becomes universal, and is spread thro' the whole shore.

The other males are also very irascible: the causes of their disputes are generally these:—The first and the most terrible is, when an attempt is made by another to seduce one of their mistresses, or a young female of the family. This insult produces a combat, and the conqueror is immediately followed by the whole seraglio, who are sure of deserting the unhappy vanquished. The second reason of a quarrel is, when one invades the seat of another. The third arises from their interfering in the disputes of others. These battles are very violent; the wounds they receive are very deep, and resemble the cuts of a sabre. At the end of a fight they fling themselves into the sea, to wash away the blood.

The males are very fond of their young; but very tyrannical towards the females: if any body attempts to take their cub, the male stands on the defensive, while the female makes off with the young in her mouth; should she drop it, the former instantly quits his enemy, falls on her, and beats her against the stones,

till he leaves her for dead. As soon as she recovers, she comes in the most suppliant manner to the male, crawls to his feet, and washes them with her tears: he, in the mean time, stalks about in the most insulting manner; but in case the young one is carried off, he melts into the deepest affliction, and shews all signs of deep concern. It is probable that he feels his misfortune the more sensibly, as the female generally brings but one at a time; never more than two. Even the cubs of those on the island of *New Georgia** are very fierce, barking at our sailors as they passed by, and biting at their legs. The breeding-time in this island is in the beginning of *January*.

They swim very swiftly, at the rate of seven miles an hour. If wounded, will seize on the boat, and carry it along with vast impetuosity, and oftentimes sink it. They can continue a long time under water. When they want to climb the rocks, they fasten with the fore paws, and so draw themselves up. They are very tenacious of life, and will live for a fortnight after receiving such wounds as would immediately destroy any other animal.

DESCRIPTION.

The male of this species is vastly superior in size to the female. The bodies of each are of a conic form, very thick before, and taper to the tail. The length of a large one is eight feet; the greatest circumference five feet; near the tail, twenty inches. The weight 800 lb. The nose projects like that of a pug dog, but the head rises suddenly: nostrils oval, and divided by a *septum*: the lips thick; their inside red and serrated; whiskers long and white.

The teeth lock into each other when the mouth is closed. In

* *Forsker's voy.* ii. 516, 529.

the upper jaw are four cutting teeth, each bifurcated; on both sides is a small sharp canine tooth bending inwards; near that another, larger: the grinders resemble canine teeth, and are six in number in each jaw: in the lower jaw are also four cutting teeth and two canine: but only four grinders in each jaw: in all, thirty-six teeth.

Tongue bifid: eyes large and prominent: iris black: pupil sinuagline: the eyes may be covered at pleasure with a fleshy membrane: the ears are small, sharp-pointed; hairy without, smooth and polished within.

The length of the fore-legs is twenty-four inches; like those of other quadrupeds, not immersed in the body like those of Seals: the feet are formed with toes, as those of other animals, but are covered with a naked skin, so that externally they seem a shapeless mass, and have only the rudiments of nails to five latent toes: the hind legs are twenty-two inches long, are fixed to the body quite behind, like those of Seals, but are capable of being brought forward, so that the animal makes use of them to scratch its head: these feet are divided into five toes, each divided by a great web, and are a foot broad: the tail is only two inches long.

The hair is long and rough; beneath which is a soft down, of a bay-color: on the neck of the old males the hair is erect, and a little longer than the rest. The general color of these animals is black, but the hairs of the old ones are tipped with grey. The females are cinereous. The skins of the young, cut out of the bellies of their dams, are very useful for cloathing, and cost about 3s. 4d. each; the skin of an old one, 4s.

The fat and flesh of the old males is very nauseous; but the
flesh

flesh of the females resembles lamb; and the young ones roasted are as good as sucking-pigs.

486. BOTTLE-
NOSE.

Sea Lion. *Dampier's voy.* i. 90. iv. 15. *Le Lion Marin. Brisson quad.* 167. *De Rogers's voy.* 136. *Ansen's voy.* 122. *Buffon*, xiii. 351. *Schreber*, cxxxiii. *Phoca Leonina. Ph. capite anticè cristato. Lin. syst.* 55. *Le Lame. Phoca elephantina. Molina Chili.* 261.

S. (*the male*) with a projecting snout, hanging five or six inches below the lower jaw: the upper part consists of a loose wrinkled skin, which the animal, when angry, has the power of blowing up, so as to give the nose an hooked or arched appearance: the feet short and dusky; five toes on each, furnished with nails: the hind feet have the appearance of great lacinated fins: large eyes: great whiskers: hair on the body short, and of a dun color; that on the neck a little longer: the skin very thick. Length of an old male twenty feet; greatest circumference, fifteen.

SIZE.

Female. Nose blunt, tuberos at the top: nostrils wide: mouth breaking very little into the jaws; two small cutting teeth below, two small and two larger above; two canine teeth, remote from the preceding; five grinders in each jaw; all the teeth conic: eyes oblique and small: auricles none: fore legs twenty inches long: toes furnished with flat oblong nails: hind parts, instead of legs, divided into two great bifurcated fins: no tail: the whole covered with short rust-colored hair. Length, from nose to the end of the fins, four yards: greatest circumference two yards and a half*.

SIZE.

* Described from a well-preserved specimen in the *Museum* of the **ROYAL SOCIETY**. This is the animal called by *Dr. Parsons*, a *Manati*.

Inhabits

Inhabits the seas about *New Zealand* *, the island of *Juan Fernandez* † and the *Falkland islands* ‡, and that of *New Georgia* §, S. lat. 54—40. Are seen in great numbers, in *June* and *July*, the breeding-season, on the island of *Juan Fernandez*, which they resort to for the purpose of suckling their young on shore, and continue there till *September*. They bring two at a time. The female, during that season, is very fierce: one of Lord *Anson's* sailors was killed by the enraged dam of a whelp, which he had robbed her of. The male shews little attachment to its young, but the female is excessively fond of it: the former will suffer it to be killed before his face without shewing any resentment. Towards evening, both male and female swim a little way to sea, the last with the young on its back, which the male will push off, as if to teach it to swim.

They arrive on the breeding-islands very fat and full of blood: when they are in motion, they seem like a great skin full of oil, from the tremulous movement of the blubber, which has been found to be a foot thick. The *Spaniards* very properly call these, and the *Urigne lobos de Aceyte*, or *oil wolves*, from their looking like a skin full of oil, from the motion of the vast quantity of fat or blubber, of which their bodies consist §. One has been known to yield a butt of oil; and so full of blood, that what has run out of a single animal † has filled two hogheads. The flesh is eatable: Lord *Anson's* people eat it under the denomination of beef, to distinguish it from that of Seal, which they called lamb.

* *Forster's obs.* 190.† *Anson's voy.* 122.‡ *Pernetti* 202.§ *Cook's voy.* ii. 213. *Forster's voy.* 527.§ *Ulloa's voy.* ii. 227.† *Anson's voy.* 123.

The old animals have a tremendous appearance, yet are excessively timid, except at the breeding-season, when they seem to lose their apprehensions, and are less disturbed at the sight of man. At other times, they hurry into the water; or, if awakened out of their sleep by a loud noise, or blows, fall into vast confusion, tumble down, and tremble in every part, thro' fear.

These animals associate in families, like the former, but not in such great numbers: the males shew equal jealousy about their mistresses, and have bloody combats on their accounts: oft-times there is one of superior courage to the rest, and procures by dint of valour a greater number of females than others. They are of a very lethargic nature, fond of wallowing in miry places, and will lie like swine on one another: they grunt like those animals, and will sometimes snort like horses in full vigor. They are very inactive on land: to prevent surprize, each herd places a centinel, who gives certain signals at appearance of danger: during the breeding-season, they abstain from food, and before that is elapsed become very lean; at other times they feed on fish and the smaller Seals.

487. LEONINE.

Bestia Marina, Kurillis, Kamtschadalis et
Ruffis, Kurilico nomine Siwutscha dicta.
Nov. Com. Petrop. ii. 360.
Phora Leonina. Melina Chili. 262.

Sea Lion. *Cook's voy. ii. 203. Ferster's*
voy. ii. 513. Pernetty's voy. 240. tab.
xvi.

S. with a short nose turning a little up: great head: eyes large: whiskers long and thick, and strong enough to serve for pick-tooths: on the neck and shoulders of the MALE is a great mane



Lionine Seal — . N^o 487.



mane of coarse, long, waving hair, not unlike the shaggy appearance of a lion: the rest of the body covered with a very short, smooth, and glossy coat. The whole color is a deep brown: those of the *Kamtschatkan* islands are reddish; the females tawny.

The fore feet are like those of the *Urşine Seal*, resembling a flat fin, formed of a black coriaceous substance, without the least external appearance of toes, as most erroneously represented by *Pernetti*: the hind feet are very broad, furnished with very small nails, with a narrow stripe of membrane extending far beyond each: tail very short: hind parts vastly large, swelling out with the vast quantity of fat.

The old males are from ten to fourteen feet long, and of great circumference about the shoulders; they weigh from twelve to fifteen hundred pounds: the females are from six to eight feet in length, of a more slender form than the males, and are quite smooth.

SIZE.

Penrose and *Pernetti* ascribe a much greater size to those of the *Falkland* isles. The former says, that some of the males are twenty-six feet long*; and the latter affirms that their length is twenty-five feet, and their girth round the shoulders from nineteen to twenty†.

They inhabit in vast numbers *Penguin* and *Seal* islands, near *Cape Desire*, on the coast of *Patagonia*‡; are found within the straits of *Magellan*, and on *Falkland isles*: they have not yet been discovered in any other part of the southern hemisphere, or in any other place nearer than the sea between *Kamtschatka* and *America*. The inhabitants of *Chili* call them *Thapel lame*, or the Seal with a mane.

PLACE.

* *Exped. Falkland Isles*, 28.

† *Voy. Malouines*, 240.

‡ *Narborough*, 31.

MANNERS.

They live in families separate from the *Ursine* and other Seals: these possess the beach nearest to the sea: they have much of the lethargic nature of the former; and, like them, are polygamous: they have from two to thirty females apiece: they have a fierce look; the old ones snort and roar like enraged bulls; but on the approach of mankind, fly with great precipitation: the females make a noise like calves: the young bleat like lambs.

The old males lie apart, and possess some large stone, which no other dare approach; if they do, a dreadful combat ensues, and the marks of their rage appear in the deep gashes on various parts of their bodies. The males frequently go into the water, take a large circuit, land, and caress their females with great affection; put snout to snout as if they were kissing one another. The females, on seeing their male destroyed, will sometimes attempt to carry away a cub in their mouth, but oftener desert them through fear.

The food of these animals is the lesser Seals, Pinguins, and fish; but while they are ashore they keep, in the breeding-time, a fast of three or four months; but to keep their stomachs distended, will swallow a number of large stones, each as big as two fists.

S. with the body very thick at the shoulders, gradually lessening to the hind legs: head like a dog, with the ears close cut: nose short and blunt: upper lip cunilineated: six cutting teeth above; four below: fore foot has four toes inclosed in a membranous sheath, so as to resemble fins: the hind feet are hid in a
continuation.

continuation of the skin of the back, and have five toes of unequal length, like those of the human hand: tail three inches long: the skin is covered with two sorts of hairs, one like that of an ox, the other more hard: the colors various: length from three to eight feet.

These are the *Sea wolves* which navigators speak of off the island of *Lobos*, near the river *Plata*. They appear in vast multitudes, meet the ships, and will even hang by the sides with their paws, and seem to stare at and admire the crew: then drop off and return to their haunts*. They swim with incredible swiftness. The natives of *Chili* kill them for the skins, and for the oil.

PLACES:

* Father *Cattaneo's* first Letter in the missions of *Paraguay*. p. 227.

XLIII.
MANATI.

Pinniform fore-legs: hind parts ending in a tail, horizontally flat. Two teats between the legs.

489. WHALE-
TAILED.

Manati. *Ruforum* Morſkuia Korowa. *De Buffon, Supplem.* vi. 399.
Steller in Nov. Com. Petrop. ii. 294. *Tricheus Borealis. Gm. Lin.* i. 61. p.
Schreter, ii. 95. *Hiſt. Kamſchatka,* 132.

THIS animal in nature ſo nearly approaches the cetaceous tribe, that it is merely in conformity to the ſyſtematic writers, that I continue it in this claſs: it ſcarce deſerves the name of a biped; what are called feet are little more than pectoral fins; they ſerve only for ſwimming; they are never uſed to aſſiſt the animal in walking, or landing; for it never goes aſhore, nor ever attempts to climb the rocks, like the *Walrus* and *Seal*. It brings forth in the water, and, like the whale, ſuckles its young in that element: like the whale, it has no voice; and, like that animal, has an horizontal broad tail in form of a crescent, without even the rudiments of hind feet.

PLACE.

Inhabits the ſeas about *Bering's* and the other *Aleutian* iſlands, which intervene between *Kamſchatka* and *America*, but never appears off *Kamſchatka*, unleſs blown aſhore by a tempeſt. Is probably the ſame ſpecies which is found above *Mindanao* *; but is certainly that which inhabits near *Rodriguez*, vulgarly called *Diego Reys*, an iſland to the eaſt of *Mauritius*, or the iſle of

* *Dampier's voy.* i. 321.

France, near which it is likewise found *: Sir *Joseph Banks* favored me with the sketch of one drawn off this island in 1761, by *Uliuke Mole*, of the *Norfolk* man of war. It is likely that this species extends to *New Holland*, where *Dampier* says he has seen it †.

They live perpetually in the water, and frequent the edges of the shores; and in calm weather swim in great droves near the mouths of rivers; in the time of flood they come so near the land that a person may stroke them with his hand: if hurt, they swim out to the sea: but presently return again. They live in families, one near another; each consists of a male, a female, a half-grown young one, and a very small one. The females oblige the young to swim before them, while the other old ones surround, and, as it were, guard them on all sides. The affection between the male and female is very great; for if she is attacked, he will defend her to the utmost, and if she is killed, will follow her corpse to the very shore, and swim for some days near the place it has been landed at.

MANNERS.

They copulate in the spring, in the same manner as the human kind, especially in calm weather, towards the evening. The female swims gently about; the male pursues, till, tired with wantoning, she flings herself on her back, and admits his embraces ‡. *Steller* thinks they go with young above a year: it is certain that they bring but one young at a time, which they suckle by two teats placed between the breast.

They are vastly voracious and gluttonous, and feed not only on the *fuci* that grow in the sea, but such as are flung on the

* *Voy. de la Caille*, 229.† *Voy.* i. 33.‡ The *Leonine* and *Ursine* Seals copulate in the same manner, only, after sporting in the sea for some time, they come on shore for that purpose.

edges

edges of the shore. When they are filled, they fall asleep on their backs. During their meals, they are so intent on their food, that any one may go among them and chuse which he likes best.

Their back and their sides are generally above water; and as their skin is filled with a species of louse peculiar to them, numbers of gulls are continually perching on their backs and picking out the insects.

They continue in the *Kamtschatkan* and *American* seas the whole year; but in winter are very lean, so that you may count their ribs. They are taken by harpoons fastened to a strong cord, and after they are struck it requires the united force of thirty men to draw them on shore. Sometimes, when they are transfixed, they will lay hold of the rocks with their paws, and stick so fast as to leave the skin behind before they can be forced off. When a *Manati* is struck, its companions swim to its assistance; some will attempt to overturn the boat, by getting under it; others will press down the rope, in order to break it; and others will strike at the harpoon with their tails, with a view of getting it out, which they often succeed in. They have not any voice, but make a noise by hard breathing, like the snorting of a horse.

DESCRIPTION.

They are of an enormous size: some are 28 feet long, and 8000 lb. in weight; but if the *Mindanao* species is the same with this, it decreases greatly in size as it advances southward, for the largest which *Dampier* saw there, weighed only six hundred pounds*. The head, in proportion to the bulk of the animal,

* *Dampier*, i. 321. Voyagers are requested to observe, whether there are not the two species about this and the other islands of the *Indian* ocean.

is small, oblong and almost square: the nostrils are filled with short bristles: the gape, or *rietus*, is small: the lips are double: near the junction of the two jaws the mouth is full of white tubular bristles, which serve the same use as the laminæ in whales, to prevent the food running out with the water: the lips are also full of bristles, which serve instead of teeth to cut the strong roots of the sea-plants, which floating ashore are a sign of the vicinity of these animals. In the mouth are no teeth, only two flat white bones, one each jaw; one above, another below, with undulated surfaces, which serve instead of grinders.

The eyes are extremely small, not larger than those of a sheep: the iris black: it is destitute of ears, having only two orifices, so minute that a quill will scarcely enter them: the tongue is pointed, and small: the neck is thick, and its junction with the head scarce distinguishable; and the last always hangs down. The circumference of the body near the shoulders is twelve feet; about the belly twenty; near the tail only four feet eight: the head thirty-one inches: the neck near seven feet: and from these measurements may be collected the deformity of this animal. Near the shoulders are two feet, or rather fins, which are only two feet two inches long, and have neither fingers nor nails; beneath are concave, and covered with hard bristles: the tail is thick, strong, and horizontal, ending in a stiff black fin, and like the substance of whalebone, and much split in the fore part, and slightly forked; but both ends are of equal lengths, like that of a whale.

The skin is very thick, black, and full of inequalities, like the bark of oak, and so hard as scarcely to be cut with an ax, and has no hair on it: beneath the skin is a thick blubber, which tastes like oil of almonds. The flesh is coarser than beef, and will not
soon.

SIZE.

TAIL.

soon putrify. The young ones taste like veal. The skin used for shoes, and for covering the sides of boats.

The *Russians* call this animal *Morškaia korowa*, or Sea-cow; and *Kapušnik*, or Eater of herbs.

490. ROUND-
TAILED.

Adanson's Senegal. 259. L. E. V. M. U. S.

M with thick lips: eyes as minute as a pea: two very small orifices in the place of ears: in each jaw are nine grinding teeth; in all thirty six: neck short, and thicker than the head; the greatest thickness of the body is about the shoulders, from which it grows gradually smaller to the tail: the tail lies horizontally, is broad, and thickest in the middle, growing thinner to the edges, and quite round.


The feet are placed at the shoulders: beneath the skins are bones for five complete toes, and externally are three or four nails flat and rounded: near the base of each leg, in the female, is a small teat.

The skin is very thick and hard, having a few hairs scattered over it.

SIZE.

The length of the specimen in the LEVERIAN MUSEUM is six feet and a half; the greatest circumference, three feet eight inches; that near the tail, two feet two. This was taken near the *Marigot* of *Kantai*, in the river *Senegal*: they grow to the length of fourteen or fifteen feet: they are very fat, and both fat and lean resemble veal: but the fat adheres to the skin, in form of blubber: the negroes take them by harpooning, and sell them at the rate of

two



Round-bellied Walrus. — . 1. 190.



two long bars of iron apiece. The season is only in the months of *December* and *January*. *Manati* are found in most of the *African* rivers to the south of the *Niger*, and possibly to those on the eastern coast. The woman-fish taken off the isles *Boçicas*, to the south of the river *Cuama*, is seemingly of this species, notwithstanding the pious describer, Father *Jonanes dos Santos*, furnishes it with four tremendous tushes*.

De Buffon, xiii. 425. tab. lviii. *Raii syn.* *Trichechus Manatus.* *Lin. syst.* 49. quad. 193.

Schreber, tab. lxxx.

491. GUIANA.

M. with a head hanging downward; the feet furnished with five toes: body almost to the tail of an uniform thickness; near its junction with that part grows suddenly thin: tail flat, and in form of a *spatula*; thickest in the middle, growing thinner towards the edges.

Inhabits the rivers and sea of *Guiana*: it grows to the length of sixteen or eighteen feet: is covered with a dusky skin with a few hairs†. Those measured by *Dampier* were ten or twelve feet long: their tail twenty inches in length; fourteen in breadth; four or five thick in the middle; two at the edges: the largest (according to the same voyager) weighed twelve hundred pounds. But they arrive at far greater magnitude: *Clusius* examined one which was sixteen feet and a half long; and *Gomora* speaks of them as sometimes of the length of twenty feet.

* *Purchas*. ii. 1446.

† *Bancroft's Guiana*, 186.

492. MANATI
CLUSII.

CLUSIUS, in his *Exotics*, p. 132, gives a print and description of a *Manati* brought from the *West Indies*: but neither one or the other enables us to define the species. He says that it had short nails and broad feet; and that the tail was broad and shapeless. Till we are better informed we shall suppose it to be the same with the *Guiana*. *M. de Buffon*, in his *Supplement*, vi. 396, makes it a distinct species, under the title of *Le grand Lamantia des Antilles*.

493. ORONOKO.

THIS is the species to which *M. de Buffon* has in his *Supplement*, p. 400, given the name of *Le petit Lamantia de L' Amerique*, and says it is found in the *Oronoko*, *Oyapoc*, and the rivers of *Amazons*. This pushes its way to the amazing distance we have mentioned. By the description *Gumilla* has given of the tail, it is circular *, and probably must be referred to this species. I do not understand why *M. de Buffon* calls it *Le petit*, for it grows to a vast size. Father *Gumilla* had one taken in a distant lake, near the *Oronoko*, which was so large that twenty-seven men could not draw it out of the water: on cutting it open, he found two young ones, which weighed twenty-five pounds apiece.

We suspect that the *Manati* of the *Amazons*, &c. never visit the sea, but are perpetually resident in the fresh waters.

PLACE.

These animals abound in certain parts of the eastern coasts and rivers of *South America*, about the Bay of *Honduras*, some of the greater *Antilles* †, the rivers of *Oronoque* ‡, and the lakes formed by it; and lastly, in that of the *Amazons*, and the

* *Gumilla*, 54.† *Dampier*, i. 33.‡ *Gumilla*, ii. 43.*Guallaga*,

Gualaga, the *Pastaga*, and most of the others which fall into that vast river: they are found even a thousand leagues from its mouth, and seem to be stop'd from making even an higher advance, only by the great cataract, the *Pongo* of *Borja* *. They sometimes live in the sea, and often near the mouth of some river, into which they come once or twice in twenty-four hours, for the sake of brouzing on the marine plants which grow within their reach: they altogether delight more in brackish or sweet water, than in the salt; and in shallow water near low land, and in places secure from surges, and where the tides run gently †. It is said, that at times they frolick and leap to great heights out of the water ‡. Their uses were very considerable to the privateers or buccaneers in the time of *Dampier*. Their flesh and fat are white, very sweet and salubrious; and the tail of a young female was particularly esteemed. A suckling was held to be most delicious, and eaten roasted, as were great pieces cut out of the belly of the old animals.

The skin cut out of the belly (for that of the back was too thick) was in great request for the purpose of fastening to the sides of canoes, and forming a place for the insertion of the oars. The thicker part of the skin, cut fresh into lengths of two or three feet, serves for whips, and become, when dried, as tough as wood.

In the head, it was pretended that there were certain stones, or bones of great value, on account of their virtues in curing the gravel and colic ||.

* *Condamine*, 77.

† *Dampier*, i. 34.

‡ *Gumilla*, ii. 55.

|| *Clusii Exot.* 233. *Monardus simp. Med.* 326.

They are taken by an harpoon stuck in the end of a staff, which the *Indians* use with great dexterity. They go in a small canoe with the utmost silence, as the animal is very quick of hearing. The harpoon is loose, but fastened to a cord of some fathoms in length; for as soon as the *Manati* is struck, it swims away with the barb infixed in its body, attended by the canoe, till spent with pain and fatigue: in some places the lesser are taken in nets. If a female, which has a young one, is struck, she takes it under its fins or feet, if not too large, and shews, even in extremity, the greatest affection for its offspring; which makes an equal return, never forsaking the captured parent, but is always a sure prey to the harpooner*.

The *Indians* of the *Maragnon*, or the river of *Amazons*, take them by the means of intoxicating herbs, or by shooting them with those poisoned arrows†, whose left touch is fatal, yet imparts no degree of venom to the thing stricken, whose flesh is eaten with the utmost safety‡.

At the time the waters of the *Oronogue* (which annually overflow the banks) begin to return into the bed of the river, the *Indians* make dams across the mouths of the shallow lakes formed by the floods, and in that manner take vast numbers of *Manati*, or *Pexi-buey*, or *Fish-cows*, as the *Spaniards* call them, together with tortoises, and variety of fish||.

I conclude this account with the extraordinary history of a tame *Manati*, preserved by a certain prince of *Hispaniola*, at the time of the arrival of the *Spaniards*, in a lake adjoining to his re-

* *Dampier*, i. 37. † *Ulloa*, i. 412. *Gumilla*, ii. 46.
Trav. 34. *Pé. Tr.* xlvii. 81. ‡ *Condamine's*
 || *Gumilla*, ii. 43.

fidence. It was, on account of its gentle nature, called in the language of the country *Matum*. It would appear as soon as it was called by any of its familiars; for it hated the *Spaniards*, on account of an injury it had received from one of these adventurers. The fable of *Arion* was here realised. It would offer itself to the *Indian* favorites, and carry over the lake ten at a time, singing and playing on its back; one youth it was particularly enamoured with, which reminds me of the classical parallel in the Dolphin of *Hippo*, so beautifully related by the younger *Pliny*. The fates of the two animals were very different; *Matum* escaped to its native waters, by means of a violent flood; the *Hipponense* fish fell a sacrifice to the poverty of the retired Colonists*.

MR. *Steller* saw on the coast of *America*† another very singular animal, which he calls a *Sea Ape*; it was five feet long: the head like a dog's: ears sharp and erect: eyes large; on both lips a sort of beard: the form of its body thick and round, thickest near the head, tapering to the tail, which was bifurcated, the upper lobe the longest: the body covered with thick hair, grey on the back, red on the belly. *Steller* could discover neither feet nor paws. It was full of frolick, and played a

494. SEA APE.

* See both relations; the first in *Peter Martyr's Decades of the Indies*, Dec. iii. book 8; the other in lib. ix. epist. 33, of *Pliny*. The elder *Pliny* also relates the same story, lib. ix. c. 8.

† The *Beluga*, which I placed here in my former edition, from the misrepresentation of other writers, is an animal of the cetaceous tribe, called by the *Germans*, *Witfish*. See *Pallas Itin.* iii. 84, tab. iv. and *Grantz Greenland*, i. 114. N° 10.

- thousand

thousand monkey tricks; sometimes swimming on one side, sometimes on the other side of the ship, looking at it with great amazement. It would come so near the ship, that it might be touched with a pole; but if any body stirred, would immediately retire. It often raised one-third of its body above the water, and stood erect for a considerable time; then suddenly darted under the ship, and appeared in the same attitude on the other side; and would repeat this for thirty times together. It would frequently bring up a sea-plant, not unlike the bottle gourd, which it would toss about, and catch again in its mouth, playing numberless fantastic tricks with it.

DIV. IV.

WINGED QUADRUPEDS.

D I V. IV. Winged Quadrupeds :

XLIV. BAT.

With long extended toes to the fore feet, connected by thin broad membranes, extending to the hind legs.

* Without Tails.

495. TERNATE.

Vespertilio ingens. Chuf. exot. 94.

Canis volans ternatanus orientalis. Sel.

Muf. i. 91. *tab.* lvii.

Vespertilio Vampyrus. V. ecaudatus, naso simplici, membrana inter femora divisa. Lin. Syst. 46.

La Rouffette & la Rougette. De Buffon,

x. 55. tab. xiv. xvii.. Schreber, 185. tab. xlii.*

Pteropus rufus aut niger auriculis brevibus acutiusculis. Briffon quad. 153, & 154. *No. 2. Shaw Spec. Lin.* viii.

Great Bat. Edw. 180. Br. Mus. Afr. Mus. Lev. Mus.

THE ROUSETTE.

B. with large canine teeth: four cutting teeth above, the same below: sharp black nose: large naked ears: the tongue is pointed, terminated by sharp aculeated *papillæ*: exterior toe detached from the membrane: the claw strong, and hooked: five toes on the hind feet: talons very crooked, strong, and compressed sideways: no tail: the membrane divided behind quite to the rump: head of a dark ferruginous color: on the neck, shoulders, and under side, of a much lighter and brighter red: on the back the hair shorter, dusky, and smooth: the membranes of the wings dusky: varies in color; some entirely of a reddish brown; others dusky. This now described was one foot long: its extent from tip to tip of the wings four feet; but they are found vastly larger.


SIZE.

* The History of these bats has been greatly elucidated by M. *De la Nux*, who resided fifty years in the *Ile de Bourbon*, where they are found. See M. *de Buffon*, Suppl. iii. 253.

This



Fecula Bat — N^o 493.



This species is not gregarious, yet they are found in numbers on the same tree, by accidentally meeting there in search of food: they fly by day, and are seen arriving one by one to the spot which furnishes subsistence. If by any accident they are frightened, they will then quit the tree in numbers, and thus fortuitously form a flock. It is different with the other species.

The Rougette*, or BAT, with the same kind of teeth as the other, and the shape of head and body the same: the whole body and head cinereous, mixed with some black; but on the neck is a great bed of lively orange, or red.

THE ROUGETTE.

The size is much less; the extent of wings being little more than two feet.

SIZE.

They are both inhabitants of the same countries, agree in their food, but differ in some of their manners, which I shall distinguish in the following history of them.

These monsters inhabit *Guinea*, *Madagascar*, and all the islands from thence to the remotest in the *Indian* ocean. They are found again in *New Holland*†, the *Friendly islands*, the *New Hebrides*, and *New Caledonia*‡. The *Rougettes* fly in flocks, and perfectly obscure the air with their numbers: they begin their flight from one neighboring island to another immediately on sun-set, and return in clouds from the time it is light till sun-rise||, and lodge during day in hollow trees: both live on fruits; and are so fond of the juice of the *palm-tree*, that they will intoxicate themselves with it till they drop on the ground§.

PLACE.

Notwithstanding the size of their teeth, they are not carnivorous. Mr. *Edwards* relates, that they will dip into the sea for

* *LEV. MUS.* † *Cook's voy.* iii. 626. ‡ *Forster's obs.* 187. || *Dampier's voy.* i. 381. § *Museum Hafniæ, Pars i. Sect. 2. No. 18.*

fish. I suspect that fact; but it is known that they skim the water with wonderful ease, perhaps in sportive moods. They also frequent that element to wash themselves from any vermin which might adhere to them*. They swarm like bees, hanging near one another from the trees in great clusters†; at least five hundred were observed hanging, some by their fore, others by their hind legs, in a large *Casuarina*-tree, in one of the *Friendly islands*. When shot at, they flew from the boughs very heavily, uttering a shrill piping note; others again, arrived at intervals from remote places to the tree‡. In *Neve Caledonia*, the natives use their hair in ropes, and in the tassels of their clubs, interweaving it with the threads of the *Cyperus squarrosus*. The *Indians* eat them, and declare the flesh to be very good: they grow excessively fat at certain times of the year. The *French*, who live in the *Isle de Bourbon*, boil them in their *bouillon*, to give it a relish||. The *Negroes* have them in abhorrence§. Many of the *Roussettes* are of an enormous size: *Beeckman*** measured one, whose extent from tip to tip of the wing was five feet four inches; and *Dampier*†† another, which extended further than he could reach with stretched-out arms. Their bodies are from the size of a pullet to that of a dove: while eating, they make a great noise: their smell rank; their bite, resistance, and fierceness great when taken.

They bring but one young at a time.

The ancients had some knowledge of these animals. *Herodotus*‡

* *Forster's obs.* 188.

† *Argensola Philip. isles*, 158. *Des Marchais*, ii. 261.

‡ *Forster's voy.* i. 450.

|| *Voy. de la Caille*, 233. § *Des Marchais*, *ibid.*

** *Voy. to Borneo*, 39.

†† i. 381.

‡ *Σηρια πλεγμα, της νυκτερις*

περσικηλα. Lib. iii.

mentions

mentions certain winged wild beasts, like bats, that molested the *Arabs*, who collected the *Cassia*, to such a degree that they were obliged to cover their bodies and faces, all but their eyes, with skins. It is very probable, as *M. de Buffon* remarks, it was from such relations the Poets formed their fictions of *Harpies*.

Linnaeus gives this species the title of *Vampyre*, conjecturing it to be the kind which draws blood from people in their sleep. *M. de Buffon* denies it, ascribing that faculty to a species only found in *S. America*: but there is reason to imagine, that this thirst after blood is not confined to the bats of one continent, nor to one species; for *Bontius* and *Nieuboff* inform us, that they of *Java** seldom fail attacking those who lie with their feet uncovered, whenever they can get access; and *Gumilla*†, after mentioning a greater and lesser species, found on the banks of the *Orenoque*, declares them to be equally greedy after human blood. Persons thus attacked, have been known to be near passing from a sound sleep into eternity. The Bat is so dextrous a bleeder as to insinuate its aculeated tongue into a vein without being perceived, and then suck the blood till it is satiated; all the while fanning with its wings, and agitating the air, in that hot climate, in so pleasing a manner, as to lull the sufferer into a still sounder sleep‡. It is therefore very unsafe to rest either in the open air, or to leave open any entrance to these dangerous animals; but they do not confine themselves to human blood; for *M. Con-*

* *Bontius India*, 70. *Nieuboff*, 255. These writers say that this kind is as big as a pigeon. I suspect that the species just described is common to *India* and *S. America*; *Mr. Greenwood*, painter, long resident at *Surinam*, informing me that there is in that colony a fox-colored bat, whose extent of wings is above four feet.

† *Hist. Orenoque*, iii. 100.

‡ *Ulloa's voy.* i. 61.

damine * says, that in certain parts of *America* they have destroyed all the great cattle introduced there by the missionaries.

β. LESSER. B. with head like a grehound: large teeth like the former: ears long, broad, and naked: whole body covered with soft short hair of a straw-color: shaped like the other in all respects: length, eight inches three quarters; extent, two feet two inches. Place unknown to the gentleman who favored me with it. LEV. MUS.

496. SPECTRE.

Andira-gnacu, *vespertilio cornutus*. *Piso*
Brazil. 190. *Margrave* *Er. fil.* 213.
Canis volans maxima aurita *fam. ex*
Nov. Hispania. *Seb. Mus. i. tab. lvii.*
Vespertilio spectrum. *V. ecaudatus*, *naso*
infundibuliformi lanceolato. *Lin. fff.*

46. *Klein quad.* 62.
Pteropus auriculis longis, patulis, naso
membrana antrorsum inflexa aucto.
Eriffon quad. 154.
Le Vampire. *De Buffon*, x. 55. *Schreber*,
192. *tab.* xlv.

B. with a long nose: large teeth: long, broad, and upright ears: at the end of the nose a long conic erect membrane, bending at the end, and flexible: hair on the body cinereous, and pretty long: wings full of ramified fibres: the membrane extends from hind leg to hind leg: no tail; but from the rump extend three tendons, terminating at the edge of the membrane. By *Seba's* figure, the extent of the wings is two feet two inches; from the end of the nose to the rump seven inches and a half.

SIZE.

PLACE.

Inhabits *South America*: lives in the palm-trees: grows very fat: called *Vampire* by *M. de Buffon*, who supposes it to be the species that sucks human blood: but neither *Piso*, or any other writers who mention the fact, give the least description of the kind.

* *Voy. S. America*, 85.

Vespertilio Americanus vulgaris. *Seb. Mus.* i. tab. lv. fig. 2.

Son quad. 161.

Vespertilio periphiatus *V. ocaudatus* naso foliato acuminate. *Lac. f. fig.* 47.

La chauve souris fer de Lance. *D. Buffon.* xiii. 276. t. b. xxxiii. *Suz. l'oe.* vii. 202 tab. lxxv. *Schreber.* 194. tab. xlv. B.

V. murina coloris pedibus anticis tetradactylis, posticis pentadactylis. *Erzf.*

497. JAVELIN.

B. with large pointed ears: an erect membrane at the end of the nose, in form of the head of an ancient javelin, having on each side two upright processes: no tail: fur cinereous: size of a common bat.

SIZE.

Inhabits the warm parts of *America*.

PLACE.

The bat described by Mr. *Schreber*, p. 193. tab. xlv. A. under the title of *La Chauve souris pelle*, has so much resemblance, that I place it here as a variety of the former: the nasal membrane being nearly of the same form; the color differs, the fur being ferruginous.

Vespertilio, rostro appendice auriculæ forma donata *Sloane Jam.* ii. 330. Small bat. *Edw.* 201. fig. 1. *La Feuille.* *De Buffon*, xiii. 227.

Vespertilio forcinus. *Pallas Miscel.* 48. tab. v.* *Schreber*, 195. tab. xlvii. LEV. MUS.

498. LEAF.

B. with small rounded ears: membrane on the nose of the form of an ovated leaf: no tail: a web between the hind legs: fur of a mouse-color, tinged with red: size of the last.

SIZE.

* This seems to be one of the blood-sucking species, the tongue being furnished with aculeated *papillæ*, and is twice the length of the nose; so is well adapted for that purpose.

Inhabits

PLACE.

Inhabits *Jamaica*, *Surinam*, and *Senegal*: in the first lives in caves in woods, which are found full of its dung, productive of salt-petre : feeds on the prickly pear.

499. CORDATED.

Glis volans Ternatanus. Seb. Mus. i. tab. lvi. fig. 1. Schreber, 191. tab. xlviii. *Vespertilio spafina. V. ecaudatus naso foliato obcordato. Lin. ffs. 47.*

B. with very broad and long ears : at the end of the nose a heart-shaped membrane : no tail : a web between the hind legs : color of the face a very light red ; that of the body still paler.

PLACE.

Inhabits *Ceylon*, and the isle of *Ternate*, one of the *Moluccas*.

* * * With tails.

500. PERUVIAN.

Chauve-fouris de la Vallée d'Ylo. Feuille tab. lx. obs. Peru, 1714. p. 623. Schreber 196. Vespertilio Leporinus. Gm. Lin. 47.

B. with a head like a pug-dog : large strait ears, sharp at the ends and pointing forwards : two canine teeth, and two small cutting teeth between each, in each jaw : tail enclosed in the membrane which joins to each hind leg, and is also supported by two long cartilaginous ligaments involved in the membrane : color of the fur iron-grey ; but erroneously colored in the print, of a straw color : body equal to that of a middle-sized rat : extent of wings two feet five inches.

SIZE.

β. With

β. With a large head and hanging lips, like the chops of a mastiff: nose bilobated: upper lip divided: strait, long, and narrow ears, sharp-pointed: teeth like the former: tail short; a few joints of it stand out of the membrane, which extends far beyond it; is angular, and ends in a point: claws on the hind feet large, hooked, and compressed sideways: membranes of the wings dusky, very thin: fur on the head and back brown; on the belly, cinereous. Length, from the nose to the end of the membrane, above five inches; extent of wings, twenty.

SIZE.

Inhabits *Peru* and the *Mosquito* shore: the last was given me by *John Ellis*, Esq; F.R.S. It differed from the former in size, being less; in all other respects agreed.

PLACE.

Linnaeus, carried away by love of system, places this, on account of its having only two cutting teeth in each jaw, among the *Glires*, next to the squirrels, under the name of *Noctilio Americanus*. But such is the variety in the numbers and disposition of the teeth in the animals of this genus, that he might form almost as many genera out of it as there are species. But as the Bats have other such striking characters, it is unnecessary to have recourse to the more latent marks to form its definition. The same may be said of several other animals.

Autre Chauve souris. *De Buffon*, x. 84, 87. tab. xix. fig. 1, 2. *Schreber*, 207. tab. xlix. *LEV. MUS.*

501. BULL-DOG.

B. with broad round ears, the edges touching each other in front: nose thick: lips pendulous: upper part of the body of a deep ash-color; the lower paler: tail long; the five last joints

SIZE. joints quite disengaged from the membrane. Length above two inches; extent nine and a half.

PLACE. Inhabits the *West Indies*.

502. SENEGAL.

Chauve-fouris etrangere. *De Buffon*, x. LEV. MUS.
82. tab. xvii. *Schreber*, 206. tab. lviii. *Vespertilio nigrita*. *Gm. Lin.* 49.

B. with a long head: nose a little pointed: ears short, and pointed: head and body a tawny brown mixed with ash-color: belly paler: two last joints of the tail extend beyond the membrane. Length from nose to rump, above four inches; extent 21.

PLACE. Inhabits *Senegal*.

503. POUCH.

La Chauve-fouris a bourse. *Schreber*, 209. tab. lvii.

WITH the nose somewhat produced: the end thickest, and beset with fine whiskers: the chin divided by a sulcus: ears long, rounded at their ends: on each wing, near the second joint, is a small purse, or pouch: the tail is only partly involved in the membrane; the end hanging out: color of the body a cinereous brown: the belly paler.

SIZE. Length an inch and a half.

PLACE. Inhabits *Surinam*.

Autre Chauve-fouris *de la Guyanne. De Buffon, Supplem. vii. 214. tab. lxxv.*

504. SLOUCH-
EARED.

B. with large pendulous ears, pointed at the ends: nose obtuse at the end: tail long, included in the membrane, and ending with a hook: color above, a deep chefnut; lighter on the belly, and cinereous on the sides: length three inches and four lines: extent of wings fifteen inches.

Inhabits *Guiana*.

PLACE.

Autre Chauve-fouris. *De Buffon, x. 92. tab. xx. fig. 3. Schreber, 204. tab. lvi.*
LEV. MUS.

505. BEARDED.

B. with the nostrils open for a great way up the nose: hair on the forehead and under the chin very long: ears long and narrow: upper part of the head and body of a reddish brown; the lower of a dirty white tinged with yellow: tail included in a membrane very full of nerves. A small species.

B. with a head shaped like that of a mouse: top of the nose a little bifid: ears short, broad, and rounded: no cutting teeth; two canine in each jaw: tail very long, inclosed in the membrane, which is of a conic shape: head, body, and the whole upper side of the membrane which incloses the tail, covered with long very soft hair of a bright tawny color, lightest on the head and beginning of the back; the belly paler: at the base of each wing a white spot: wings thin, naked, and

506. NEW YORK.

VOL. II.

S s

dusky:

- SIZE.** dusky: bones of the hind legs very slender. Length, from nose to tail two inches and a half; tail one inch eight-tenths; extent of wings ten and a half.
- PLACE.** Inhabits *North America*. Communicated by Mr. *Astton Blackburne* *. It is also found in *New Zealand* †. Mr. *Schreber* describes it from me, in p. 212. *LEV. Mus.*

507. STRIPED. Autre Chauve-souris. *De Buffon*, x. 92. tab. xx. fig. 3. *Zooph. Grenov. No. 25.*
Schreber, 205. tab. xlix.


- B.** with a small short nose; ears short, broad, and pointing forward: body brown: wings striped with black, and sometimes with tawny and brown. Length, from nose to the end of the tail, two inches: varies in color; the upper part of the body being sometimes of a clear reddish brown, the lower whitish.
- SIZE.**
- PLACE.** Inhabits *Ceylon*; called there, *Kiriwoula* ‡. I may add to this little species of Bat, the mention of a minute kind seen and heard in myriads of numbers in the isle of *Tanna*, one of the *New Hebrides*, but which escaped every attempt of our voyagers to obtain a near examination ||.

* The Rev. M. *Clayton* mentions another species of *North American Bat*; large, with great ears, and long straggling hairs. *Phil. Transf. abridg.* iii. 594.

† *Forster's observ.* 189.

‡ *Pallas Miscel.* 49.

|| *Forster's obs.* 188.



P. Magell. Sculp.

1. *Liber. Ternate. Pa. p. 308.*
2. *New York. 1806*



Vespertilio Cephalotes. *Pallas Spicil. Zool. fasc.* iii. 10. *tab. i. Schreber*, 203. *tab.*
lxi. LEV. MUS.

503. *MOLUCCA.*

B. with a large head: thick nose: small ears: tubular nostrils, terminating outwards in form of a screw: upper lip divided: tongue covered with papillæ and minute spines: claw, or thumb, joined to the wing by a membrane: first ray of the wing terminated by a claw: end of the tail reaches beyond the membrane: color of the head and back greyish ash-color; that in the *LEVERIAN MUSEUM* of a fine straw-color: the belly dull white. Length, from nose to rump, three inches three quarters; extent of wings about fifteen.

SIZE.

Inhabits the *Molucca* isles. Described first by that very able naturalist Doctor *Pallas*.

PLACE.

Vespertilio Lepturus. *Schreber*, *tab. lvii. Gm. Lin.* 50.

509. *SLENDER-
TAILED.*

B. with tubular nostrils: long erect ears: color dusky above, cinereous beneath.

Inhabits *Surinam*.

PLACE.

Vespertilio Lasiurus. *Schreber*, *tab. lxii. Gm. Lin.* 50.

510. *ROUGH-
TAIL'D.*

B. with upright small ears: tail broad at the base, terminating in a point thickly covered with hair: color a reddish brown: a small species.

Place unknown.

511. LASCOPE-
RUS.Vespertilio Lascopterus. *Schreber*, tab. lviii. B. *Gm. Lin.* 50.

B. with a most prominent rounded forehead : short nose : color a bright rust : upper part of the wings of a paler rust : ends and lower parts of the wings black. By Mr. *Schreber's* figure it seems a large species.

Place unknown.

512. HORSE-SHOE.

La Chauve-fouris fer a Cheval. *De Buffon*, viii. 131, 132. tab. xvii. xx. *Schreber*, 210. tab. lxii. *Br. Zool.* i. 129.

B. with a membrane at the end of the nose in form of a horse-shoe : ears large, broad at their base, and sharp-pointed, inclining backward : wants the little or internal ear : color of the upper part of the body deep cinereous ; of the lower, whitish. There is a greater and lesser variety ; the greater is above three inches and a half long from the nose to the tip of the tail : its extent above fourteen. This and all the following have the tail inclosed in the membrane.

SIZE.

PLACE.

Inhabits *Burgundy* ; and has lately been discovered in *Kent*, by Mr. *Latham*, of *Dartford* ; found also about the *Caspian* sea. The long-eared Bat, No 519, has also been observed there, and at *Petersburg*. This and the four next were first discovered by M. *de Buffon*, whose names I retain.

Is. Noctule. De Buffon, viii. 128, tab. xviii. Schreber, 200. tab. lii.
Great Bat. Br. Zool. illustr. tab. ciii. Br. Zool. i. 128.

513. NOCTULE.

B. with the nose slightly bilobated: ears small and rounded: on the chin a minute *verruca*: hair of a reddish ash-color. Length to the rump two inches eight-tenths; tail one seven-tenth; extent of wings thirteen inches.

SIZE.

PLACE.

Inhabits *Great Britain* and *France*; very common in the open deserts of *Russia*, wherever they can find shelter in caverns: flies high in search of food, not skimming near the ground. A gentleman informed me of the following fact, relating to those animals, which he was witness to:—that he saw taken under the eaves of *Queen's College, Cambridge*, in one night, one hundred and eighty-five; the second night sixty-three; the third night two; and that each that was measured had fifteen inches extent of wings*.

La Serotine. De Buffon, viii. 129, tab. xviii. Schreber, 201. tab. liii.

514. SEROTINE.

B. with a longish nose: ears short, but broad at the base: hair on the upper part of the body brown, mixed with ferruginous; the belly of a paler color. Length from nose to rump, two inches and a half: no tail.

SIZE.

* No notice was taken of the species; but, by the size, it could be neither of the common kinds. I never saw but one specimen of the *Noctule*, which was caught during winter in *Flintshire*.

Inhabits.

PLACE.

Inhabits *France*; found in caverns of rocks upon the river *Argun*, beyond lake *Baikal*; but as yet not discovered in any other part of the vast *Russian* dominions.

515. GREATER
SEROTINE.

La Grande Serotine de la *Guyanne*. *De Buffon Supplem.* vii. 289. tab. lxiii.

B. with a very long, strait and strong nose, sloping down at the end: ears long, erect, dilated towards the bottom, rounded at the end: color of the upper parts of a reddish chestnut; sides a clear yellow, rest of a dirty white. Length five inches eight lines: extent of wings two feet: no tail.

PLACE.

Inhabits *Guiana*: assembles in vast numbers in open places, particularly meadows; and flies in company with the goat-suckers, and both together, in such numbers as to darken the air.

516. PIPIST-
RELLE.

La Pipistrelle. *De Buffon*, viii. 129. tab. xix. fig. 2. *Schreber*, 202. tab. liv.

SIZE.

B. with a small nose: the upper lip swelling out a little on each side: the ears broad: the forehead covered with long hair: color of the upper part of the body a yellowish brown; the lower part dusky; the lips yellow. The least of Bats; not an inch and a quarter long to the rump: extent of wings six and a half.

PLACE.

Inhabits *France*: common in the rocky and mountainous parts of *Russia* and *Siberia*.

La Barbastelle. *De Buffon*, viii. 130. *tab. xix. fig. 1. Schreber*, 203. *tab. lv.*

517. BARBAS-
TELLE.

B. with a sunk forehead: long and broad ears; the lower part of the inner sides touching each other, and conceal the face and head when looked at in front: the nose short; the end flattened: cheeks full: the upper part of the body of a dusky brown; the lower, ash-colored and brown. Its length to the rump about two inches; its extent ten and a half.

SIZE.

Inhabits *France*.

PLACE.

Nexregis. Arist. hist. an. lib. i. c. 5.

Vespertilio. Plin. lib. x. c. 61. Gefner

quad. 766. Agricola Anim. Subter. 483.

Bat. Flitter-moufe. Raii syn. quad. 243.

Rear-moufe. Chariton Ex. 80.

Vespertilio major. Speck-maus, Fleder-

maus. Klein quad. 61.

Vespertilio murinus. V. caudatus naso

oreque simplici, auribus capite minoribus. Lin. syst. 47.

Laderlap, Fladermus. Faun. succ. No. 2.

La grande Chauve-souris de notre pais.

Briffon quad. 158. De Buffon, viii. 113.

tab. xvi.

Short-eared Bat. Br. Zool. i. 130. Edw.

201. Schreber, 199. tab. li. Lev. Mus.

518. COMMON.

B. with short ears: mouse-colored fur tinged with red. Length two inches and a half; extent of wings nine.

SIZE.

Inhabits *Europe*: the most common species in *England*.

PLACE.

512. LONG-EAR.
ED.

Souris Chauve, Rattepenade. *Belin* *exf.*
147.

Vesperugo auritus. V. naso oreque simplici, auriculis duplicatis, capite majoribus. *Lac. f. 47. Faun. f. 4c. No. 3.*
Klin quad. 61.

La petite Chauve-fouris de notre pais.

Briffon quad. 160. Sharv spec. Lin. vii.
L'Orcillar. De Buffon, viii. 118. tab.

xvii. *Schreber, 197. tab. l.*

Long-eared Bat. *Edw. 201. Br. Zool. i.*
129. *Br. Zool. illustr. tab. ciii. LEV.*
Mus.

SIZE.

B. with ears above an inch long, thin, and almost pellucid: body and tail only one inch three quarters long. This and all other Bats, except the *Ternate* and the *Horse-shoe*, have a lesser or internal ear, serving as a valve to close the greater when the animal is asleep.

PLACE.

Inhabits *Europe*, and is found in *Great Britain*. Bats appear abroad in this country early in the spring; sometimes are tempted by a warm day to sally out in winter; fly in the evenings; live on moths and other nocturnal insects; skim along the water in quest of gnats; fly by jerks, not with the regular motion of birds, for which the antients mistake them; frequent glades and shady places; will go into larders, and gnaw any meat they find: bring two young at a time, which they suckle at their breast: retire at the end of summer into caves, the eaves of houses, and into ruined buildings, in vast multitudes, where they generally remain torpid, suspended by the hind legs, enveloped in their wings: are the prey of owls: their voice weak. *Ovid* takes notice both of that and the origin of the *Latin* name.

Minimam pro corpore vocem

Emittunt; peraguntque leves stridore querelas.

Tesque, non sylvas celebrant: lucemque perose

Nocte volant: seroque trahunt a vespere nomen.

ADDITIONS.

A PROOF of their being prolific was produced by Mr. *Tullo*, in the parish of *Newtyle*, in the shire of *Forfar*, about twenty years ago, when a she-mule, which he turned to a horse, brought a foal which much resembled the female parent. But as there is a superstition in *Scotland* about these productions, the foal was put to death, being considered as a monster.

MULES.
page 8.

SUMATRAN ANTELOPE,

As communicated by Doctor *Shaw*.

ANTELOPE.
p. 104.

Camblingootan, or Goat of the woods, *Marsden's Sumatra*, 93.

SIZE of a common goat, but stands considerably higher on its legs: color an uniform black, but each hair when narrowly examined is grey towards the base: on the top of the neck just above the shoulders a patch of whitish, bristly, long straight hair, much stronger than the rest, and having somewhat the appearance of a partial mane: on each side of the lower jaw a longitudinal patch of yellowish white: ears moderate, marked internally with three obscure longitudinal bands of white, as in some of the antelopes: horns six inches long, bending slightly backwards, sharp-pointed, black and annulated near half their length with

VOL. II. T t prominent

prominent rings: tail about the length of horns, and sharpish: hoofs rather small and black: hair on the whole animal rather harsh, and not lighter-colored below, or on the belly, than above.

In its disposition it is wild and fierce, and is said by the natives to be remarkably swift: we are obliged to the author of the elegant history of *Sumatra* for the discovery of this animal.

MONKEY.
p. 226.

PROBOSCIS MONKEY.

La Guenon a long nez. *De Buffon Supplém.* vii. 53. tab. xi. xii.

M. with the nose projecting very far beyond the mouth, like the human, but divided in the middle by a shallow furrow: in the profile it exactly resembles a long proboscis, and makes a ridiculous appearance: the forehead hangs far over the base of the nose: the face is hooked, of a brown color, marked with blue and red: the head covered with thick hair of a chestnut brown: the ears broad, thin, and naked, hid in the fur: the body is large, clothed with hair of a brown chestnut color; orange on the breast: round the throat, neck, and shoulders, the hair is longer than that on the rest of the body, and forms a sort of short cloak, of a color contrasting that of the face: the legs are covered with short tawny hair: the length from the tip of the nose to the base of the tail is two feet: of the tail, above two feet.

SIZE.

PLACE.

Inhabits the *East Indies*; but the particular part is not mentioned.

THE



Proboscis . Monkey.

J. Stanger del.






Proboscis - Monkey.







Heart-marked . Marmoset.

THE HEART-MARK'D MAUCAUCO.

MAUCAUCO.
P. 234.

I HAVE totally forgotten the friend who obliged me with the drawing of this animal, and the place it came from; but probably from *Madagascar*, or the neighboring isles, the seat of most of the congenerous species.

All the upper parts of the body are of a deep cinereous brown: the face marked with large white heart-shaped spots: the broader part extends between the ears; the point reaches almost to the nose: the belly, legs, and feet, are white. I am at a loss for the size; but possibly the gentleman from whom I received the drawing may reveal himself, and communicate the wanted particulars.

ONE which was examined at the *Cape of Good Hope*, by Captain *Blanket*, had ears like those of a lurcher, but larger, and more on the top of the head. It could turn them on all sides with great facility: feet flatter than those of other dogs. It could not bark or howl, but only cried: was very fierce, and mastered the tame dogs it was with, though it was only a young one.

WILD DOGS.
P. 236.

SLENDER TOED WEESEL.

W with short rounded ears: fur soft and fine, grizzled minutely with black and rufous: toes very long and slender; five in number; each lobated at the bottom of the first

T t 2

joint:

A D D I T I O N S.

joint: claws small: the upper part of the toes and part of the legs covered with short velvet-like down.

Length from the tip of the nose to the base of the tail seven inches: tail about the same length; bushy or covered with long hairs of the same color with the rat.

PLACE.

A native of *Cochin China*.


THE ERMINED WEESEL.

W with ears short, round and naked: within of a fine pink color: tip of the nose black: head white and plain; the rest of the body and tail white: the first spotted with ermine-like black spots, disposed in rows from neck to tail, on the sides as well as back: the tail annulated with black: the hairs on all parts of the tail short, only the end is tufted with black.

The legs remarkably strong, and thick covered to the very claws with long bright ferruginous hairs: claws sharp and white: length of the head three inches and a half; of the neck and body from head to tail sixteen inches and a half; of the tail eleven and a half.


This elegant animal is likewise a native of *Cochin China*, and with the former, communicated to us by the friendship of Lieut. Col. *Davies*, of the artillery.

INDEX.



Slender toed Weasel.



*Ermine Weasel*



I N D E X *.

A		Vol. Page			Vol. Page
ANT-EATER, or Ant-Bear	—	II. 256	Bull-Dog	—	I. 242
ANTELOPES, their general history	—	I. 68	C		
— Species of	—	I. 70	CAMEL, Arabian	—	I. 129
APES, their general history	—	I. 178	— Baetrian	—	I. 132
— Sea	—	II. 301	— Peruvian, or Llama	—	I. 133
ARMADILLO	—	II. 246	— The only native beast of		
AS	—	I. 8	— burden in America	—	I. 134
— Wild	—	I. 8	— Vicunna	—	I. 136
Axis	—	I. 117	— Paicos	—	I. 137
— Greater	—	I. 118	— Guanaco	—	I. 138
B			— Chilihucque	—	I. 138
Baboons	—	I. 188	Camelopard	—	I. 65
Baby-rouffa	—	I. 148	Castoreum	—	II. 118
BADGER	—	II. 14	CAT, Common	—	I. 295
BATS	—	II. 304	— Wild	—	I. 296
BEAR	—	II. 1	— Tiger	—	I. 277
— Polar	—	II. 5	— Mountain	—	I. 300
BEAVER	—	II. 114	— Civet	—	II. 70
— Its wondrous œconomy	—	II. 115	— Angora	—	I. 296
— Sea, <i>vide</i> Sea Otter.	—	II. 83	CAVY, various species of	—	II. 88
Beaver-Eater	—	II. 9	Chamois	—	I. 72
Beluga	—	II. 301	Chimpanzee	—	I. 180
Bezoar	—	I. 58	Civet	—	II. 70
Bison, Scottish	—	I. 17	D		
Buck	—	I. 113	DEER,	—	I. 105
Buffalo, Indian	—	I. 28	— Elk, or Moofe	—	I. 105
— When introduced into			— Rein	—	I. 111
— Europe	—	I. 29	— Fallow	—	I. 113
— American	—	I. 23	— Mexican	—	I. 122
— Dwarf, or Anoa	—	I. 30, 36	— Porcine	—	I. 119
— Naked, or Bonafus	—	I. 30	— Grey	—	I. 123
— Ceylon	—	I. 31	— Virginian	—	I. 116
Bull	—	I. 16	— Red, Stag, or Hart	—	I. 114

* In this Index very few of the Species are enumerated, that having been amply done under the INDEX of GENERA; to which the Reader is referred, the Genera being here printed in capitals for that purpose, under which he will find all the Species belonging to each.

DEER,

I N D E X.

	Vol.	Page		Vol.	Page
Mandrill	—	I. 190	Panther	—	I. 280
Manis	—	II. 252	Pecary	—	I. 147
MARMOTS	—	II. 128	Pekan	—	II. 51
Martin	—	II. 41	Pig, Guinea	—	II. 90
— Pine	—	II. 42	Pole-cat	—	II. 37
MAUCAUCOS	—	I. 227	— American	—	II. 64
Minx	—	II. 81	PORCUPINE	—	II. 122
Mococo	—	I. 230	— incapable of darting	—	II. 123
Mole Rats	—	II. 214	— its quills	—	II. 123
MOLES	—	II. 229	Potto	—	II. 59
Mongooz	—	I. 229	Puma	—	I. 289
MONKIES	—	I. 199	Pygmies, what	—	I. 183
Moose	—	I. 105			
MORSE	—	II. 266			
Moufe	—	II. 184	Quagga	—	I. 14
MULE, Wild	—	I. 4	Quick-hatch	—	II. 8
Mule	—	I. 8	Quojas Morrou	—	I. 180
Mufimon	—	I. 44			
Musk, Animal	—	I. 124			
— Rat	—	II. 221			
	N			R	
Norway Rat	—	II. 178	Rabbit	—	II. 103
	O		Raccoon	—	II. 12
Once	—	I. 285	RAT	—	II. 172
OPOSSUM	—	II. 18	— Norway	—	II. 178
Orang Outang	—	I. 180	— Water	—	II. 182
OTTER	—	II. 77	— Musk	—	II. 221
Otter, Sea	—	II. 83	Ratel	—	II. 66
Ox	—	I. 16	Rein Deer	—	I. 111
— Great Indian	—	I. 20, 21	RHINOCEROS	—	I. 150
— Abyssinian	—	I. 21	River Hog	—	II. 88
— Madagascar	—	I. 21	Roebuck	—	I. 120
— Tinian	—	I. 21			
— Lant or Dant	—	I. 21		S	
— Holstein and Jutland	—	I. 21	Sable	—	II. 43
— Podolian and Hungarian	—	I. 22	Schakal	—	I. 261
— Grunting	—	I. 24	Sea Bear	—	II. 281
— Musk	—	I. 31	— Ape	—	II. 301
— Cape	—	I. 32	— Calf	—	II. 270
— American	—	I. 23	— Cow	—	I. 266
	P		— Horfe	—	II. 266
Pacos	—	I. 137	— Lion	—	II. 286
			SEAL	—	II. 270
			SH+EP	—	I. 37
			— Cretan	—	I. 38
			— Hornlefs	—	I. 39
			— Many-horned	—	I. 39
					African

I N D E X.

	Vol. Page		Vol. Page
SHEEP, African	— I. 40	V	
— Broad-tailed	— I. 41	Vampire	— II. 307
— Sibirian	— I. 44	Vanfire	— II. 51
— Corfican	— I. 45	Vicunna	— I. 136
— Bearded	— I. 52	Vifon	— II. 51
SHREW Mouse	— II. 221		
Siyah Ghufh	— I. 305	W	
Skunk	— II. 65	WALRUS	— II. 266
SLOTH	— II. 240	Warree	— I. 141
SQUIRRELS	— II. 138	Water Elephant	— I. 157
Stag	— I. 114	WEESEL	— II. 33
Stoat	— II. 35	Wolf	— I. 248
Strepsiceros	I. 38, 88	Wolverene	— II. 8
Sukotyro	— I. 175		
		Y	
T		Yfarus	— I. 72
Tapiir	— I. 163		
Tiger	— I. 277	Z	
— Hunting	— I. 284	Zebra	— I. 13
		Zerda	— I. 267
U		Zibet	— II. 72
Unicorn	— I. 155	Zorrina	— II. 66
Urchin	— II. 234		

I N D E X

OF THE

NAMES of QUADRUPEDS,

IN THE

ANCIENT CLASSIC WRITERS,

IN THE

WORKS of M. DE BUFFON,

AND IN OTHER AUTHORS.

A		Vol.	Page			Vol.	Page
ABBADOS	—	I.	154	Ane	—	I.	8
Acanthion	—	II.	122	Anta	—	I.	163
Addax	—	I.	89	Antelope	—	I.	68
Adil	—	I.	261	Apar	—	II.	246
Adimain	—	I.	40	Aper	—	I.	140
Adive	—	I.	261	Aperea	—	II.	90
Ægagrus	—	I.	57	Aquiqui	—	I.	214
Agouti	—	II.	94	Arabata	—	I.	215
Ahu	—	I.	93	Arctomys	—	II.	131
Ai	—	II.	240	Arctopithecus	—	II.	240
Aigrette	—	I.	207	Argali	—	I.	44
Akouchy	—	II.	95	Arnæb	—	II.	98
Alagh-daagha	—	II.	169	Arctæb	—	II.	1
Alce	—	I.	105	Armadillo	—	II.	246
Algazel	—	I.	77	Athnoko	—	II.	92
Allo-camelus	—	I.	133	Aspalax	—	II.	216
Allouatte	—	I.	215	Asapanick	—	II.	153
Alpaco	—	I.	137	Attarfoak	—	II.	279
Amboimenes	—	I.	230	Aurochs	—	I.	17
Anak el Ard	—	I.	306	Axis	—	I.	117
Andira-guacu	—	II.	308	Aye, Aye	—	II.	142
Vol. II.				U u		Ayotochtli	

INDEX OF CLASSICAL NAMES, &c.

Ayotochtli	_____	Vol. Page II. 248	Brebis	_____	Vol. Page I. - 37
	B		Bubalus	_____	I. 28, 102
Babouin	_____	I. - 194	Bubel	_____	I. - 24
Baby-rouffa	_____	I. - 148	Bucula	_____	I. - 103
Bafwer	_____	II. 114	Buffle	_____	I. - 28
Baieu	_____	I. - 122	Bufelaphus	_____	I. - 102
Bar	_____	II. 1		C	
Baraba	_____	II. 213	Caaigora	_____	I. - 147
Barbarefque	_____	II. 150	Cabiai	_____	II. 88
Barbastelle	_____	II. 319	Cabionora	_____	II. 88
Barrys	_____	I. - 180	Cachicame	_____	II. 248
Becheti	_____	I. - 132	Cagui	_____	I. 222, 224
Behemoth	_____	I. - 157	Caitaia	_____	I. - 220
Bekker el Wash	_____	I. - 102	Callitriche	_____	I. - 203
Belette	_____	II. 33	Callitrichus	_____	I. - 203
Belier	_____	I. - 40	Camelus	_____	I. - 132
Beluga	_____	II. 301	Camelo pardalis	_____	I. - 65
Bey	_____	I. - 212	Campagnol	_____	II. 205
Biber	_____	II. 114	Cani-apro-lupo-vulpes	_____	I. - 272
Bievre	_____	II. 114	Capefch	_____	I. - 265
Biggel	_____	I. - 83	Capibara	_____	II. 88
Biorn	_____	II. 1	Caprea	_____	I. 96, 120
Bifon d'Amerique	_____	I. - 23	Capreolus	_____	I. - 120
Biur	_____	II. 114	Capricorne	_____	I. - 55
Blaireau	_____	II. 14	Caracal	_____	I. - 305
Bobak	_____	II. 131	Caraco Mus	_____	II. 181
Bobr	_____	II. 114	Carcajou	_____	II. 8
Bocht	_____	I. - 132	Cariacou	_____	I. - 122
Boern-doeckie	_____	II. 157	Caribou	_____	I. - 111
Bœuf	_____	I. - 16	Carigueya	_____	II. 20
Bonnet Chinois	_____	I. - 209	Cariguibeiu	_____	II. 79
Bofchratte	_____	II. 90	Caftor	_____	II. 114
Bœzard	_____	I. - 102	Cataphractus	_____	II. 246
Bouc	_____	I. - 60	Cavia, cobaya	_____	II. 80
d' Angora	_____	I. - 61	Cavia, genus	_____	II. 88
d' Afrique	_____	I. - 63	Cay	_____	I. - 219
d' Juda	_____	I. - 63	Cayopollin	_____	II. 24
Baumrüttler	_____	I. - 295	Cemas	_____	I. - 85
Bouquetin	_____	I. - 55	Cerf	_____	I. - 114
Boury	_____	I. - 28	Chacal	_____	I. - 261
Buz argus	_____	II. 240	Chameau	_____	I. - 132
Bradypus	_____	I. - 116	Chamois	_____	I. - 44
Brandhirtz	_____		Chat	_____	I. - 295

INDEX OF CLASSICAL NAMES, &c.

	Vol.	Page		Vol.	Page
Chat d' Angora ————	I.	- 296	Cuetclatchtli ————	I.	- 250
d' Espagne ————	I.	- 296	Cuguaca-apara ————	I.	- 122
Chat-pard ————	I.	- 300	Cugacuara ————	I.	- 289
Chaus <i>Pinnii</i> ————	I.	- 301	Cuguacu ete ————	I.	- 126
Chauve-fouris ————	II.	- 311	Cugacuarana ————	I.	- 289
Chérôpotamus ————	I.	- 156	Culpeu ————	I.	- 258
Chérofo ————	II.	- 186	Cuniculus ————	II.	- 173
Cheval ————	I.	- 1	Cynocephalus ————	I.	- 186
Chevre ————	I.	- 60			
Chevreuil ————	I.	- 120	D		
Chevrotain d'Afrique ————	I.	- 81	Dachs ————	II.	- 14
de Guinée ————	I.	- 82	Dain ————	I.	- 113
des Indes ————	I.	- 127	Dama, Antelope ————	I.	- 85
Chien ————	I.	- 235	Daman Israél ————	II.	- 92
Chimpanzee ————	I.	- 150	Dam-tanhirfelch ————	I.	- 113
Chinchê ————	II.	- 65	Dandoelana ————	II.	- 140
Chinchimen ————	II.	- 82	Daniel ————	I.	- 113
Chinchilla ————	II.	- 196	Dant ————	I.	- 21
Chomik ————	II.	- 206	Defman * ————	II.	- 221
Choras ————	I.	- 188	Dafypus ————	II.	- 246
Cirquingon ————	II.	- 250	Djammel ————	I.	- 129
Citillus ————	II.	- 135	Diane ————	I.	- 201
Civetle ————	II.	- 70	Dof, Dof-hiort ————	I.	- 113
Clap-Myfs ————	II.	- 279	Dorcas ————	I. 92,	102, 120
Coaita ————	I.	- 216	Douc ————	I.	- 211
Coafe ————	II.	- 62	Dromedaire ————	I.	- 129
Coati ————	II.	- 12, 61	Dferen ————	I.	- 96
Cochon ————	I.	- 140	Dhikketaci ————	I.	- 4
Cochon d'Inde ————	II.	- 89	Dubha ————	I.	- 270
Coendu ————	II.	- 125	Dugon ————	II.	- 269
Coymatl ————	I.	- 147	Dzik ————	I.	- 140
Colus ————	I.	- 98	E		
Condoma ————	I.	- 88	Echinus Terrestris ————	II.	- 234
Conepate ————	II.	- 64	Ecureuil ————	II.	- 138
Coquallin ————	II.	- 147	Eichhorn ————	II.	- 138
Corine ————	I.	- 101	Elan ————	I.	- 105
Coudous ————	I.	- 78	Elephant, Elephas ————	I.	- 165
Couguar ————	I.	- 289, 290	Elk ————	I.	- 105
Coyotl ————	I.	- 257	Encourbert ————	II.	- 247
Coypu ————	II.	- 177	Engalla ————	I.	- 144
Crabier ————	II.	- 24	Erdzeifl ————	II.	- 205
Cricetus ————	II.	- 206	Erinaceus ————	II.	- 234, 248
Crocota ————	I.	- 272	Exquima ————	I.	- 201
Cuandu ————	II.	- 124			

* *De Buffon*, x. I. tab. II.

U V 2

Fara

INDEX OF CLASSICAL NAMES, &c.

	Vol.	Page		Vol.	Page
F					
Fara	II.	18	Guerlinguets	II.	162
Felis Catus	I.	295	Guevei	I.	82
Feuille	II.	309	Guib	I.	81
Fial racka	I.	255	Guillino	II.	120
Fiber	II.	114	Gulo	II.	110
Filander	II.	22	Gundi	II.	137
Filfreis	II.	10		H	
Fiskatta	II.	65	Hamster	II.	206
Fistal	I.	53	Handl	I.	150
Flader-mus	II.	319	Hardlooper	I.	144
Fong kyo fo	I.	133	Hafe	II.	98
Fossane	II.	75	Hæbe	II.	234
Fouine	II.	41	Heriflon	II.	234
Fourmillier	II.	260	Hermine	II.	35
Foyna	II.	41	Hippelaphus	I.	115
Fret	II.	40	Hippopotamus	I.	157
Fuchs	I.	251	Hirax	II.	92
Furo	II.	40	Hirco cervus	I.	52
G					
Ganfud	II.	234	Hiori	I.	114
Galera	II.	53	Hirsch	I.	114
Galgopithecus	I.	234	Hoang yang	I.	96
Gazelle	I.	75, 77, 89	Huanucu-Llama	I.	133
Gemse	I.	72	Hugiun	I.	129
Genette	II.	74	Hyæna	I.	270
Gerbo	II.	170	Hydrochærus	I.	88
Ghainouk	I.	27	Hyfrix	II.	122
Gibbon	I.	184	I		
Giraffe	I.	65	Jæerven	II.	10
Glis	II.	158	Jaguar	I.	284, 286
Glutton	II.	10	Jaguarete	I.	290
Glouton	II.	8	Jagura	I.	286
Gnou	I.	70	Jarf	II.	10
Gornostay	II.	35	Javaris	I.	147
Grafkin	II.	138	Ibex	I.	55
Grimme	I.	81	Ichneumon	II.	54
Grifon	II.	52	Jelen	I.	114
Guachi	II.	80	Jez	II.	234
Guanaco	I.	138	Jerboa	II.	164
Guanque	II.	183	Igel, Igelkott	II.	234
Guareba	I.	214	Ignavus	II.	240
Guepard *	I.	284	Indri	I.	228
			Jocko	I.	180

* *De Buffon* viii, 254. The same with the Hunting Leopard, No. 184.

INDEX OF CLASSICAL NAMES, &c.

	Vol.	Page		Vol.	Page
Ἰάβυβος	I.	157	Lerot	II.	159
Irabubos	II.	88	Lerwee	I.	53
Itatis	I.	255	Leucoryx	I.	76
			Levrier	I.	241
			Lidmee	I.	91
K			Lievre	II.	98
Kabarga	I.	174	Lion	I.	274
Kabafiou	II.	249	Loir	II.	158
Kalan	II.	83	Loris	I.	228
Kangaru	II.	29	Loup	I.	248
Kanin	II.	103	de Mexique	I.	250
Karagan	I.	252	Loup-Cervier	I.	301
Kassigiak	II.	270	Loup-Renard	I.	257
Kattlo	I.	301	Loutre	II.	77
Κατῶ	I.	210	Lowe	I.	274
Kenlie	I.	265	Lupus	I.	248
Kevel	I.	92	Lummick	II.	198
Kidang	I.	119	Lutra	II.	77
Kinkajou	II.	60	Lux	I.	301
Kob	I.	104	λυξ	I.	301
Koba	I.	103	Lynx	I.	301
Kolonnok	II.	39			
Kot dr.ki	I.	205			
Koulau	I.	8			
Kret	II.	229	M		
Kron-hiort	I.	114	Maucauco	I.	227
Krietsch	II.	206	Machlis	I.	105
Krylatca	II.	279	Mafutiliqui	II.	66
Kuna	II.	41	Magot	I.	186
			Magu	I.	213
L			Maimon	I.	190
Llama	I.	133	Malbrouck*	I.	201
Lacertus	II.	252	Mammouth	I.	172
Laderlap	II.	319	Manati	II.	292
Lame	II.	285	Mandrill	I.	190
Lant	I.	21	Mangabey	I.	204
Lapin	II.	103	Mangouffe	II.	54
d' Angora	II.	104	Manicou	II.	18
Lar	I.	185	Manipouris	I.	163
Latax	II.	80	Manis	II.	252
Lemur	I.	227	Manul	I.	294
Lemmar, Leming	II.	198	Mapach	II.	12
Lemni	II.	214	Maraguo	I.	292
Leo	I.	274	Mard	II.	41
Leopard	I.	282	Margay	I.	292
Lepus	II.	98	Marikina	I.	225
			Mariputa	II.	66

* *De Buffon*, xiv. 224. tab. xxix. A variety of our *Egret*, No. 119.

Marmose

INDEX OF CLASSICAL NAMES, &c.

	Vol.	Page		Vol.	Page
Marmose	II.	23	Nanguer	I.	85
Marmotte	II.	128	Neidek	II.	278
Martes, Marte	II.	41, 42	Nems	II.	54
Mejangan Banjoe	I.	118	Niedzwiadz	II.	1
Meles	II.	14	Nietferfoak	II.	279
Meminna	I.	127	Nil-ghau	I.	83
Mico	I.	226	Noctule	II.	317
Mocawk	I.	230	Noerza	II.	80
Mococo	I.	230	Norka	II.	80
Molle	II.	205	Nuxtegis	II.	319
Monax	II.	130			
Mone	I.	210		O	
Monèa	I.	207	Ocelot	I.	287
Mongooz	I.	229	Ochis	I.	16
Moofe	I.	107	Odobenus	II.	266
Morfe	II.	266	Ogotona	II.	109
Morkuia Korawa	II.	292	Onager	I.	11
Mouffettes *	II.	62	Once	I.	285, 290
Mouffon	I.	44	Ondatra	II.	119
Mouffac	I.	205	Opeagha	I.	14
Mouton de Barbarie	I.	41	Ophion	I.	44
Mufro	I.	46	Orang Outang	I.	180
Mulet	I.	8	Oreillar	II.	320
Muzza	II.	224	Oreotragus	I.	79
Mullvad	II.	229	Original. See Elk.		
Mulet	II.	184	Oryx	I.	76
Munt-jak	I.	119	Oitrowidz	I.	301
Murmelthier	II.	128	Ouaikarè	II.	240
Mus Alpinus	II.	128	Ouanderou	I.	198
Mus	II.	134	Ouarine	I.	214
Mufaraigne	II.	224	Cuikiti	I.	224
Mus Araneus	II.	224	Ourebi	I.	79
Musc	I.	124	Ourico	II.	124
Muscardin	II.	160	Ours	II.	1
Mufimon	I.	44	Ours blanc de mer	II.	5
Mufquash	II.	119	Ours marin	II.	281
Muffafcus	II.	119	Ovis	I.	37
Mutela	II.	33		P	
Myrmecophaga	II.	256	Paca	II.	91
	N		Pacaffe	I.	78
Nabbmus	II.	224	Paco, Pacos	I.	136
Nagor	I.	86	Palatine	I.	200

* *M. De Buffon's* generic name for the Polecats which exhale fo putridential a vapour.

Palmiste

INDEX OF CLASSICAL NAMES, &c.

	Vol.	Page		Vol.	Page
Palmiste	—	II. 149	Quahatechalotl-thlitic	—	II. 145
Pangolin	—	II. 253	Quapizotl	—	I. - 147
Panthera, Panthere	—	I. - 280	Quato	—	I. - 216
Papio	—	I. - 188	Quauhila	—	I. - 147
Παρδαλις	—	I. 280, 285	Quil, Quirpele	—	II. 54
Pardus	—	I. - 280	Quojas Morrou	—	I. - 180
Pareffieux	—	II. 240	Quoll	—	II. 69
Pafan, Pafan	—	I. 57-75	Quouata	—	I. - 216
Patas	—	I. - 208	Quumbengo	—	I. - 272
Pecary	—	I. - 147			
Pekan	—	II. 51		R	
Pelander Aroe	—	II. 21	Radjur	—	I. - 120
Perchal	—	II. 179	Racé	—	I. - 251
Pere	—	I. - 12	Rangier	—	I. - 111
Perugufna	—	II. 38	Rangwo	—	II. 105
Petit Gris	—	II. 144	Rat	—	II. 176
Phalanger	—	II. 27	d'Eau	—	II. 182
Phatagin	—	II. 252	de Madagafcar	—	I. - 233
Philandre	—	II. 27	Ratel	—	II. 66
Philodotus	—	II. 252	Raton	—	II. 12
Phoca	—	II. 270	Rattepenade	—	II. 320
Phoque	—	II. 270	Rein Deer	—	I. 111
Pichou	—	I. - 292	Renard	—	I. - 251
Piloris	—	II. 97	Renne	—	I. - 111
Pilofello	—	II. 74	Rennthier	—	I. - 111
Pinche	—	I. - 25	Rhen	—	I. - 111
Pipitrelle	—	II. 318	Rhinoceros	—	I. - 150
Piffay	—	I. - 127	Rillow	—	I. - 209
Πιθνε, Pitheque	—	I. - 183	River Paard	—	I. - 157
Platogna	—	I. - 113	Roloway	—	I. - 200
Platyceros	—	I. - 113	Root	—	I. - 213
Poephagus	—	I. - 27	Rofelet	—	II. 35
Poulatouche	—	II. 155	Rofimarus	—	II. 266
Pongo	—	I. - 180	Rofomak	—	II. 10
Porc-epic	—	II. 122	Rougette	—	II. 304
Πρωξ	—	I. - 113	Rouffette	—	II. 304
Przewiafska	—	II. 38	Rukkaia	—	II. 140
Pteropus	—	II. 304, 308	Rupicapra	—	I. 44, 72
Puma	—	I. - 289	Rulla	—	II. 112
Putois	—	II. 37	Rys	—	I. - 301
Putorius	—	II. 37		S	
	Q		Saca	—	I. - 294
Quagga	—	I. - 14	Saccawinkee	—	I. - 222
			Saguin	—	Saguin

INDEX OF CLASSICAL NAMES, &c.

	Vol.	Page		Vol.	Page
Sagouin	I.	244	Squiflachi	I.	261
Sai	I.	218	Squinaton	I.	122
Saiga	I.	98	Steinbock	I.	55
Salmiri	I.	220	Stink bingfem	II.	66
Sajou	I.	217	Stock	I.	150
Saki	I.	222	Strepsiceros	I.	38, 88
Sangler	I.	140	Suhak	I.	98
de Capvert	I.	144	Suisse, Ecureuil	II.	157
Sanglin	I.	224	Sumxi	I.	296
Sapajou	I.	222	Surikate	II.	57
Saragoy	II.	20	Surk	II.	229
Saricovienne	II.	82	Surmulot	II.	178
Sarigue	II.	20	Sus	I.	140
Sarlyk	I.	27	Sus Aquaticus	I.	163
Schakal	II.	45	Swiftch	II.	131
Satyrus	I.	180			
Scenoontung	I.	122	T		
Schakal	I.	261	Taguan	II.	151
Schwein	I.	140	Tajacu	I.	147
Sciurus	II.	138	Tajibi	II.	18
Semlanoi Sactshik	II.	112	Taifon	II.	14
Serotine	II.	317	Talapoin	I.	206
Serval	I.	301	Talpa	II.	229
Shitnik	II.	190	Tamandua	II.	256
Sial	II.	270	Tamanoir	II.	256
Siegen Bock	I.	60	Tamarin	I.	223
Sifac	I.	211	Tanrec	II.	236
Simia	I.	183	Tapeti	II.	107
Siwutscha	II.	288	Tapir	I.	163
Siya	II.	79	Tapoa Tafa	II.	69
Siyah Ghusha	I.	305	Tarandus	I.	111
Skrzeczek	II.	206	Tardigradus	II.	240
Siepez	II.	214	Tarfier	I.	231
Sno-mus	II.	33	Tartarin	I.	194
Sobol	II.	47	Tatou	II.	248
Sogur	II.	131	Tatu apara	II.	246
Songar	II.	212	Tatuete	II.	248
Sorex	II.	224	Taupe	II.	229
Souris	II.	184	Taupe dorée	II.	231
Souflik	II.	135	Taureau	I.	16
Speck-maus	II.	319	Taxus	I.	270
Spring-bock	I.	64	Tayra	II.	53
Springen Haas	II.	170	Tchoriz	II.	37
			Tegoulitchek	II.	194
			Tegul		

INDEX OF CLASSICAL NAMES, &c.

	Vol.	Page		Vol.	Page
Tegul			Ulf	I.	248
Temamaçama	I.	103	Unau	II.	242
Tendrac	II.	236	Uncia	I.	282
Tenlie	I.	265	Vormela	II.	210
Tepe Maxlaton	I.	292	Urigne	II.	290
Teutlalmaçama	I.	122	Urson	II.	126
Tgao	I.	157	Urfus	II.	1
Thous	I.	267	Urus	I.	16
Tigris, Tigre	I.	277	Υς τερπνεύς	I.	148
Tlaloceloti	I.	287	Υγις	II.	122
Tla-coozelotl	I.	287	Vulpes	I.	251
Tlalmorotli	II.	149	Ufuk	II.	277
Tlaquatzin	II.	18. 124	Utter	II.	77
Tolai	II.	104			
Touan	II.	34		W	
Tragelaphus	I.	52	Walrus	II.	266
Trago-Camelus	I.	83	Warg	I.	248
Tragulus	I.	124	Warglo	I.	301
Tretretrete	I.	191	Warree	I.	141
Trichechus Rosmarus	II.	266	Weefel	II.	33
Manatus	II.	297	Wettfk	I.	150
Tŷchotŷchot	II.		Whang, Yang	I.	96
Tŷtsjan	II.	135	Wha Tapoua Row	II.	13
Tucan	II.	223	Wiewiorka	II.	138
Tzeiran	I.	74	Wirrebocarra	I.	126
			Whydra	II.	77
V				X	
Vache Marine	II.	266	Χοιρωιθνας	I.	187
de Tartarie	I.	24	Xoloitzcuintli	I.	250
Yalaz	I.	270		Y	
Vampire	II.	308	Yerboa	II.	29
Vanfire	II.	51	Yltis	II.	37
Vary	I.	229	Yfard, Yfarus	I.	72
Varia	I.	280	Yzquiepatl	II.	62
Vavi	I.	261		Z	
Veldratte	II.	90	Zbik	I.	295
Verdadeiro	II.	248	Zebre, Zebra	I.	13
Vespertilio	II.	304	Zebu	I.	28
Vigogne, Vicunna	I.	136		Zecora	
Vilon	II.	51			
Viverra	II.	40			
Viverra tigrina	I.	298			
Vol. II.			X x		

INDEX OF CLASSICAL NAMES, &c.


Zecora	_____	Vol. Page	Zibet	_____	Vol. Page
Zenik	_____	I. - 13	Zizel	_____	II. 72
Zerda	_____	II. 193	Zobela	_____	II. 135
Zibeline	_____	I. - 267	Zorrina*	_____	II. 43
		II. 43			II. 66

* *De Buffon*, xiii. 302. tab. xli.

THE END.


ERRATUM.

Nos. 273 and 276, Pages 70 and 72, being the same animal, the reader is desired to correct this mistake.













1000073430

NOT FOR CIRCULATION

